

27
1-22-76
to DTIS

MASTER

ORNL-TM-5204
ENDF-228

**SB3. Experiment on Secondary Gamma-Ray
Production Cross Sections Averaged Over
a Fast-Neutron Spectrum for Each
of 13 Different Elements Plus
a Stainless Steel**

R. E. Maerker



OAK RIDGE NATIONAL LABORATORY

OPERATED BY UNION CARBIDE CORPORATION • FOR THE U.S. ATOMIC ENERGY COMMISSION

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency Thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

Printed in the United States of America. Available from
National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road, Springfield, Virginia 22161
Price: Printed Copy \$5.00; Microfiche \$2.25

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the Energy Research and Development Administration, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

Contract No. W-7405-eng-26


Neutron Physics Division

**SB3. Experiment on Secondary Gamma-Ray Production Cross
Sections Averaged Over a Fast-Neutron Spectrum for
Each of 13 Different Elements Plus a Stainless Steel**

R. E. Maerker

Reference: R. E. Maerker and F. J. Muckenthaler, "Gamma-Ray Spectra
Arising from Fast-Neutron Interactions in Elements Found in
Soil, Concretes, and Structural Materials", ORNL-4475 (1969).

JANUARY 1976



OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37830
operated by
Union Carbide Corporation
for the
U.S. ENERGY RESEARCH AND DEVELOPMENT ADMINISTRATION

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED
fy

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

Abstract

The experimental and calculational details for a CSEWG integral data testing shielding experiment are presented. This particular experiment measured the secondary gamma-ray production cross sections averaged over a fast-neutron spectrum for iron, oxygen, sodium, aluminum, copper, titanium, calcium, potassium, silicon, nickel, zinc, barium, sulfur, and a type 321 stainless steel. The gamma-ray production cross sections were binned into ~ 0.5 -MeV wide gamma-ray energy intervals.

Description

Gamma-ray production cross sections averaged over a fast-neutron (> 1 MeV) spectrum for iron, stainless steel, oxygen, and sodium, obtained in 0.5-MeV bins from gamma-ray spectra measurements described in the reference, were originally collated and documented in SDT7*. The present document supplements these results with additional data on aluminum, copper, titanium, calcium, potassium, silicon, nickel, zinc, barium, and sulfur. All the measurements were made at the Oak Ridge National Laboratory Tower Shielding Facility using a carefully calibrated 5 x 5 in. NaI(Tl) detector in good geometry. The experimental arrangement is shown in Fig. 1.

Experimental Data

The experimental data are differential in the gamma-ray energy from approximately 1 to 6.5 MeV and are expressed as values of 4π times the differential gamma-ray production cross section in millibarns at 90 degrees to the incident neutron beam averaged over a neutron spectrum lying above 1 MeV. Gamma-ray production measurements for all 13 elements plus the stainless steel are presented in Tables 1A through 15A. These results are estimated to have an accuracy of $\pm 30\%$ and include the contributions from both discrete and continuum gamma rays. The tabulated relative fast-neutron spectra for each experiment are given in Tables 1B through 3B.

Methods of Calculation

No transport calculations are necessary for this benchmark, hence model description, atom densities, etc. are not needed. Averaging can be done as follows:

Let $P(\Delta E_n)$ be the entries in the tabulated relative fast-neutron spectrum. Then for the continuum or unresolved contribution to the

* SB3 supercedes SDT7 in the new shielding benchmark series (1976).




Fig. 1. Schematic Diagram of Geometry for Determining Average Secondary Gamma-Ray Production Cross Sections from Fast-Neutron Interactions. In addition, a 6-in. thickness of lithium hydride, not shown, was placed in the detector beam between the slab and the detector collimator.

production cross section,

$$\bar{\sigma}_c(\Delta E\gamma) = \sum_{\Delta E_n} P(\Delta E_n) \bar{\sigma}_c(\Delta E_n, \Delta E\gamma) ,$$

where

$$\bar{\sigma}_c(\Delta E_n, \Delta E\gamma) = \int_{\Delta E_n} \sigma_c(E_n, \Delta E\gamma) dE_n / \Delta E_n .$$

The evaluation of the integral depends on the neutron energy grid of the ENDF/B data, and may have to be rather coarsely estimated.

Similarly, for the discrete contribution to the production cross section,

$$\bar{\sigma}_d(E\gamma) = \sum_{\Delta E_n} P(\Delta E_n) \bar{\sigma}_d(\Delta E_n, E\gamma) ,$$

where

$$\bar{\sigma}_d(\Delta E_n, E\gamma) = \int_{\Delta E_n} \sigma_d(E_n, E\gamma) dE_n / \Delta E_n ,$$

and again the accuracy of the evaluation of this integral depends on the neutron energy grid of the data.

Finally,

$$\bar{\sigma}(\Delta E\gamma) = \bar{\sigma}_c(\Delta E\gamma) + \sum_{\Delta E\gamma} \bar{\sigma}_d(E\gamma) ,$$

and it is this value that is to be compared with experiment.

It is recommended, however, that a "standard" ENDF/B photon production group averaging code such as LAPHANO (written by Los Alamos) or LAPHFOR (Oak Ridge's modified version of LAPHANO) be used to accomplish the above.

The input for LAPHANO or LAPHFOR, for example, is relatively straightforward. Note the following input items:

NGG = number of gamma groups for each element is determined from
Tables 1A - 15A;

NGF = number of neutron fine groups is determined from Tables 1B -
3B plus one for a sink group;

NBF = number of neutron broad groups = 1.

The gamma energy bounds are given in Tables 1A - 15A and the neutron energy bounds and weighting spectrum are given in Tables 1B - 3B. The neutron broad group is specified from thermal energy to 14 MeV, which covers the energy region of interest. The sink group, thermal to 1 MeV, has a negligibly small weight, say 10^{-8} . The MT flag is set to call for all reactions given in the files. All other flags for numbers of nuclides, mixtures, zones, etc. are set for one material, one mixture, one zone with radius = 1.0 etc. Note that the cross sections are outputted in barns.*

Method of Reporting

The calculational results should be tabulated as shown in Tables 16 through 29.

*Acknowledgments should be made to R. J. LaBauve of Los Alamos Scientific Laboratory and W. E. Ford III of Oak Ridge National Laboratory for the general description of the input for the LAPHANO and LAPHFOR processing codes.

Table 1

- A. Experimental values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum for iron are the following, in millibarns:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.0 - 1.5	278
1.5 - 2.0	132
2.0 - 2.5	101
2.5 - 3.0	74
3.0 - 3.5	44
3.5 - 4.0	33
4.0 - 4.5	8.7
4.5 - 5.0	5.6
5.0 - 5.5	3.8
5.5 - 6.0	2.4
6.0 - 6.5	< 2.1
6.5 - 7.0	< 0.6
7.0 - 7.5	< 1.5
≥ 7.5	0

*Upper energy limit of each group not included in summation.

- B. Tabulated relative fast-neutron spectrum:

Neutron Energy Group (MeV)	Relative Number in Group	Neutron Energy Group (MeV)	Relative Number in Group
1.0 - 1.5	0.174	5.5 - 6.0	0.026
1.5 - 2.0	0.163	6.0 - 6.5	0.021
2.0 - 2.5	0.169	6.5 - 7.0	0.015
2.5 - 3.0	0.118	7.0 - 7.5	0.011
3.0 - 3.5	0.079	7.5 - 8.0	0.008
3.5 - 4.0	0.063	8.0 - 9.0	0.009
4.0 - 4.5	0.056	9.0 - 10.0	0.005
4.5 - 5.0	0.045	10.0 - 12.0	0.003
5.0 - 5.5	0.034	12.0 - 14.0	0.001

Totals 1.0 - 14.0

1.000

Table 2

- A. Experimental value of 4π times the differential gamma-ray production cross section at 90° averaged over the tabulated fast-neutron spectrum for the 6.13-MeV gamma ray in oxygen is the following, in millibarns:

<u>Gamma-Ray Energy</u> (MeV)	<u>Average Cross Section</u> (mb)
6.13	90

- B. Tabulated relative fast-neutron spectrum for the 6.13-MeV gamma-ray production test:

<u>Neutron Energy</u> <u>Group (MeV)</u>	<u>Relative Number</u> <u>in Group</u>	<u>Neutron Energy</u> <u>Group (MeV)</u>	<u>Relative Number</u> <u>in Group</u>
6.5 - 7.0	0.290	10.5 - 11.0	0.018
7.0 - 7.5	0.204	11.0 - 11.5	0.013
7.5 - 8.0	0.150	11.5 - 12.0	0.009
8.0 - 8.5	0.107	12.0 - 12.5	0.006
8.5 - 9.0	0.075	12.5 - 13.0	0.004
9.0 - 9.5	0.054	13.0 - 13.5	0.003
9.5 - 10.0	0.038	13.5 - 14.0	0.002
<u>10.0 - 10.5</u>	0.027		
Totals 6.5 - 14.0			1.000

Table 3

- A. Experimental value of 4π times the differential gamma-ray production cross section at 90° averaged over the tabulated fast-neutron spectrum for the 6.92 + 7.12 MeV gamma rays in oxygen is the following, in millibarns:

<u>Gamma-Ray Energy</u> (MeV)	<u>Average Cross Section</u> (mb)
6.92 + 7.12	55

- B. Tabulated relative fast-neutron spectrum for the production test for 6.92 + 7.12 MeV gamma rays:

<u>Neutron Energy</u> <u>Group (MeV)</u>	<u>Relative Number</u> <u>in Group</u>	<u>Neutron Energy</u> <u>Group (MeV)</u>	<u>Relative Number</u> <u>in Group</u>
7.4 - 7.5	0.063	10.5 - 11.0	0.034
7.5 - 8.0	0.278	11.0 - 11.5	0.024
8.0 - 8.5	0.198	11.5 - 12.0	0.016
8.5 - 9.0	0.139	12.0 - 12.5	0.012
9.0 - 9.5	0.099	12.5 - 13.0	0.008
9.5 - 10.0	0.069	13.0 - 13.5	0.006
<u>10.0 - 10.5</u>	0.050	13.5 - 14.0	<u>0.004</u>
Totals 7.4 - 14.0			1.000

Table 4

- A. Experimental values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for stainless steel (67.4 weight percent iron, 18.3 weight percent chromium, 9.7 weight percent nickel, 1.5 weight percent manganese, and the remainder may be neglected) are the following, in millibarns:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.0 - 1.5	383
1.5 - 2.0	139
2.0 - 2.5	84
2.5 - 3.0	53
3.0 - 3.5	43
3.5 - 4.0	25
4.0 - 4.5	8.8
4.5 - 5.0	5.8
5.0 - 5.5	5.1
5.5 - 6.0	3.0
6.0 - 6.5	< 2.6
6.5 - 7.5	< 4.9
≥ 7.5	0

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 5

- A. Experimental values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum[†] for sodium are the following, in millibarns:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.1 - 1.5	59
1.5 - 2.0	131
2.0 - 2.5	47
2.5 - 3.0	50
3.0 - 3.5	22
3.5 - 4.0	15
4.0 - 4.5	7.3
4.5 - 5.0	4.6
5.0 - 5.5	2.9
5.5 - 6.0	3.7
6.0 - 6.5	4.2
6.5 - 7.5	2.4
≥ 7.5	0

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 6

- A. Experimental values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for aluminum are the following, in millibarns:

<u>Gamma-Ray Energy Interval</u> (MeV)*	<u>Average Cross Section</u> (mb)
1.0 - 1.5	181
1.5 - 2.0	43.6
2.0 - 2.5	102.5
2.5 - 3.0	26.3
3.0 - 3.5	42.4
3.5 - 4.0	8.8
4.0 - 4.5	5.8
4.5 - 5.0	6.9
5.0 - 5.5	4.9
5.5 - 6.0	2.0
6.0 - 6.5	< 1.7
6.5 - 7.0	< 1.5
7.0 - 7.5	< 0.3

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 7

A. Experimental values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for copper are the following, in millibarns:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
0.95 - 1.5	1090
1.5 - 2.0	307
2.0 - 2.5	127
2.5 - 3.0	87
3.0 - 3.5	45
3.5 - 4.0	35
4.0 - 4.5	24
4.5 - 5.0	13.5
5.0 - 5.5	9.1
5.5 - 6.0	< 5.9
6.0 - 6.5	< 6.2

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 8

A. Experimental values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for titanium are the following, in millibarns:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
0.98 - 1.5	688
1.5 - 2.0	139
2.0 - 2.5	97
2.5 - 3.0	48
3.0 - 3.5	22
3.5 - 4.0	20
4.0 - 4.5	9.5
4.5 - 5.0	5.1
5.0 - 5.5	< 4.4
5.5 - 6.0	< 2.8
6.0 - 6.5	< 2.4

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 9

A. Experimental values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for calcium are the following, in millibarns:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.1 - 1.5	64
1.5 - 2.0	44
2.0 - 2.5	18
2.5 - 3.0	16.5
3.0 - 3.5	8.0
3.5 - 4.0	40.5
4.0 - 4.5	2.5
4.5 - 5.0	1.9
5.0 - 5.5	1.6
5.5 - 6.0	< 3.0
6.0 - 6.5	< 1.4
6.5 - 7.0	< 1.1

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 10

A. Experimental values of 4π times the differential gamma-ray production cross-sections at 90° averaged over the tabulated fast-neutron spectrum† for potassium are the following, in millibarns:

<u>Gamma-Ray Energy Interval</u> <u>(MeV)*</u>	<u>Average Cross Section</u> <u>(mb)</u>
1.1 - 1.5	133
1.5 - 2.0	96
2.0 - 2.5	39
2.5 - 3.0	96
3.0 - 3.5	26
3.5 - 4.0	23
4.0 - 4.5	6.1
4.5 - 5.0	8.7
5.0 - 5.5	3.6
5.5 - 6.0	4.6
6.0 - 6.5	< 3.1
6.5 - 7.0	< 1.4
7.0 - 7.5	< 1.5

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 11

A. Experimental values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for silicon are the following, in millibarns:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.1 - 1.5	55
1.5 - 2.0	344
2.0 - 2.5	30
2.5 - 3.0	11
3.0 - 3.5	5.4
3.5 - 4.0	5.4
4.0 - 4.5	1.2
4.5 - 5.0	3.2
5.0 - 5.5	1.2
5.5 - 6.0	1.5
6.0 - 6.5	< 1.5
6.5 - 7.0	< 0.7
7.0 - 7.5	< 2.6

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 12

- A. Experimental values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for nickel are the following, in millibarns:

<u>Gamma-Ray Energy Interval (MeV)*</u>	<u>Average Cross Section (mb)</u>
1.0 - 1.5	790
1.5 - 2.0	179
2.0 - 2.5	68
2.5 - 3.0	43
3.0 - 3.5	37
3.5 - 4.0	17
4.0 - 4.5	11.5
4.5 - 5.0	4.8
5.0 - 5.5	< 4.7
5.5 - 6.0	< 3.4
6.0 - 7.0	< 5.3

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 13

- A. Experimental values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for zinc are the following, in millibarns:

<u>Gamma-Ray Energy Interval</u> (MeV)*	<u>Average Cross Section</u> (mb)
1.0 - 1.5	909
1.5 - 2.0	234
2.0 - 2.5	113
2.5 - 3.0	59
3.0 - 3.5	41
3.5 - 4.0	27
4.0 - 4.5	16.1
4.5 - 5.0	< 13.3
5.0 - 5.5	< 6.4
5.5 - 6.0	< 5.6

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 14

- A. Experimental values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for barium are the following, in millibarns:

<u>Gamma-Ray Energy Interval (MeV)*</u>	<u>Average Cross Section (mb)</u>
1.1 - 1.5	1340
1.5 - 2.0	552
2.0 - 2.5	290
2.5 - 3.0	144
3.0 - 3.5	79
3.5 - 4.0	47
4.0 - 4.5	22.3
4.5 - 5.0	21.0
5.0 - 5.5	< 9.2
5.5 - 6.0	< 8.3
6.0 - 6.5	< 5.2
6.5 - 7.0	< 2.4

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 15

- A. Experimental values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for sulfur are the following, in millibarns:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.1 - 1.5	73
1.5 - 2.0	44
2.0 - 2.5	199
2.5 - 3.0	16.4
3.0 - 3.5	11.2
3.5 - 4.0	3.3
4.0 - 4.5	10.8
4.5 - 5.0	3.8
5.0 - 5.5	1.2
5.5 - 6.0	< 3.0
6.0 - 6.5	< 0.7
6.5 - 7.0	< 1.2
7.0 - 7.5	< 0.6

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 16

Report sheet for values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for iron:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.0 - 1.5	_____
1.5 - 2.0	_____
2.0 - 2.5	_____
2.5 - 3.0	_____
3.0 - 3.5	_____
3.5 - 4.0	_____
4.0 - 4.5	_____
4.5 - 5.0	_____
5.0 - 5.5	_____
5.5 - 6.0	_____
6.0 - 6.5	_____
6.5 - 7.0	_____
7.0 - 7.5	_____
≥ 7.5	_____

*Do not include upper energy limit of each group in summation.

†Use Table 1B for the neutron spectrum.

Table 17

Report sheet for values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectra † for oxygen:

<u>Gamma-Ray Energy</u> (MeV)	<u>Average Cross Section</u> (mb)
6.13	_____
6.92 + 7.12	_____

†Use Table 2B for the neutron spectrum for the 6.13 MeV gamma ray and Table 3B for the neutron spectrum for the 6.92 + 7.12 MeV gamma rays.

Table 18

Report sheet for values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for stainless steel:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.0 - 1.5	_____
1.5 - 2.0	_____
2.0 - 2.5	_____
2.5 - 3.0	_____
3.0 - 3.5	_____
3.5 - 4.0	_____
4.0 - 4.5	_____
4.5 - 5.0	_____
5.0 - 5.5	_____
5.5 - 6.0	_____
6.0 - 6.5	_____
6.5 - 7.5	_____
≥ 7.5	_____

*Do not include upper energy limit of each group in summation.

†Use Table 1B for the neutron spectrum.

Table 19

Report sheet for values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for sodium:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.1 - 1.5	_____
1.5 - 2.0	_____
2.0 - 2.5	_____
2.5 - 3.0	_____
3.0 - 3.5	_____
3.5 - 4.0	_____
4.0 - 4.5	_____
4.5 - 5.0	_____
5.0 - 5.5	_____
5.5 - 6.0	_____
6.0 - 6.5	_____
6.5 - 7.5	_____
≥ 7.5	_____

*Do not include upper energy limit of each group in summation.

†Use Table 1B for the neutron spectrum.

Table 20

Report sheet for values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for aluminum:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.0 - 1.5	_____
1.5 - 2.0	_____
2.0 - 2.5	_____
2.5 - 3.0	_____
3.0 - 3.5	_____
3.5 - 4.0	_____
4.0 - 4.5	_____
4.5 - 5.0	_____
5.0 - 5.5	_____
5.5 - 6.0	_____
6.0 - 6.5	_____
6.5 - 7.0	_____
7.0 - 7.5	_____

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 21

Report sheet for values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for copper:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
0.95 - 1.5	_____
1.5 - 2.0	_____
2.0 - 2.5	_____
2.5 - 3.0	_____
3.0 - 3.5	_____
3.5 - 4.0	_____
4.0 - 4.5	_____
4.5 - 5.0	_____
5.0 - 5.5	_____
5.5 - 6.0	_____
6.0 - 6.5	_____

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 22

Report sheet for values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for titanium:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
0.98 - 1.5	_____
1.5 - 2.0	_____
2.0 - 2.5	_____
2.5 - 3.0	_____
3.0 - 3.5	_____
3.5 - 4.0	_____
4.0 - 4.5	_____
4.5 - 5.0	_____
5.0 - 5.5	_____
5.5 - 6.0	_____
6.0 - 6.5	_____

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 23

Report sheet for values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for calcium:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.1 - 1.5	_____
1.5 - 2.0	_____
2.0 - 2.5	_____
2.5 - 3.0	_____
3.0 - 3.5	_____
3.5 - 4.0	_____
4.0 - 4.5	_____
4.5 - 5.0	_____
5.0 - 5.5	_____
5.5 - 6.0	_____
6.0 - 6.5	_____
6.5 - 7.0	_____

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 24

Report sheet for values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for potassium:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.1 - 1.5	_____
1.5 - 2.0	_____
2.0 - 2.5	_____
2.5 - 3.0	_____
3.0 - 3.5	_____
3.5 - 4.0	_____
4.0 - 4.5	_____
4.5 - 5.0	_____
5.0 - 5.5	_____
5.5 - 6.0	_____
6.0 - 6.5	_____
6.5 - 7.0	_____
7.0 - 7.5	_____

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 25

Report sheet for values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for silicon:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.1 - 1.5	_____
1.5 - 2.0	_____
2.0 - 2.5	_____
2.5 - 3.0	_____
3.0 - 3.5	_____
3.5 - 4.0	_____
4.0 - 4.5	_____
4.5 - 5.0	_____
5.0 - 5.5	_____
5.5 - 6.0	_____
6.0 - 6.5	_____
6.5 - 7.0	_____
7.0 - 7.5	_____

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 26

Report sheet for values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for nickel:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.0 - 1.5	_____
1.5 - 2.0	_____
2.0 - 2.5	_____
2.5 - 3.0	_____
3.0 - 3.5	_____
3.5 - 4.0	_____
4.0 - 4.5	_____
4.5 - 5.0	_____
5.0 - 5.5	_____
5.5 - 6.0	_____
6.0 - 7.0	_____

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 27

Report sheet for values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for zinc:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.0 - 1.5	_____
1.5 - 2.0	_____
2.0 - 2.5	_____
2.5 - 3.0	_____
3.0 - 3.5	_____
3.5 - 4.0	_____
4.0 - 4.5	_____
4.5 - 5.0	_____
5.0 - 5.5	_____
5.5 - 6.0	_____

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 28

Report sheet for values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for barium:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.1 - 1.5	_____
1.5 - 2.0	_____
2.0 - 2.5	_____
2.5 - 3.0	_____
3.0 - 3.5	_____
3.5 - 4.0	_____
4.0 - 4.5	_____
4.5 - 5.0	_____
5.0 - 5.5	_____
5.5 - 6.0	_____
6.0 - 6.5	_____
6.5 - 7.0	_____

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

Table 29

Report sheet for values of 4π times the differential gamma-ray production cross sections at 90° averaged over the tabulated fast-neutron spectrum† for sulfur:

Gamma-Ray Energy Interval (MeV)*	Average Cross Section (mb)
1.1 - 1.5	_____
1.5 - 2.0	_____
2.0 - 2.5	_____
2.5 - 3.0	_____
3.0 - 3.5	_____
3.5 - 4.0	_____
4.0 - 4.5	_____
4.5 - 5.0	_____
5.0 - 5.5	_____
5.5 - 6.0	_____
6.0 - 6.5	_____
6.5 - 7.0	_____
7.0 - 7.5	_____

*Upper energy limit of each group not included in summation.

†Use Table 1B for the neutron spectrum.

THIS PAGE
WAS INTENTIONALLY
LEFT BLANK

INTERNAL DISTRIBUTION

- | | |
|-------------------------------|---|
| 1-3. L. S. Abbott | 21. D. K. Trubey |
| 4-8. C. E. Clifford | 22. C. R. Weisbin |
| 9. G. F. Flanagan | 23. G. E. Whitesides |
| 10. W. E. Ford III | 24. A. Zucker |
| 11. H. Goldstein (Consultant) | 25-44. RSIC |
| 12. W. O. Harms | 45. W. W. Havens (Consultant) |
| 13. R. E. Maerker | 46. A. F. Henry (Consultant) |
| 14. F. C. Maienschein | 47. P. F. Fox (Consultant) |
| 15. F. J. Muckenthaler | 48-49. Central Research Library |
| 16. F. R. Mynatt | 50. ORNL Y-12 Technical Library
Document Reference Section |
| 17. E. M. Oblow | 51-52. Laboratory Records Dept. |
| 18. F. G. Perey | 53. Laboratory Records ORNL RC |
| 19. R. W. Roussin | 54. ORNL Patent Office |
| 20. D. B. Trauger | |

EXTERNAL DISTRIBUTION

55. Martin Becker, Rensselaer Polytechnic Institute, Tibbits Ave., Troy, N. Y. 12181
56. N. Francis, Knolls Atomic Power Laboratory, Schenectady, N. Y. 12301
57. R. A. Grimesey, Idaho Nuclear Corp., P. O. Box 1845, Idaho Falls, Idaho 83401
58. W. H. Hannum, Div. of Reactor Research and Development, Safety, ERDA, Washington, D. C. 20545
59. P. B. Hemmig, Div. of Reactor Research and Development, Reactor Physics Branch, ERDA, Washington, D. C. 20545
60. L. Harris, Science Applications, Inc., P. O. Box 2351, La Jolla, California 92037
61. M. H. Kalos, Courant Institute of Mathematical Sciences, New York University, 251 Mercer Street, New York, N. Y. 10012
62. R. J. LaBauve, Los Alamos Scientific Laboratory, P. O. Box 1663, Los Alamos, New Mexico 87544
63. Leona Stewart, Los Alamos Scientific Laboratory, P. O. Box 1663, Los Alamos, New Mexico 87544
64. Duane Marr, WADCO, P. O. Box 1970, Richland, Washington 99352
65. R. J. Howerton, Lawrence Livermore Laboratory, P. O. Box 808, Livermore, California 94550
66. E. A. Straker, Science Applications, Inc., P. O. Box 2351, La Jolla, California 92037
- 67-86. Sol Pearlstein, NNCSC, Brookhaven National Laboratory, Upton, N. Y. 11973
87. E. M. Bohn, Argonne National Laboratory, 9700 Cass Avenue, Argonne, Illinois 60439
88. Dr. John Butler, Bldg. B21, United Kingdom Atomic Energy Authority, AEE Winfrith, Dorchester, Dorset, England
89. C. Devillers, DPRMA/SERMA, B. P. 2, CEN Saclay, 91190 Gif Sur Yvette, France

90. Dr. Carlo Ponti, CCR Curatom, 21027 ISPRA(Varese), Italy
91. Dr. G. Hehn, Institut fur Kernenergetik, Der Universitat,
7000 Stuttgart 80, Pfaffenwaldring 31, Germany
92. Tomas Lefvert, Research Institute of National Defense,
FOA 212, FACK, S-10450, Stockholm 80, Sweden
93. Makato Akanuma, Computing Center, Div. of Reactor Eng.,
Japan Atomic Energy Research Institute, Tokai Research
Establishment, Tokai-Mura, Naka-Gun, Ibaraki-Ken, Japan
- 94-120. Technical Information Center (TIC)
121. Research and Technical Support Division