Cross sections and covariance data through international collaboration

Andrej Trkov[1)], Roberto Capote[2)]

1) Jožef Stefan Institute, Ljubljana, Slovenia
2) IAEA, Nuclear Data Section, Vienna, Austria

Covariances

- Methodology:
  EMPIRE-MC + GANDR

- Evaluations:
  - Th-232
  - W isotopes (including ^{180}W)
  - Mn-55 (on-going)

^{232}Th – Current status

- MF1/MT451: Corrections & clarification
- MF1/MT458: added
- MF3/all reactions: small differences due to refinements in GANDR
- MF32/MT151: setting ISR=0
- MF33/MT2: fully correlated uncertainty added to account for ignored contributions (e.g. scattering radius uncertainty)



Ordinate scales are % relative standard deviation and barns.

Abscissa scales are energy (eV).





Correlation Matrix




Δσ/σ vs. E for 232Th(n,γ)
Ordinate scales are % relative standard deviation and barns.
Abscissa scales are energy (eV).
σ vs. E for 232Th(n,γ)
Correlation Matrix
1.0
0.8
0.6
0.4
0.2
0.0
-1.0
-0.8
-0.6
-0.4
-0.2
0.0



Ordinate scales are % relative standard deviation and barns.

Abscissa scales are energy (eV).





Correlation Matrix



^{232}Th - Validation

- Broad-group average cross sections
- Integral benchmarks

3-group average cross sections

- E-breakpoints: 1e-4, 0.55, 2e6, 20e6 eV
- Spectrum 293.6K thermal up to 0.25 eV, 1/E up to 2 MeV, fission spectrum tail (kT=1.4 MeV)

MT	Sig_th [b]	Sig_e [b]	Sig_f [b]
1	18.4 1.6%	24.8 2.0%	8.73 1.7%
2	13.0 2.2%	17.9 2.7%	6.59 2.4%
18			.0434 2.1%
102	5.17 1.5%	6.91 1.9%	0.125 11%
16+24+41			.0122 20%

Comparison with measured values

- Ref. Kayzero database: Sig_0, Q_0=RI/Sig_0
- EXFOR: Sig_Cf

Reaction	Measured	C/E-1
Sig_0(n,g) [b]	7.37 0.34 %	-0.4 1.5%
Q0(n,g)	11.53 3.6 %	-0.5 2.4%
Sig_Cf(n,g) [b]	0.0891 2.4 % (?)	+2.4 12 %
Sig_Cf(n,f) [b]	0.0829 9.1 % (?)	-4.5 2.1 %

W-isotopes - Status

- Cross-sections essentially unchanged
- Work on covariances to include elemental cross section measurements
- Validation results – big improvement compared to ENDF/B-VI.8, but criticality is overpredicted; some problems with ^{55}Mn activation in FNG-W benchmark
- LLNL EGAF levels to be used …
- New resonance measurements at Geel …



Ordinate scales are % relative standard deviation and barns.

Abscissa scales are energy (eV).



Correlation Matrix







Ordinate scales are % relative standard deviation and barns.

Abscissa scales are energy (eV).



Correlation Matrix





^{55}Mn - Status of existing files

- Limited to 20 MeV
- All recent evaluations adopt the same cross section data
- No covariances except an ad-hoc evaluation in IRDF-2002, and model-based covariances in TENDL
- RI inconsistent with Kayzero
- Problem with FNG-W benchmark Mn foil activation measurement


25-Mn-55(N,G),SIG
nMn-55_ib8:Chi/dof=603.1%
nMn-55_ib8c:Chi/dof=605.5%
nMn-55_Irdf2002:Chi/dof=681.9%
1980 Magnusson
1979 Budnar
1978 Garg
1976 Le Rigoleur
1976 Schwerer
1974 Vuletin
1969 Dovbenko
1967 Menlove
Cross Section (barns)
E (MeV)
1
10
10^{-2}
10^{-3}

Resonance data

- H. Derrien et al., PHYSOR 2008


25-Mn-55(N,G),SIG
Cross Section (barns)
E (MeV)
nMn-55_ib8c:Chi/dof=605.4%
nMn-55_Irdf2002:Chi/dof=681.9%
1978 Garg
1975 Widder
1970 Schuman

Final covariance matrix

- Prior by MC sampling of model parameters with EMPIRE-3
- GANDR adjustment of cross sections and covariances by GLSQ



Ordinate scales are % relative standard deviation and barns.

Abscissa scales are energy (eV).

Warning: some uncertainty data were suppressed.



Correlation Matrix





Verification and validation

- Formal correctness
  - CHECKR, FIZCON, PSYCHE, Pre-Pro
- Thermal capture and resonance integral
- Cf-252 spectrum averaged cross section
- FNG-W Mn-55 foil activation
- Oktavian Mn sphere leakage spectrum

Sig0 and RI

	σ_0 [b]	d σ_0 [%]	Diff. [%]	RI [b]	dRI [%]	Diff. [%]
Kayzero	13.18	0.92	Ref	13.88	2.8	Ref
Mughabghab	13.36	0.37	1.34	13.4	3.73	-3.47
IRDF-2002	13.18	0.92	0	11.31	7.56	-18.53
Present	**13.28**	**1.53**	**0.74**	**13.22**	**6.51**	**-4.73**

Cf-252 sp.aver. x.s.

Target	Csikai [mb]	Unc. [%]	IRDF-2002 [mb]	Unc. [%]	C/E-1 [%]	Unc [%]	New [mb]	Unc. [%]	C/E-1 [%]	Unc. [%]
Mn-55	2.96	7.09	2.84	15.6	-4.0	17.1	2.49	8.41	-15.9	11.0

FNG-W Mn foil activation




FNG-Tungsten Benchmark
Neutron spectra
Spectrum Integral (10^{-6})
25
20
15
10
5
10^3
2
4
6
8
10^4
2
4
6
8
10^5
2
4
6
8
10^6
2
4
6
8
10^7
Energy (eV)
IRDF-2002
Present
No background

Oktavian Mn sphere

- Measured leakage spectrum
- Good results with ENDF/B-VII
- Highly sensitive to (n,n’)
- ENDF/B-VII inelastic unsupported by differential cross section data
- Good performance due to cancellation of errors?


OKTAVIAN MANGANESE EXPERIMENT: NEUTRON LEAKAGE SPECTRA
SPECTRUM n/Lethargy
IB-8
ENDF/B-VII
EXP.
Maximum Difference
-70.43 %
at X= 0.100
Ratios
ENERGY (Mev)


Empire
Empire El1.26E+5
Empire El9.84E+5
ENDF/B-VII.0: MN-55(N,INL)MN-55
1963 Glazkov
1963 Thomson
1973 Fujita
1994 Lashuk
1977 Correia
1975 Almen-Ramstrom
1971 Degtjarev
1968 Barrows
1963 Glazkov
1963 Lamb
1959 Nath
1994 Lashuk
1978 Schweitzer
1977 Correia
1975 Almen-Ramstrom
1971 Degtjarev
1968 Barrows
1963 Glazkov
1963 Lamb
1959 Nath
1955 Beghian
1.5
1.0
0.5
0
1
Cross Section (barns)
Incident Energy (MeV)

^{55}Mn Conclusions

- Mn-55 evaluation – work in progress
- Marked improvement of RI
- “low” tendency in Cf-252 spectrum averaged cross section
- FNG-W Mn-55 activation 60% high → problem with measurement?
- Inelastic cross section to be resolved