

## <sup>290</sup>Lv

<sup>290</sup>Lv was first identified by Oganessian et al. in “Measurements of cross sections for the fusion-evaporation reactions  $^{244}\text{Pu}(^{48}\text{Ca},\text{xn})^{292-x}\text{114}$  and  $^{245}\text{Cm}(^{48}\text{Ca},\text{xn})^{293-x}\text{116}$ ” in 2004 ([2004Og07](#)). A 243 MeV  $^{48}\text{Ca}$  beam from the Dubna U400 cyclotron bombarded a  $\text{CmO}_2$  target enriched in  $^{245}\text{Cm}$ . <sup>290</sup>Lv was populated in (3n) fusion-evaporation reactions. The residues were separated with a gas-filled recoil separator and implanted in a semiconductor detector array. Subsequent  $\alpha$  particle decay and spontaneous fission events were recorded in this array and in eight detectors arranged in a box configuration around the implantation detector. “As a result, the longer ER- $\alpha$  –  $\alpha$  –  $\alpha$ -SF chains observed in the  $^{245}\text{Cm}+^{48}\text{Ca}$  reaction must arise from the decay of  $^{291}\text{116}$  produced via the 2n-evaporation channel. Then, the shorter chains should be assigned to the decay of even-even  $^{290}\text{116}$ , the product of 3n-evaporation.” Two decay chains of the isotope were observed.

Adapted from reference ([2013Th02](#))

[2004Og07](#) Yu. Ts. Oganessian, V. K. Utyonkov, Yu. V. Lobanov, F. Sh. Abdullin *et al.*, Phys. Rev. C **69**, 054607 (2004).

[2013Th02](#) M. Thoennessen, At. Data Nucl. Data Tables **99**, 312 (2013).

Please cite this abstract as: “FRIB Nuclear Data Group, *Discovery of Nuclides Project*, Isotope Database, doi:[10.11578/frib/2279152](https://doi.org/10.11578/frib/2279152)”