

## <sup>84</sup>As

In “Identification of new arsenic isotopes in fission: <sup>83</sup>As and <sup>84</sup>As” del Marmol reported the discovery of <sup>84</sup>As at the Centre d’Etude de l’Energie Nucléaire in Belgium in 1968 ([1968De19](#)). Thermal neutrons from a BR2 reactor irradiated a solution of <sup>235</sup>U, <sup>76</sup>As tracer, As<sup>+5</sup>, Sb<sup>+5</sup> and SeO<sub>3</sub> dissolved in sulfuric acid. “A least-squares analysis, weighted for initial bromine activities, gives half-lives of 14.1±1.1 sec for <sup>83</sup>As and of 5.8±0.5 sec for <sup>84</sup>As.”

Adapted from reference ([2010Sh34](#))

[1968De19](#) P. del Marmol, J. Inorg. Nucl. Chem. **30**, 2873 (1968).

[2010Sh34](#) A. Shore, A. Fritsch, M. Heim, A. Schuh, and M. Thoennessen, At. Data Nucl. Data Tables **96**, 299 (2010).

Please cite this abstract as: “FRIB Nuclear Data Group, *Discovery of Nuclides Project*, Isotope Database, doi:[10.11578/frib/2279152](https://doi.org/10.11578/frib/2279152)”