

¹⁰⁶Sr

The discovery of ¹⁰⁶Sr was reported in the 2010 article “Identification of 45 new neutron-rich isotopes produced by in-flight fission of a ²³⁸U Beam at 345 MeV/nucleon,” by Ohnishi et al. ([2010Oh02](#)). The experiment was performed at the RI Beam Factory at RIKEN, where the new isotopes were created by in-flight fission of a 345 MeV/nucleon ²³⁸U beam on a beryllium target. ¹⁰⁶Sr was separated and identified with the BigRIPS superconducting in-flight separator. The results for the new isotopes discovered in this study were summarized in a table. Twenty-two counts for ¹⁰⁶Sr were recorded.

Adapted from reference ([2012Pa21](#))

- [2010Oh02](#) T. Ohnishi, T. Kubo, K. Kusaka, A. Yoshida *et al.*, J. Phys. Soc. Jap. **79**, 073201 (2010).
- [2012Pa21](#) A. M. Parker and M. Thoennessen, At. Data Nucl. Data Tables **98**, 812 (2012).

Please cite this abstract as: “FRIB Nuclear Data Group, *Discovery of Nuclides Project*, Isotope Database, doi:[10.11578/frib/2279152](https://doi.org/10.11578/frib/2279152)”