

¹¹²Rh

In the 1971 paper “Low-energy transitions from the deexcitation of spontaneous fission fragments of ²⁵²Cf,” Hopkins et al. reported the observation of ¹¹²Rh at the Center for Nuclear Studies of the University of Austin ([1971Ho29](#)). Fission fragments from the spontaneous fission of ²⁵²Cf were measured in coincidence with X-rays and low-energy γ -rays. In a table the energy (60.5 keV) of the first excited state was identified correctly. This energy had also been reported earlier ([1970Jo20](#)), however, no specific element assignment was made.

This assignment was changed from the original compilation ([2012Pa21](#)) which credited a later paper by Hopkins et al. ([1972Ho08](#)) with the discovery of ¹¹²Rh.

- [1970Jo20](#) W. John, F. W. Guy, and J. J. Wesolowski, Phys. Rev. C **2**, 1451 (1970).
[1971Ho29](#) F. F. Hopkins, G. W. Phillips, J. R. White, C. F. Moore, and P. Richard, Phys. Rev. C **4**, 1927 (1971).
[1972Ho08](#) F. F. Hopkins, J. R. White, G. W. Phillips, C. F. Moore, and P. Richard, Phys. Rev. C **5**, 1015 (1972).
[2012Pa21](#) A. M. Parker and M. Thoennessen, At. Data Nucl. Data Tables **98**, 812 (2012).

Please cite this abstract as: “FRIB Nuclear Data Group, *Discovery of Nuclides Project*, Isotope Database, doi:[10.11578/frib/2279152](https://doi.org/10.11578/frib/2279152)”