

^{110}Te

Kirchner et al. reported the first observation of ^{110}Te in the paper “New neutron-deficient tellurium, iodine, and xenon isotopes produced by reactions of 290 MeV ^{58}Ni ions on ^{58}Ni and ^{63}Cu targets” in 1977 ([1977Ki11](#)). An enriched ^{58}Ni target was bombarded with a 290 MeV ^{58}Ni beam forming ^{110}Te in (4p2n) or (α 2p) fusion-evaporation reactions. Beta-, gamma-, and X-rays, as well as protons and α particles were measured following mass separation with the GSI on-line mass separator facility. “We wish to report in this letter the identification of the new neutron-deficient isotopes $^{108-110}\text{Te}$, $^{110-114}\text{I}$, and ^{114}Xe .”

Adapted from reference ([2013Ka01](#))

- [1977Ki11](#) R. Kirchner, O. Klepper, G. Nyman, W. Reisdorf *et al.*, Phys. Lett. B **70**, 150 (1977).
[2013Ka01](#) J. Kathawa, C. Fry, and M. Thoennessen, At. Data Nucl. Data Tables **99**, 22 (2013).

Please cite this abstract as: “FRIB Nuclear Data Group, *Discovery of Nuclides Project*, Isotope Database, doi:[10.11578/frib/2279152](https://doi.org/10.11578/frib/2279152)”