

## $^{140}\text{Te}$

In 1994, Bernas et al. published the discovery of  $^{140}\text{Te}$  in “Projectile fission at relativistic velocities: a novel and powerful source of neutron-rich isotopes well suited for in-flight isotopic separation” ([1994Be24](#)). The isotope was produced using projectile fission of  $^{238}\text{U}$  at 750 MeV/nucleon on a lead target at GSI, Germany. “Forward emitted fragments from  $^{80}\text{Zn}$  up to  $^{155}\text{Ce}$  were analyzed with the Fragment Separator (FRS) and unambiguously identified by their energy-loss and time-of-flight.” This experiment yielded 264 counts of  $^{140}\text{Te}$ .

Adapted from reference ([2013Ka01](#))

[1994Be24](#) M. Bernas, S. Czajkowski, P. Armbruster, H. Geissel *et al.*, Phys. Lett. B **331**, 19 (1994).

[2013Ka01](#) J. Kathawa, C. Fry, and M. Thoennessen, At. Data Nucl. Data Tables **99**, 22 (2013).

Please cite this abstract as: “FRIB Nuclear Data Group, *Discovery of Nuclides Project*, Isotope Database, doi:[10.11578/frib/2279152](https://doi.org/10.11578/frib/2279152)”