

^{127}Ce

The discovery of the isotope ^{127}Ce was first presented in the 1978 paper “New Neutron-Deficient Isotopes of Lanthanum and Cerium” by Bogdanov et al. (1978Bo32). A 190 MeV ^{36}S beam accelerated by the U-300 heavy-ion cyclotron of the Joint Institute for Nuclear Research (JINR) facility at Dubna, bombarded targets of ^{96}Ru and ^{98}Ru . The fusion-evaporation residues were mass separated with the on-line BEMS-2 facility and their X-ray and β emission was detected with a Ge(Li) spectrometer and a plastic counter, respectively. Half-lives were determined from the X-ray decay curves. “Seven isotopes $^{123-125}\text{La}$ and $^{124-127}\text{Ce}$ have been first observed and their half-lives and low-energy γ -ray data are reported.” The measured half-life for ^{127}Ce was 32(4) s.

Adapted from reference (2009Gi07)

- 1978Bo32 D. D. Bogdanov, A. V. Demyanov, V. A. Karnaukhov, M. Nowicki *et al.*, Nucl. Phys. A **307**, 421 (1978).
2009Gi07 J. Q. Ginepro, J. Snyder, and M. Thoennessen, At. Data Nucl. Data Tables **95**, 805 (2009).

Please cite this abstract as: “FRIB Nuclear Data Group, *Discovery of Nuclides Project*, Isotope Database, doi:[10.11578/frib/2279152](https://doi.org/10.11578/frib/2279152)”