

## $^{157}\text{Ce}$

In 2017, Wu et al. discovered  $^{157}\text{Ce}$  in “94  $\beta$ -Decay Half-Lives of Neutron-Rich  $_{55}\text{Cs}$  to  $_{67}\text{Ho}$ : Experimental Feedback and Evaluation of the r-Process Rare-Earth Peak Formation” (2017Wu04). A 345 MeV/nucleon  $^{238}\text{U}$  beam from RIBF was used to produce fission fragments which were separated with the BigRIPS separator and the ZeroDegree spectrometer. The nuclides were identified with the TOF-B $\rho$ - $\Delta E$  method and their  $\beta$ -decay half-lives were measured with the Wide range Active Silicon-Strip Stopper Array for Beta and ion detection (WAS3ABi). The extracted half-life for  $^{157}\text{Ce}$  was 0.175(41) s as listed in the supplemental material of the paper.

2017Wu04 J. Wu, S. Nishimura, G. Lorusso, P. Moller *et al.*, Phys. Rev. Lett. **118**, 072701 (2017).

Please cite this abstract as: “FRIB Nuclear Data Group, *Discovery of Nuclides Project*, Isotope Database, doi:[10.11578/frib/2279152](https://doi.org/10.11578/frib/2279152)”