

## $^{182}\text{Er}$

Fukuda et al. reported the discovery of  $^{182}\text{Er}$  in the 2026 article “First Observation of Twenty-two Exotic Isotopes Using a  $^{208}\text{Pb}$  Primary Beam at RIBF” ([2026Fu06](#)). A 345-MeV/nucleon  $^{208}\text{Pb}$  from the RI Beam Factory at RIKEN in Japan, impinged on 2–5 mm thick Be targets and resulting fragments were separated and identified with the large-acceptance two-stage BigRIPS separator “Consequently, a total of 17 new neutron-rich isotopes were identified for the first time in this study:  $^{179}_{67}\text{Ho}$ ,  $^{181,182}_{68}\text{Er}$ ,  $^{184-186}_{69}\text{Tm}$ ,  $^{188,189}_{70}\text{Yb}$ ,  $^{191}_{71}\text{Lu}$ ,  $^{193,194}_{72}\text{Hf}$ ,  $^{195,196}_{73}\text{Ta}$ ,  $^{198,199}_{74}\text{W}$ , and  $^{200,201}_{75}\text{Re}$ .”

[2026Fu06](#) N. Fukuda, S. Michimasa, H. Suzuki, Y. Shimizu *et al.*, Prog. Theor. Exp. Phys. **2026**, 061D01 (2026).

Please cite this abstract as: “FRIB Nuclear Data Group, *Discovery of Nuclides Project*, Isotope Database, doi:[10.11578/frib/2279152](https://doi.org/10.11578/frib/2279152)”