

## $^{147}\text{Yb}$

In 2026, Suzuki et al. discovered  $^{147}\text{Yb}$  in “Discovery of proton-rich radioactive isotopes in the  $Z = 63\text{-}70$  region produced by the projectile fragmentation of a 345-MeV/nucleon  $^{238}\text{U}$  beam” (2026Su06). The RIKEN RI factory delivered the  $^{238}\text{U}$  beam to a 1 mm thick beryllium target. The large-acceptance two-stage separator Big-RIPS was used to separate the fragments. A thin tantalum foil at the first dispersive focus reduced the fraction of less-exotic contaminants. The fragments were identified with the TOF– $B\rho - \Delta E$  method. “In total, 14 new isotopes were obtained with the  $^{135}\text{Tb}$  and  $^{146}\text{Yb}$  settings based on the evaluation shown in the NNDC as of October 2025:  $^{132,133}_{63}\text{Eu}$ ,  $^{133,134,136}_{64}\text{Gd}$ ,  $^{136,137,138}_{65}\text{Tb}$ ,  $^{138}_{66}\text{Dy}$ ,  $^{143}_{67}\text{Ho}$ ,  $^{143}_{68}\text{Er}$ ,  $^{144}_{69}\text{Tm}$ , and  $^{147,148}_{70}\text{Yb}$ .” 3 events of  $^{147}\text{Yb}$  were observed.

[2026Su06](#) H. Suzuki, N. Fukuda, H. Takeda, Y. Shimizu *et al.*, Prog. Theor. Exp. Phys. **2026**, 023 (2026).

Please cite this abstract as: “FRIB Nuclear Data Group, *Discovery of Nuclides Project*, Isotope Database, doi:[10.11578/frib/2279152](https://doi.org/10.11578/frib/2279152)”