

²¹⁰Fr

In 1964, Griffioen and MacFarlane reported the identification of ²¹⁰Fr in the paper “Alpha-decay properties of some francium isotopes near the 126-neutron closed shell” (1964Gr04). ¹⁹⁷Au, ^{203,205}Tl, and ²⁰⁸Pb targets were bombarded with ¹⁶O, ¹²C, and ¹¹B beams with energies up to 10.38 MeV/amu from the Berkeley HILAC. Recoil products were collected on a catcher foil which was positioned in front of gold surface-barrier detector which measured subsequent α decay. “Fr²¹¹ and Fr²¹⁰: ... These facts seem to indicate that this group is due to two different isotopes, Fr²¹¹ and Fr²¹⁰, which were formed by (C¹²,6n) and (C¹²,7n) reactions, respectively... ” The measured half-life was 159(5) s.

Adapted from reference (2013Fr09)

1964Gr04 R. D. Griffioen and R. D. Macfarlane, Phys. Rev. **133**, B1373 (1964).
2013Fr09 C. Fry and M. Thoennessen, At. Data Nucl. Data Tables **99**, 497 (2013).

Please cite this abstract as: “FRIB Nuclear Data Group, *Discovery of Nuclides Project*, Isotope Database, doi:[10.11578/frib/2279152](https://doi.org/10.11578/frib/2279152)”