

## <sup>218</sup>Ac

In the 1970 article “Production and decay properties of protactinium isotopes of mass 222 to 225 formed in heavy-ion reactions,” Borggreen et al. identified <sup>218</sup>Ac (1970Bo13). The Berkeley heavy-ion linear accelerator (HILAC) was used to bombard <sup>209</sup>Bi, <sup>208</sup>Pb and <sup>205</sup>Tl targets with <sup>16</sup>O, <sup>19</sup>F and <sup>22</sup>Ne beams forming <sup>224</sup>Pa, <sup>223</sup>Pa, and <sup>222</sup>Pa in (xn) fusion-evaporation reactions. <sup>218</sup>Ac was then populated by  $\alpha$ -decay. Recoil products were deposited by a helium gas stream on a metal surface located in front of a gold surface-barrier detector which recorded the subsequent  $\alpha$  decay. “<sup>218</sup>Ac has a 0.27- $\mu$ sec half-life, so that it appears in these spectra only by its continued replenishment by the decay of <sup>222</sup>Pa.”

Adapted from reference (2013Fr03)

1970Bo13 J. Borggreen, K. Valli, and E. K. Hyde, Phys. Rev. C **2**, 1841 (1970).

2013Fr03 C. Fry and M. Thoennessen, At. Data Nucl. Data Tables **99**, 345 (2013).

Please cite this abstract as: “FRIB Nuclear Data Group, *Discovery of Nuclides Project*, Isotope Database, doi:10.11578/frib/2279152”