

²⁴⁰Cf

The 1970 discovery of ²⁴⁰Cf by Silva et al. was described in the paper “New isotopes ²⁴¹Cf and ²⁴⁰Cf” (1970Si19). Enriched ²³³U, ²³⁴U, and ²³⁵U targets were bombarded with 118 MeV ¹²C beams from the Oak Ridge isochronous cyclotron. Reaction products were deposited with a helium jet on a platinum disk which was then moved in front of a Si(Au) detector to measure α -particles. “The half-lives of the 7.335- and 7.590-MeV α activities were determined to be 3.78 ± 0.70 and 1.06 ± 0.15 min, respectively, from computer least squares fits to 200-300 recorded decay events... the excitation functions for the 7.335- and 7.590-MeV activities support the assignment of the former α -particle activity to the decay of ²⁴¹Cf and of the latter to ²⁴⁰Cf.”

Adapted from reference (2013Fr02)

1970Si19 R. J. Silva, R. L. Hahn, K. S. Toth, M. L. Mallory *et al.*, Phys. Rev. C **2**, 1948 (1970).

2013Fr02 C. Fry and M. Thoennessen, At. Data Nucl. Data Tables **99**, 96 (2013).

Please cite this abstract as: “FRIB Nuclear Data Group, *Discovery of Nuclides Project*, Isotope Database, doi:[10.11578/frib/2279152](https://doi.org/10.11578/frib/2279152)”