

¹⁰⁰Ag-1

The 2.24 m isomer of ¹⁰⁰Ag was first observed in 1980 by H.I.Hayakawa et al. at McGill University, Montreal, Canada ([1980Ha20](#)). The assignment of a 2.3(1) m isomer to ¹⁰⁰Ag in 1970 ([1970Hn03](#)) was based on an incorrect assignment of a 8 m state to ¹⁰⁰Ag ([1966Bu05,1967Do06](#)).

- [1966Bu05](#) F. D. S. Butement and M. Y. Mirza, J. Inorg. Nucl. Chem. **28**, 303 (1966).
[1967Do06](#) D. R. Doty, J. R. Richardson, and J. W. Sunier, Nucl. Phys. A **103**, 417 (1967).
[1970Hn03](#) D. J. Hnatowich, E. Hagebo, A. Kjelberg, R. Mohr, and P. Patzelt, J. Inorg. Nucl. Chem. **32**, 3137 (1970).
[1980Ha20](#) H. I. Hayakawa, I. Hyman, and J. K. P. Lee, Phys. Rev. C **22**, 247 (1980).

Please cite this abstract as: “FRIB Nuclear Data Group, *Discovery of Nuclides Project*, Isomer Database, doi:[10.11578/frib/2572219](https://doi.org/10.11578/frib/2572219)”