

Adopted Levels

Type	History		Literature Cutoff Date
	Author	Citation	
Full Evaluation	Balraj Singh	ENSDF	20-Jul-2015

$Q(\beta^-)=12210$ SY; $S(n)=2740$ SY; $S(p)=17670$ CA; $Q(\alpha)=-11700$ CA [2012Wa38,1997Mo25](#)

Estimated uncertainties ([2012Wa38](#)): 640 for $Q(\beta^-)$, 710 for $S(n)$.

$Q(\beta^-)$ and $S(n)$ from [2012Wa38](#); $S(p)$ and $Q(\alpha)$ from [1997Mo25](#).

$Q(\beta^-n)=7210$ 580, $S(2n)=7560$ 640 (syst,[2012Wa38](#)). $S(2p)=33520$ ([1997Mo25](#),theory).

[2010Oh02](#): ^{117}Mo nuclide identified in $\text{Be}(^{238}\text{U},\text{F})$ and $\text{Pb}(^{238}\text{U},\text{F})$ reactions with a $^{238}\text{U}^{86+}$ beam energy of 345 MeV/nucleon produced by the cascade operation of the RBIF accelerator complex of the linear accelerator RILAC and four cyclotrons RRC, fRC, IRC and SRC. Identification of ^{117}Mo nuclei was made on the basis of magnetic rigidity, time-of-flight and energy loss of the fragments using BigRIPS fragment separator. Experiments performed at RIKEN facility. Based on A/Q spectrum and Z versus A/Q plot, 78 counts were assigned to ^{117}Mo isotope. (Q=charge state).

[2015Lo04](#): ^{117}Mo nuclide produced at RIBF-RIKEN facility in $^9\text{Be}(^{238}\text{U},\text{F})$ reaction at $E=345$ MeV/nucleon with an average intensity of 6×10^{10} ions/s. Identification of ^{117}Mo was made by determining atomic Z and mass-to-charge ratio A/Q, where Q=charge state of the ions. The selectivity of ions was based on magnetic rigidity, time-of-flight and energy loss. The separated nuclei were implanted at a rate of 50 ions/s in a stack of eight double-sided silicon-strip detector (WAS3ABi), surrounded by EURICA array of 84 HPGe detectors. Correlations were recorded between the implanted ions and β rays. The half-life of ^{117}Mo isotope was measured from the correlated ion- β decay curves and maximum likelihood analysis technique as described in [2014Xu07](#). Comparison of measured half-lives with FRDM+QRPA, KTUY+GT2 and DF3+CQRPA theoretical calculations.

 ^{117}Mo Levels

E(level)	$T_{1/2}$	Comments
0	22 ms 5	$\% \beta^- = ?$; $\% \beta^- n = ?$; $\% \beta^- 2n = ?$ Theoretical $T_{1/2}=42.9$ ms, $\% \beta^- n=16.8$, $\% \beta^- 2n=0.32$ (2003Mo09). Measured $\sigma=4$ pb (2010Oh02), systematic uncertainty $\approx 40\%$. $T_{1/2}$: measured by 2015Lo04 from (implanted ions) β correlated curves in time and position using maximum likelihood method. See 2015Lo04 for comparison of their experimental value with theoretical values. Probability of misidentification of ^{117}Mo isotope $< 0.001\%$ (2010Oh02).