¹²C(¹⁶O, ¹²C) **1979Do01**

History

Type Author Citation Literature Cutoff Date
Full Evaluation J. H. Kelley, J. E. Purcell and C. G. Sheu NP A968,71 (2017)

1-Jan-2017

1979Do01: ${}^{12}\text{C}({}^{16}\text{O}, {}^{12}\text{C})$ E=315 MeV; measured $\sigma(\text{E},\theta)$.

1994Su09: ¹²C(¹⁶O, ¹²C) E=28.5-33.5 MeV; measured magnetic substate population for ¹²C*(4.44 MeV), deduced intermediate structure resonances.

1995Fr05: ${}^{12}C({}^{16}O, {}^{12}C)$ E=99 MeV; measured $Q(\beta^-)$ value spectra.

1996Fr09: 12 C(16 O, 12 C) E=51-66 MeV; measured Q(β^-)value spectra.

2004Su10: 12 C(16 O, 12 C) E=17.4-23 MeV; measured E_{γ}, I_{γ} (particle) γ coincidences.

2006Sz06: ¹²C(¹⁶O, ¹²C) E=62-124 MeV; measured particle spectra.

2011Ha23: 12 C(16 O, 12 C) E=20,24,28 MeV, measured E(particle), I(particle, θ). Deduced $\sigma(\theta)$, optical potential parameters.

2014Oh04: XUNDL dataset compiled by TUNL, 2014.

The authors analyzed the Airy structures present in inelastic $^{16}O+^{12}C$ scattering to $^{12}C*(4.44 \text{ MeV})$ using 170-280 MeV ^{16}O beams, from the Jyvaskyla cyclotron. Scattered particles were detected at $\theta_{c.m.}=7^{\circ}-40^{\circ}$ using a position sensitive $\Delta E-E$ Si detector telescope; at larger angles ($\theta_{c.m.}>40^{\circ}$) a position sensitive gas proportional counter/Si detector $\Delta E-E$ array was used. Analyzed angular distributions for scattering to $^{12}C*(4.44 \text{ MeV}: J^{\pi}=2^{+})$ via an extended double folding coupled-channels model. The angular distributions are well reproduced with an emphasis on the large angle so-called rainbow region where diffraction effects and Airy structures are prominent. Discussed the couplings between elastic and inelastic components. See also (2015Ma12).

¹²C Levels

E(level)	$T_{1/2}$
0	
4.44×10^3	
7.65×10^3	
9.64×10^{3}	
10.8×10^{3}	
14.1×10^3	
15.8×10^{3}	
21.6×10^{3}	
25.3×10^{3}	≈4 [†] MeV
26.7×10^{3}	≈4 [†] MeV

[†] From (1979Do01).