

$^{12}\text{C}(^3\text{He},^3\text{He})$ 1971St22,1977Bu03

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	J. H. Kelley, J. E. Purcell and C. G. Sheu		NP A968,71 (2017)	1-Jan-2017

- 1966Sc12: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=2.0-6.1 MeV, measured $\sigma(E,\theta)$.
- 1966Sc22: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=8.5-10.0 MeV, measured $\sigma(E,\theta)$.
- 1968Fo06: $^{12}\text{C}(^3\text{He},^3\text{He}),(^3\text{He},^3\text{He}')$ E=12-19 MeV, measured $\sigma(E,\theta)$, $\sigma(E,E(^3\text{He}'),\theta)$. Deduced optical model parameters.
- 1968La19: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=5.29-5.50 MeV, measured $\sigma(E,E_p,\theta)$. Deduced optical potentials, level overlap parameter.
- 1968We15,1969We03,1971Ja01: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=2.0-8.0 MeV, measured $\sigma(E,E(^3\text{He}),\theta)$.
- 1969Ar08: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=36 MeV, measured $\sigma(\theta)$. Deduced optical model parameters.
- 1969En03: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=31.6 MeV, measured polarization P. Deduced optical model spin-orbit potential.
- 1969Zu02: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=15 MeV, measured $\sigma(\theta)$.
- 1970Mc10,1972Mc01: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=18,20 MeV, measured P(θ), $\sigma(\theta)$. Deduced optical model parameters.
- 1970Sc23: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=11.0 MeV, measured $\sigma(\theta)$. Deduced optical-model parameters.
- 1971St22: $^{12}\text{C}(^3\text{He},^3\text{He}')$, E=19,19.5,20 MeV; measured $\sigma(E_p)$, $\sigma(^3\text{He}')$. ^{12}C deduced energies of 1st, 2nd excited states.
- 1973Fu03: $^{12}\text{C}(^3\text{He},^3\text{He}),(^3\text{He},^3\text{He}')$ E=24.0,29.2,34.7,39.6 MeV, measured $\sigma(E(^3\text{He}'),\theta)$. Deduced optical model parameters. ^{12}C deduced levels, β .
- 1973Wi07: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=217 MeV, measured $\sigma(E(^3\text{He}),\theta)$. Deduced optical model parameters.
- 1975Bo06: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=28 MeV, measured polarization angular distribution. Deduced optical model spin-orbit potential.
- 1976Ma26: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=18,20,22,24.5 MeV, measured $\sigma(\theta)$.
- 1976Ta12: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=82.1 MeV, measured $\sigma(\theta)$. Deduced optical potential parameters. $^{12}\text{C}(^3\text{He},^3\text{He}')$ E=81.2 MeV, measured $\sigma(\theta)$. ^{12}C levels deduced β .
- 1977Ba05: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=39.1 MeV, measured $\sigma(\theta)$. Deduced optical model parameters.
- 1977Bu03: $^{12}\text{C}(^3\text{He},^3\text{He}')$, E=130 MeV; measured $\sigma(E(^3\text{He}'),\theta)$. ^{12}C deduced giant resonances, deformation length.
- 1975Bu11,1977Ka25: $^{12}\text{C}(\text{pol. } ^3\text{He},^3\text{He})$ E=20.5-33 MeV, measured $\sigma(\theta,E)$, A(θ,E). Deduced optical model potential, phase shifts, β_2 .
- 1979Go07: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=44.04 MeV, measured $\sigma(\theta)$.
- 1980Hy02: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=119.0 MeV, measured $\sigma(\theta)$. Deduced optical model parameters.
- 1980Tr02: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=41 MeV, measured $\sigma(\theta)$.
- 1982Al14: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=40.9 MeV, measured $\sigma(\theta)$. Deduced absorptive term characteristics.
- 1991Go25: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=98 MeV, measured $\sigma(\theta)$. Deduced model parameters.
- 1992Ad06: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=50,60 MeV, measured $\sigma(\theta)$. Deduced model parameters. ^{12}C levels deduced multipole deformation parameters. Optical model and DWBA analyses.
- 1995Da08,1995Da21: $^{12}\text{C}(^3\text{He},^3\text{He}),(^3\text{He},^3\text{He}')$ E=98 MeV, measured $\sigma(\theta)$. Deduced nuclear rainbow effect evidence, model parameters.
- 1995Ya06: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=450 MeV, measured $\sigma(\theta)$. Deduced optical potential parameters.
- 1997Kh07: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=27.4-217 MeV, analyzed $\sigma(\theta)$. Deduced parameters, reaction σ .
- 2001Ku20: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=17-20 MeV/nucleon, analyzed $\sigma(\theta)$. Deduced S-matrix parameters.
- 2003Ka24: $^{12}\text{C}(^3\text{He},^3\text{He})$ E=443 MeV, measured $\sigma(\theta)$.
- 2009Da22: $^{12}\text{C}(^3\text{He},^3\text{He}),(^3\text{He},^3\text{He}')$ E=34.7,50,60,72,82 MeV, analyzed elastic and inelastic scattering cross section and $\sigma(\theta)$ data using diffraction model of scattering. Deduced nuclear rms radii for excited states in ^{12}C .
- 2013Ha01: XUNDL dataset compiled by TUNL, 2013.
- Beams of 50.5 and 60 MeV ^3He ions from the Almaty-Kazakhstan U-150M cyclotron impinged on a $30 \mu\text{g}/\text{cm}^2$ and the angular distribution of scattered ions was measured for $15^\circ < \theta_{\text{c.m.}} < 170^\circ$ by a set of four moveable ΔE -E Si telescopes. Angular distributions were analyzed using a coupled channels method and the location of the Airy minimum was deduced. Analysis of scattering to the Hoyle state suggests it is a 3α particle condensate with a large radius of dilute matter.
- 2014Wh02: XUNDL dataset compiled by TUNL, 2014.
- Unbound ^{12}C states were studied using 46 MeV ^3He ions, from the Birmingham MC40 cyclotron, that impinged on a $300 \mu\text{g}/\text{cm}^2$ carbon target. The scattered ^3He ejectiles were detected using a position sensitive ΔE -E Si-strip detector telescope that was placed 9.8 cm from the target at $\theta_{\text{lab}} = -32.5^\circ$. Additional details on the reaction were determined by measurement of α -particles from the breakup of recoiling ^{12}C . The α -particles were detected in an array of 2 ΔE -E Si-strip telescopes placed at $\theta_{\text{lab}} = 27.5^\circ$ and 57.5° . The data were analyzed with a kinematic gate that required a sequential decay via $^{12}\text{C} \rightarrow ^8\text{Be}_{\text{g.s.}} + \alpha_0$. This implies only states with natural parity were selected. Discussion on backgrounds and contaminant reactions, such as $^{12}\text{C}(^3\text{He},^7\text{Be} \rightarrow ^3\text{He} + \alpha)$ and

$^{12}\text{C}(^3\text{He},^3\text{He})$ 1971St22,1977Bu03 (continued)

$^{12}\text{C}(^3\text{He},^{11}\text{C}\rightarrow^8\text{Be}+^3\text{He})$ are given in the text.

The known states at $E_x=7.65, 9.64, 10.84$ and 14.08 were easily resolved with an experimental resolution of ≈ 530 keV. A variety of states up to $E_x=25.1$ MeV are observed and compared with literature values. By comparing the ^3He “singles” rate with the $^{12}\text{C}\rightarrow^8\text{Be}_{\text{g.s.}}+\alpha_0$ data, the authors estimated the Γ_{α_0}/Γ branching ratios for several states.

1980Le25: Unconfirmed $J^\pi=0^+$ states are reported at 9.25 MeV 20 ($\Gamma=1.8$ MeV 2) and 20.3 MeV 2 ($\Gamma=1.1$ MeV 2).

 ^{12}C Levels

E(level)	J^π	$T_{1/2}$	L	Comments
0			0	
4442.2	15		2	E(level): From (1971St22).
7655.9	25		2	E(level): From (1971St22: includes (p,p')). $R_{\text{r.m.s.}}\approx 2.94$ fm (2008De35).
9.6×10^3			3	
10.83×10^3				
10.84×10^3				
12.7×10^3			0	
14.08×10^3			0	$\Gamma_{\alpha_0}/\Gamma=0.20$ 10 (2014Wh02).
15.11×10^3			0	T=1 $\Gamma_{\alpha_0}/\Gamma<0.08$ (2014Wh02).
15.2×10^3	3	1.8 MeV 3	2	E(level), $T_{1/2}$: From (1977Bu03).
16.11×10^3			2	T=1 $\Gamma_{\alpha_0}/\Gamma=0.18$ 10 (2014Wh02).
16.58×10^3			2	T=1
18.4×10^3	6	0.4 MeV 1	2	E(level), $T_{1/2}$: From (1977Bu03). $\Gamma_{\alpha_0}/\Gamma=0.25$ 10 (2014Wh02). J^π : From (2014Wh02).
18.90×10^3	15	0.70 MeV 15	2	T=1 E(level), $T_{1/2}$: From (1977Bu03).
19.58×10^3	6		2	T=1 E(level), $T_{1/2}$: From (1969Ba06). $\Gamma_{\alpha_0}/\Gamma=0.21$ 10 (2014Wh02).
21.30×10^3	15	1.4 MeV 2	2	E(level), $T_{1/2}$: From (1977Bu03). Possibly unresolved states with $\Gamma=1.4$ MeV 2 and $\Gamma=0.43$ MeV 8.
22.2×10^3	3	<0.7 MeV	2	E(level): From (2014Wh02). This state can likely be associated with the $E_x=22.4$ MeV $J^\pi=5^-$ state that was populated in $^{12}\text{C}(\alpha,3\alpha)$ (2014Ma37).
23.5×10^3	2	0.6 MeV 2	2	E(level), $T_{1/2}$: From (1977Bu03).
25.1×10^3	3	<0.8 MeV	2	E(level): From (2014Wh02).
25.9×10^3	3	2.2 MeV 3	2	E(level), $T_{1/2}$: From (1977Bu03).
28.8×10^3	4	2.7 MeV 4	2	E(level), $T_{1/2}$: From (1977Bu03).