

Adopted Levels

Type	History		Literature Cutoff Date
	Author	Citation	
Full Evaluation	Balraj Singh	ENSDF	20-Jul-2015

$Q(\beta^-)=11380$  SY;  $S(n)=3110$  SY;  $S(p)=16820$  SY;  $Q(\alpha)=-11300$  SY [2012Wa38](#)

Estimated uncertainties ([2012Wa38](#)): 500 for  $Q(\beta^-)$ , 570 for  $S(n)$ , 640 for  $S(p)$  and  $Q(\alpha)$ .

$S(2n)=8630$  500,  $Q(\beta^-n)=5700$  450.  $S(2p)=32030$  ([1997Mo25](#),theory).

[2010Oh02](#):  $^{121}\text{Ru}$  nuclide identified in  $\text{Be}(^{238}\text{U},\text{F})$  and  $\text{Pb}(^{238}\text{U},\text{F})$  reactions with a  $^{238}\text{U}^{86+}$  beam energy of 345 MeV/nucleon produced by the cascade operation of the RBIF accelerator complex of the linear accelerator RILAC and four cyclotrons RRC, fRC, IRC and SRC. Identification of  $^{121}\text{Ru}$  nuclei was made on the basis of magnetic rigidity, time-of-flight and energy loss of the fragments using BigRIPS fragment separator. Experiments performed at RIKEN facility. Based on A/Q spectrum and Z versus A/Q plot, 170 counts were assigned to  $^{121}\text{Ru}$  isotope. (Q=charge state).

[2015Lo04](#):  $^{121}\text{Ru}$  nuclide produced at RIBF-RIKEN facility in  $^9\text{Be}(^{238}\text{U},\text{F})$  reaction at  $E=345$  MeV/nucleon with an average intensity of  $6 \times 10^{10}$  ions/s. Identification of  $^{121}\text{Ru}$  was made by determining atomic Z and mass-to-charge ratio A/Q, where Q=charge state of the ions. The selectivity of ions was based on magnetic rigidity, time-of-flight and energy loss. The separated nuclei were implanted at a rate of 50 ions/s in a stack of eight double-sided silicon-strip detector (WAS3ABi), surrounded by EURICA array of 84 HPGe detectors. Correlations were recorded between the implanted ions and  $\beta$  rays. The half-life of  $^{121}\text{Ru}$  isotope was measured from the correlated ion- $\beta$  decay curves and maximum likelihood analysis technique as described in [2014Xu07](#). Comparison of measured half-lives with FRDM+QRPA, KTUY+GT2 and DF3+CQRPA theoretical calculations.

[Additional information 1](#).

 $^{121}\text{Ru}$  Levels

E(level)	$T_{1/2}$	Comments
0	29 ms 2	$\% \beta^- = 100$ ; $\% \beta^- n = ?$ ; $\% \beta^- 2n = ?$ Theoretical $T_{1/2} = 88.3$ ms, $\% \beta^- n = 7.1$ , $\% \beta^- 2n = 0.0$ ( <a href="#">2003Mo09</a> ). E(level): measured half-life is assumed to correspond to the ground state of $^{121}\text{Ru}$ . Measured $\sigma = 143$ pb ( <a href="#">2010Oh02</a> ), systematic uncertainty $\approx 40\%$ . Probability of misidentification of $^{121}\text{Ru}$ isotope $< 0.001\%$ ( <a href="#">2010Oh02</a> ). $J^\pi$ : $3/2^-$ in theoretical calculations ( <a href="#">1997Mo25</a> ). $T_{1/2}$ : measured by <a href="#">2015Lo04</a> from (implanted ions) $\beta$ correlated curves in time and position using maximum likelihood method. See <a href="#">2015Lo04</a> for comparison of their experimental value with theoretical values.