

^{129}Sn IT decay (3.40 μs) 2002Ge07,2008Lo07

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	Janos Timar and Zoltan Elekes, Balraj Singh		NDS 121, 143 (2014)	31-May-2014

Parent: ^{129}Sn : E=1761.6 10; $J^\pi=(19/2^+)$; $T_{1/2}=3.40 \mu\text{s}$ 13; %IT decay=100.0

 ^{129}Sn Levels

E(level) [†]	J^π [†]	$T_{1/2}$ [†]	Comments
35.15 5	11/2 ⁻	6.9 min 1	% β^- =100; %IT< 2×10^{-3}
1171.48 7	(15/2 ⁻)		
1359.40 7	(13/2 ⁻)		
1741.89 7	(15/2 ⁺)		
1761.6 10	(19/2 ⁺)	3.40 μs 13	%IT=100

[†] From Adopted Levels.

 $\gamma(^{129}\text{Sn})$

E_γ [†]	I_γ [‡]	$E_i(\text{level})$	J_i^π	E_f	J_f^π	Mult. [†]	$\alpha^\#$	Comments
19.7 10		1761.6	(19/2 ⁺)	1741.89	(15/2 ⁺)	(E2)	1.0×10^3 3	$\alpha(\text{L})=7.8\times 10^2$ 23; $\alpha(\text{M})=1.6\times 10^2$ 5; $\alpha(\text{N})=27$ 9; $\alpha(\text{O})=0.58$ 17 Mult.: half-life is characteristic of an E2 transition (2002Ge07).
382.49 2	69 7	1741.89	(15/2 ⁺)	1359.40	(13/2 ⁻)			
570.41 3	100 10	1741.89	(15/2 ⁺)	1171.48	(15/2 ⁻)			
1136.31 5	88 9	1171.48	(15/2 ⁻)	35.15	11/2 ⁻			
1324.25 5	65 7	1359.40	(13/2 ⁻)	35.15	11/2 ⁻			

[†] From Adopted Gammas.

[‡] From 2008Lo07.

[#] Total theoretical internal conversion coefficients, calculated using the BrIcc code (2008Ki07) with Frozen orbital approximation based on γ -ray energies, assigned multipolarities, and mixing ratios, unless otherwise specified.

^{129}Sn IT decay (3.40 μs) 2002Ge07,2008Lo07

Decay Scheme

Intensities: Relative I_γ
 %IT=100.0

Legend

- $I_\gamma < 2\% \times I_\gamma^{\text{max}}$
- $I_\gamma < 10\% \times I_\gamma^{\text{max}}$
- $I_\gamma > 10\% \times I_\gamma^{\text{max}}$

