

¹⁴⁹Er εp decay (9.6 s) 1989Fi01

Type	Author	History Citation	Literature Cutoff Date
Full Evaluation	N. Nica	NDS 208,1 (2026)	17-Jan-2026

Parent: ¹⁴⁹Er: E=741.8 4; J^π=(11/2⁻); T_{1/2}=9.6 s 6; Q(εp)=6829 29; %εp decay=0.18 7

¹⁴⁹Er-Q(εp): From 2021Wa16.

Measured p, γ, x, β⁺ in singles and coin. FWHM for protons ≈35 keV.

¹⁴⁸Dy Levels

E(level)	J ^π †	T _{1/2} †
0.0	0 ⁺	3.3 min 2
1677.9	2 ⁺	
1688.4	3 ⁻	
2349.4	5 ⁻	
2427.6	4 ⁺	
2732.3	6 ⁺	
2739.3	7 ⁻	

† From Adopted Levels.

γ(¹⁴⁸Dy)

I_γ normalization: according to 1989Fi01, 24% of ^{149m}Er εp decay goes to ¹⁴⁸Dy g.s., whence %ΣI(γ+ce)(g.s.)=100-24=76; to get the absolute intensity per 100 decays normalization factor, multiply %ΣI(γ+ce)=76 by %εp decay=0.18 7, which gives 0.0014 5.

E _γ	I _γ [#]	E _i (level)	J _i ^π	E _f	J _f ^π	Mult.†	α‡	I _(γ+ce) [#]	Comments
10.5	0.2	1688.4	3 ⁻	1677.9	2 ⁺	[E1]	26.1 4	5	ce(L)/(γ+ce)=0.747 7; ce(M)/(γ+ce)=0.1757 31 ce(N)/(γ+ce)=0.0370 7; ce(O)/(γ+ce)=0.00348 7; ce(P)/(γ+ce)=6.89×10 ⁻⁵ 13 α(L)=20.24 28; α(M)=4.76 7 α(N)=1.003 14; α(O)=0.0943 13; α(P)=0.001867 26
304.6	2	2732.3	6 ⁺	2427.6	4 ⁺	E2	0.0616 9		α(K)=0.0463 6; α(L)=0.01187 17; α(M)=0.00273 4 α(N)=0.000621 9; α(O)=8.17×10 ⁻⁵ 11; α(P)=2.418×10 ⁻⁶ 34
382.9	3	2732.3	6 ⁺	2349.4	5 ⁻	E1	0.00950 13		α(K)=0.00806 11; α(L)=0.001124 16; α(M)=0.0002451 34 α(N)=5.63×10 ⁻⁵ 8; α(O)=8.09×10 ⁻⁶ 11; α(P)=4.33×10 ⁻⁷ 6
389.9	3	2739.3	7 ⁻	2349.4	5 ⁻	E2	0.0298 4		α(K)=0.02331 33; α(L)=0.00504 7; α(M)=0.001147 16 α(N)=0.000262 4; α(O)=3.53×10 ⁻⁵ 5; α(P)=1.267×10 ⁻⁶ 18
661.0	14	2349.4	5 ⁻	1688.4	3 ⁻	E2	0.00759 11		α(K)=0.00625 9; α(L)=0.001046 15; α(M)=0.0002329 33 α(N)=5.35×10 ⁻⁵ 7; α(O)=7.53×10 ⁻⁶ 11; α(P)=3.56×10 ⁻⁷ 5
739.2	6	2427.6	4 ⁺	1688.4	3 ⁻				

Continued on next page (footnotes at end of table)

^{149}Er εp decay (9.6 s) **1989Fi01** (continued) $\gamma(^{148}\text{Dy})$ (continued)

E_γ	$I_\gamma^\#$	$E_i(\text{level})$	J_i^π	E_f	J_f^π	Mult. [†]	α^\ddagger	Comments
749.7	4	2427.6	4 ⁺	1677.9	2 ⁺	E2	0.00567 8	$\alpha(\text{K})=0.00471$ 7; $\alpha(\text{L})=0.000755$ 11; $\alpha(\text{M})=0.0001674$ 23 $\alpha(\text{N})=3.85\times 10^{-5}$ 5; $\alpha(\text{O})=5.46\times 10^{-6}$ 8; $\alpha(\text{P})=2.70\times 10^{-7}$ 4
1677.9	13	1677.9	2 ⁺	0.0	0 ⁺	E2	1.24×10^{-3} 2	$\alpha(\text{K})=0.000938$ 13; $\alpha(\text{L})=0.0001290$ 18; $\alpha(\text{M})=2.81\times 10^{-5}$ 4 $\alpha(\text{N})=6.49\times 10^{-6}$ 9; $\alpha(\text{O})=9.49\times 10^{-7}$ 13; $\alpha(\text{P})=5.42\times 10^{-8}$ 8; $\alpha(\text{IPF})=0.0001400$ 20
1688.4	24	1688.4	3 ⁻	0.0	0 ⁺	E3	2.12×10^{-3} 3	$\alpha(\text{K})=0.001724$ 24; $\alpha(\text{L})=0.000257$ 4; $\alpha(\text{M})=5.67\times 10^{-5}$ 8 $\alpha(\text{N})=1.308\times 10^{-5}$ 18; $\alpha(\text{O})=1.897\times 10^{-6}$ 27; $\alpha(\text{P})=1.034\times 10^{-7}$ 14; $\alpha(\text{IPF})=7.11\times 10^{-5}$ 10

[†] From adopted gammas.

[‡] [Additional information 1.](#)

[#] For absolute intensity per 100 decays, multiply by 0.0014.

Delayed Protons (^{148}Dy)

Particle normalization: $\% \varepsilon + \% \beta^+ = 96.5$ 7; $\% \varepsilon \text{p} = 0.18$ 7 (**1989Fi01**).

$E(^{148}\text{Dy})$	$I(\text{p})^\dagger$
0.0	72
1677.9	3
1688.4	7
2349.4	5
2427.6	6
2732.3	4
2739.3	3

[†] For absolute intensity per 100 decays, multiply by 0.0018 7.

^{149}Er ϵp decay (9.6 s) 1989Fi01

Decay Scheme

Intensities: $I_{(\gamma+ce)}$ per 100 parent decays

Legend

- $I_{\gamma} < 2\% \times I_{\gamma}^{\max}$
- $I_{\gamma} < 10\% \times I_{\gamma}^{\max}$
- $I_{\gamma} > 10\% \times I_{\gamma}^{\max}$

