

Adopted Levels

Type	Author	Citation	History	Literature Cutoff Date
Full Evaluation	F. G. Kondev	ENSDF		20-Feb-2017

$Q(\beta^-)=7.9\times10^3$ 4; $S(n)=5.3\times10^3$ 6; $S(p)=10699$ (syst) 529; $Q(\alpha)=-3.8\times10^3$ 4
 $S(2n)=9.2\times10^3$ 6; $S(2p)=24638$ (syst) 529; $Q(\beta^-n)=3.5\times10^3$ 4 [2017Wa10](#)

Additional information 1.

[2017Wu04](#): The ^{151}La nuclide was produced at the RIBF-RIKEN facility using the $^9\text{Be}(^{238}\text{U},\text{F})$ reaction at $E=345$ MeV/nucleon.

Two experiments, optimized for the transmission of ^{158}Nd and ^{170}Dy ions, were carried out with average beam intensities of 7 pnA and 12 pnA, respectively. The identification of the nuclide of interest was made in the BigRIPS separator by determining the atomic number and the mass-to-charge ratio of the ion using the TOF- $B\rho-\Delta E$ method. The reaction products were transported through the ZeroDegree Spectrometer and implanted into the beta-counting system WAS3ABi that was surrounded by the EURICA array comprising of 84 HPGe detectors. The typical implantation rate was 100 ions/s. Measured: implanted ion- β^- -t, implanted ion- β^- - γ -t and implanted ions- γ -t correlations. Deduced: $T_{1/2}$.

[1994Be24](#): Identification in reaction: $\text{Pb}(^{238}\text{U},\text{F})$ at 750 MeV/nucleon. Residual products Fragment Recoil Separator (FRS), time-of-flight technique. A total of 106 counts were observed corresponding to a cross section of $>52 \mu\text{b}$.

 ^{151}La Levels

E(level)	J^π	$T_{1/2}$	Comments
0.0	(1/2 ⁺)	0.457 s +30-18	% β^- =100; % β^- n=? % β^- : Only β^- decay mode is expected. J^π : From systematics of known quasiparticle states in neighboring nuclei and the proposed configuration (by the evaluator). The assignment is tentative. $T_{1/2}$: From 2017Wu04 , using a fit to the implanted ion- β^- -t spectrum using the least-squares and maximum-likelihood methods. The data analysis included contributions from the parent, daughter and ground-daughter decays, as well as a constant background. configuration: $\pi 1/2[420]$ ($d_{5/2}$) Nilsson orbital, expected from systematics of well-deformed nuclei in the region (by the evaluator). The assignment is tentative.