

**<sup>169</sup>Hf ε+β<sup>+</sup> decay (3.24 min) 1973Me09,1973FoYE**

Type	Author	Citation	Literature Cutoff Date
Full Evaluation	M. Shamsuzzoha Basunia	NDS 209,1 (2026)	1-Oct-2025

Parent: <sup>169</sup>Hf: E=0.0; J<sup>π</sup>=5/2<sup>-</sup>; T<sub>1/2</sub>=3.24 min 5; Q(ε)=3366 28; %ε+%β<sup>+</sup> decay=100

<sup>169</sup>Hf-J<sup>π</sup>,T<sub>1/2</sub>: from <sup>169</sup>Hf Adopted Levels.

<sup>169</sup>Hf-Q(ε+β<sup>+</sup>): from 2021Wa16.

1973Me09: sources from <sup>170</sup>Yb(<sup>3</sup>He,4n); Yb oxide targets enriched to 67% in <sup>170</sup>Yb; measured E<sub>γ</sub>, I<sub>γ</sub> (Compton-suppressed Ge(Li) (FWHM=1.9 keV at 1332 keV), surface-barrier Ge(Li) (FWHM=0.8 keV at 122 keV)), Ice (Si(Li)), half-life of <sup>169</sup>Hf.

1973FoYE: sources from <sup>168</sup>Yb(α,3n) and <sup>170</sup>Yb(α,5n); measured E<sub>γ</sub>, I<sub>γ</sub> (Ge(Li), FWHM=1.4 keV at 100 keV, 2.3 keV at 1 MeV); measured E<sub>γ</sub>, I<sub>γ</sub> γγ coin; placed 72.9γ and 68.4γ.

1969Ar23: Hf sources from <sup>169</sup>Ta parent decay; measured E<sub>γ</sub>, I<sub>γ</sub>, γ(t), γγ coin using coaxial Ge(Li) and NaI(Tl) detectors, and β<sup>+</sup> data using a 2 cm x 2 cm anthracene crystal (14% energy resolution for 624-keV ce line from <sup>137</sup>Cs), also I(γ<sup>±</sup>).

1970At01: Measured E<sub>γ</sub>, I<sub>γ</sub> and half-life of <sup>169</sup>Hf.

1970Ch17: Measured E<sub>γ</sub>, I<sub>γ</sub>, I(γ<sup>±</sup>), and half-life of <sup>169</sup>Hf.

1975Gr44: Measured E<sub>γ</sub>, I<sub>γ</sub> and half-life of <sup>169</sup>Hf.

The decay scheme is tentative and incomplete. The measured I(K x ray) cannot be corrected for K conversion of possible 71γ and 85γ because their intensities are unknown; if significant, this would lower εK/β<sup>+</sup> and, hence, the calculated ε+β<sup>+</sup> feeding to the 493 level. However, a very large change in feeding to the 493 level would be required to invalidate the conclusion that this is an allowed, unhindered transition. Feeding to the (5/2<sup>-</sup>) 43 level may be significant, but is unknown. There may also exist feeding to levels with E>493 that are as yet unknown.

<sup>169</sup>Lu Levels

E(level) <sup>†</sup>	J <sup>π</sup> <sup>‡</sup>	T <sub>1/2</sub> <sup>‡</sup>	Comments
0.0	7/2 <sup>+</sup>	34.06 h 5	
29.0 5	1/2 <sup>-</sup>	160 s 10	
42.9 6	(5/2 <sup>-</sup> )		
97.4 5	(1/2 <sup>+</sup> )		log f <sup>A<sub>t</sub></sup> >8.5 implies %ε+%β <sup>+</sup> feeding<0.4 to this level.
113.8 6	(3/2 <sup>+</sup> )		log ft>5.9 implies %ε+%β <sup>+</sup> <5 to this level.
123.49 15	(9/2 <sup>+</sup> ) <sup>+</sup>		log f <sup>A<sub>t</sub></sup> >8.5 implies %ε+%β <sup>+</sup> feeding<0.4 to this level, consistent with intensity balance at this level.
186.7 6	(5/2 <sup>+</sup> )		
492.89 10	7/2 <sup>-</sup>		

<sup>†</sup> From least-squares fit to E<sub>γ</sub>.

<sup>‡</sup> From Adopted Levels.

ε,β<sup>+</sup> radiations

Additional information 1.

E(decay)	E(level)	Iβ <sup>+</sup> <sup>†</sup>	Iε <sup>†</sup>	Log ft	I(ε+β <sup>+</sup> ) <sup>†</sup>	Comments
(2873 28)	492.89	>12.1	>75.9	<4.6	>88	av Eβ=834 12; εK=0.714 5; εL=0.1125 8; εM+=0.03576 22 E(decay): 2872 200 from measured Eβ+=1850 200 (1969Ar23). I(ε+β <sup>+</sup> ): estimated in 1973Me09 from I(K x ray)=86 5 and I(γ <sup>±</sup> )=33.0 15, relative to Iγ=100 for 492.9γ (1973Me09).
(3179 28)	186.7	<1.0	<4.0	>5.9	<5	av Eβ=970 12; εK=0.659 6; εL=0.1035 9; εM+=0.03288 25 I(ε+β <sup>+</sup> ): 6.8 from intensity imbalance; however, uncertainty may be large (1973FoYE do not state uncertainty in I(73γ)). <5% feeding is expected based on log ft>5.9 for first-forbidden transition.

Continued on next page (footnotes at end of table)

$^{169}\text{Hf}$   $\varepsilon+\beta^+$  decay (3.24 min)  $^{1973}\text{Me09},^{1973}\text{FoYE}$  (continued) $\varepsilon, \beta^+$  radiations (continued)

<u>E(decay)</u>	<u>E(level)</u>	<u><math>I\beta^+</math></u> †	<u><math>I\varepsilon</math></u> †	<u>Log <math>ft</math></u>	<u><math>I(\varepsilon+\beta^+)</math></u> †	Comments
(3323 ‡ 28)	42.9					
(3366 ‡ 28)	0.0	<1.2	<3.7	>6.0	<5	av $E\beta=1053$ 13; $\varepsilon\text{K}=0.622$ 6; $\varepsilon\text{L}=0.0976$ 10; $\varepsilon\text{M}+=0.03099$ 26 $I(\varepsilon+\beta^+)$ : from $\log ft > 5.9$ for first-forbidden transition.

† Absolute intensity per 100 decays.

‡ Existence of this branch is questionable.

<sup>169</sup>Hf ε+β<sup>+</sup> decay (3.24 min) **1973Me09,1973FoYE (continued)**

γ(<sup>169</sup>Lu)

I<sub>γ</sub> normalization: from ΣI(γ+ce) 94 6 from 492.9 keV level, based on the estimated total of >88% to satisfy K/β=5.2 14-4 from measured I(K x ray)=86 5 and I(γ<sup>±</sup>)=33.0 15 relative to I(493γ)=100 (1973Me09). This normalization should be considered to be highly tentative. Other I(γ<sup>±</sup>)=27 (1969Ar23), 82 20 relative to I(493γ)=100 (1970Ch17).  
I<sub>γ</sub>(K x ray)=86 5, I(γ<sup>±</sup>)=33.0 15, relative to I<sub>γ</sub>=100 for 492.9γ (1973Me09).

<u>E<sub>γ</sub><sup>†</sup></u>	<u>I<sub>γ</sub><sup>†α</sup></u>	<u>E<sub>i</sub>(level)</u>	<u>J<sub>i</sub><sup>π</sup></u>	<u>E<sub>f</sub></u>	<u>J<sub>f</sub><sup>π</sup></u>	<u>Mult.<sup>‡</sup></u>	<u>α&amp;</u>	<u>I<sub>(γ+ce)</sub><sup>a</sup></u>	<u>Comments</u>
(14.1 @)		42.9	(5/2 <sup>-</sup> )	29.0	1/2 <sup>-</sup>				
(16.4 @)		113.8	(3/2 <sup>+</sup> )	97.4	(1/2 <sup>+</sup> )			≈3.7	I <sub>(γ+ce)</sub> : since negligible ε+β <sup>+</sup> feeding of the 97 level is expected (ΔJ <sup>π</sup> =2, no), the level scheme implies I(γ+ce)(16γ)≈I(γ+ce)(68γ)=3.7.
(29.0 5)		29.0	1/2 <sup>-</sup>	0.0	7/2 <sup>+</sup>	E3	9.4×10 <sup>4</sup> 10		α(L)=6.8×10 <sup>4</sup> 7; α(M)=2.08×10 <sup>4</sup> 24 α(N)=4.9×10 <sup>3</sup> 6; α(O)=5.4×10 <sup>2</sup> 6; α(P)=0.346 35 E <sub>γ</sub> ,Mult.: from Adopted Gammas.
68.4 @ 1	1.9#	97.4	(1/2 <sup>+</sup> )	29.0	1/2 <sup>-</sup>	(E1) @	0.931 13		α(K)=0.754 11; α(L)=0.1379 20; α(M)=0.0311 5 α(N)=0.00716 10; α(O)=0.000951 14; α(P)=3.88×10 <sup>-5</sup> 6
(70.9 @ 2)		113.8	(3/2 <sup>+</sup> )	42.9	(5/2 <sup>-</sup> )	[E1]	0.852 13		α(K)=0.691 11; α(L)=0.1248 20; α(M)=0.0282 5 α(N)=0.00648 10; α(O)=0.000864 14; α(P)=3.56×10 <sup>-5</sup> 6
72.9 @ 2	0.7#	186.7	(5/2 <sup>+</sup> )	113.8	(3/2 <sup>+</sup> )	[M1,E2]	10.6 15		α(K)=4.6 30; α(L)=4.6 34; α(M)=1.1 9 α(N)=0.26 20; α(O)=0.032 23; α(P)=3.4×10 <sup>-4</sup> 23
(84.8 @ 2)		113.8	(3/2 <sup>+</sup> )	29.0	1/2 <sup>-</sup>	[E1]	0.538 8		α(K)=0.440 7; α(L)=0.0760 12; α(M)=0.01712 26 α(N)=0.00395 6; α(O)=0.000534 8; α(P)=2.317×10 <sup>-5</sup> 35
123.6 2	5.0 5	123.49	(9/2) <sup>+</sup>	0.0	7/2 <sup>+</sup>	M1,E2	1.76 25		α(K)=1.1 5; α(L)=0.47 21; α(M)=0.11 6 α(N)=0.026 13; α(O)=0.0034 14; α(P)=8 E <sub>γ</sub> : others: 123.2 (1970At01), 123 (1970Ch17), 123 (1975Gr44).
369.5 2	11.7 8	492.89	7/2 <sup>-</sup>	123.49	(9/2) <sup>+</sup>	[E1]	0.01253 18		I <sub>γ</sub> : weighted average of 4.6 5 (1973Me09) and 5.6 6 (1970Ch17). Others: 8 (1970At01), 7 (1975Gr44). Mult.: from 1.16<α(exp)<5.0 (1973Me09 – scaled from 1.17<α(exp)<5.1) with α <sub>1</sub> =0.0814(Bricc)/0.0823(1973Me09) for the 198.84 keV 3 4 <sup>+</sup> to 2 <sup>+</sup> transition in 168Yb. α(K)=0.01054 15; α(L)=0.001548 22; α(M)=0.000346 5 α(N)=8.11×10 <sup>-5</sup> 11; α(O)=1.173×10 <sup>-5</sup> 16; α(P)=6.64×10 <sup>-7</sup> 9 E <sub>γ</sub> : others: 369.3 (1970At01), 368 (1970Ch17), 369 (1975Gr44).
492.86 10	100	492.89	7/2 <sup>-</sup>	0.0	7/2 <sup>+</sup>	E1	0.00651 9		I <sub>γ</sub> : weighted average of 11.6 10 (1973Me09) and 11.9 12 (1970Ch17). Others: 11 (1970At01), 13 (1975Gr44). α(K)=0.00550 8; α(L)=0.000792 11; α(M)=0.0001765 25 α(N)=4.14×10 <sup>-5</sup> 6; α(O)=6.04×10 <sup>-6</sup> 8; α(P)=3.53×10 <sup>-7</sup> 5

<sup>169</sup>Hf  $\epsilon+\beta^+$  decay (3.24 min) 1973Me09,1973FoYE (continued)

$\gamma(^{169}\text{Lu})$  (continued)

<u><math>E_\gamma</math></u> <sup>†</sup>	<u><math>E_i</math>(level)</u>	Comments
		$E_\gamma$ : others: 492.3 (1970At01), 491 (1970Ch17). $I_\gamma$ : same value in 1970At01, 1970Ch17, and 1975Gr44. $\alpha(\text{K})_{\text{exp}}=0.004$ I (1973Me09).

<sup>†</sup> From 1973Me09, except as noted.

<sup>‡</sup> From  $\alpha(\text{K})_{\text{exp}}$ , as deduced from simultaneous measurement of  $I_{\text{ce}}$  and  $I_\gamma$ , except where noted (detector calibration assumes  $\alpha(\text{L})=0.0814$  (E2 theory) for 198.8 $\gamma$  in <sup>168</sup>Yb) (1973Me09). Same in the adopted dataset.

# From 1973FoYE; uncertainty unstated by authors.

@ From Adopted Gammas.

& Additional information 2.

<sup>a</sup> For absolute intensity per 100 decays, multiply by  $\approx 0.82$ .

**$^{169}\text{Hf } \epsilon + \beta^+ \text{ decay (3.24 min) } 1973\text{Me09,1973FoYE}$**

Decay Scheme

Legend

- $I_\gamma < 2\% \times I_\gamma^{\text{max}}$
- $I_\gamma < 10\% \times I_\gamma^{\text{max}}$
- $I_\gamma > 10\% \times I_\gamma^{\text{max}}$
- - - - -  $\gamma$  Decay (Uncertain)

Intensities:  $I_{(\gamma+ce)}$  per 100 parent decays

