

**$^{12}\text{C}(^7\text{Li},\text{d})$  2008Cr03**

Type	Author	Citation	Literature Cutoff Date
Full Evaluation	C. G. Sheu, J. H. Kelley, J. Purcell	ENSDF	5-Aug-2021

**1971Sc21:** The reactions  $^{12}\text{C}(^7\text{Li},\text{d})$  and  $^{13}\text{C}(^7\text{Li},\text{t})$  were studied at  $E_{\text{cm}}=13.3$  MeV using the lithium beam, from the E(n)-tandem-van-de-Graaff-Accelerator of the Max-Planck-Institut für Kernphysik at Heidelberg, impinged on a  $^{13}\text{C}$  target (50%  $^{13}\text{C}$ , 50%  $^{12}\text{C}$  and  $^{16}\text{O}$ ). The reactions products were identified by the  $\Delta E$ -E information. The overall resolutions for deuterons was about 90 keV.

The integrated cross sections  $\sigma_{\text{int}}$  were measured in both reactions. Spin assignments were extracted from  $\sigma_{\text{int}}$  in the reaction  $^{12}\text{C}(^7\text{Li},\text{d})$  and a modified DWBA code was used to analyze the reaction  $^{13}\text{C}(^7\text{Li},\text{t})$ . Energy levels and  $J^\pi$  values of  $^{17}\text{O}$  were deduced.

**1982Ta23:**  $^{12}\text{C}(^7\text{Li},\text{d})$ ,  $E=36,32,28$  MeV; measured yield vs particle energy,  $\sigma(\theta)$ , fusion  $\sigma$ , breakup  $\sigma$  vs  $E$ ; deduced reaction mechanism. Optical, simple breakup model analyses.

**2008Cr03:** XUNDL dataset compiled by McMaster, 2008.

$E=34$  MeV beam provided by FN tandem accelerator at Florida State. Detected charged particles using two  $\Delta E$ -E Si telescopes. Measured absolute cross sections and  $\sigma(\theta)$ . DWBA analysis assuming a  $^5\text{He}$  cluster transfer. FWHM=110 keV.

**Theory:**

**1987Ar13:**  $^{12}\text{C}(^7\text{Li},\text{d})$ ,  $E(\text{cm})=7.4,9.4$  MeV; calculated (np), d emission  $\sigma$ , residual production  $\sigma(E)$  ratio. Hauser-Feshbach theory.

 $^{17}\text{O}$  Levels

E(level) <sup>‡</sup>	$J^\pi$ <sup>‡</sup>	L <sup>‡</sup>	C <sup>2</sup> S <sup>‡#</sup>	Comments
0 <sup>@</sup>	5/2 <sup>+</sup>			E(level), $J^\pi$ : See also (1971Sc21).
870 <sup>@</sup>	1/2 <sup>+</sup>			E(level), $J^\pi$ : See also (1971Sc21).
3060 <sup>@</sup>	1/2 <sup>-</sup>			E(level), $J^\pi$ : See also (1971Sc21).
3840 <sup>@</sup>	5/2 <sup>-</sup>			E(level), $J^\pi$ : See also (1971Sc21).
4550 <sup>@</sup>	3/2 <sup>-</sup>			E(level), $J^\pi$ : See also (1971Sc21).
5080				E(level): from (1971Sc21).
5220 <sup>@</sup>	9/2 <sup>-</sup>			$J^\pi$ : 7/2 (1971Sc21).
5380 <sup>@</sup>	3/2 <sup>-</sup>			
5700 <sup>@&amp;</sup>				E(level): See also doublet 5.69-MeV and 5.72-MeV (1971Sc21).
5900 <sup>@&amp;</sup>				E(level): See also doublet 5.87-MeV: $J^\pi=5/2$ and 5.94-MeV: $J^\pi=1/2$ (1971Sc21).
6360 <sup>@</sup>	1/2 <sup>+</sup>			
6860 13	5/2 <sup>+</sup>	3	0.53	Configuration= $^{12}\text{C}_{\text{g.s.}}+1\text{p}_{1/2}^2, 1\text{d}_{5/2}^3-(3\text{p}-2\text{h})$ . E(level): See also 6.87-MeV: $J^\pi=7/2$ (1971Sc21).
6990	5/2			E(level), $J^\pi$ : from (1971Sc21).
7170 <sup>@</sup>	5/2 <sup>-</sup>			E(level), $J^\pi$ : See also (1971Sc21).
7380 <sup>@&amp;</sup>	9/2			$J^\pi$ : from (1971Sc21).
7580 13	7/2 <sup>+</sup>	5	0.59	Configuration= $^{12}\text{C}_{\text{g.s.}}+1\text{p}_{1/2}^2, 1\text{d}_{5/2}^3-(3\text{p}-2\text{h})$ . E(level): See also 7.56-MeV: $J^\pi=9/2$ (1971Sc21).
7760 <sup>@</sup>	11/2 <sup>-</sup>			E(level): See also triplet 7.69-MeV: $J^\pi=3/2$ , 7.71-MeV: $J^\pi=7/2$ and 7.72-MeV: $J^\pi=3/2$ (1971Sc21).
8470 13	9/2 <sup>+</sup>	3	1.06	Configuration= $^{12}\text{C}_{\text{g.s.}}+1\text{p}_{1/2}^2, 1\text{d}_{5/2}^3-(3\text{p}-2\text{h})$ . E(level): See also triplet 8.40-MeV: $J^\pi=5/2$ , 8.47-MeV: $J^\pi=9/2$ and 8.50-MeV: $J^\pi=5/2$ (1971Sc21).
8680 <sup>@</sup>	3/2 <sup>-</sup>			
8900 <sup>@</sup>				E(level): triplet. See also triplet 8.87-MeV: $J^\pi=3/2$ , 8.88-MeV: $J^\pi=7/2$ and 8.95-MeV: $J^\pi=7/2$ (1971Sc21).
9190 <sup>@</sup>				E(level): quadruplet. E(level): See also (1971Sc21).

Continued on next page (footnotes at end of table)

$^{12}\text{C}(^7\text{Li,d})$  2008Cr03 (continued) $^{17}\text{O}$  Levels (continued)

E(level) <sup>‡</sup>	J <sup>π</sup> <sup>‡</sup>	Γ <sup>†</sup>	L <sup>‡</sup>	C <sup>2</sup> S <sup>‡#</sup>	Comments
9490 @	5/2 <sup>-</sup>				E(level): See also (1971Sc21).
9710 @	7/2 <sup>+</sup>				E(level): See also (1971Sc21).
9870 @&					E(level): See also doublet 9.88-MeV and 9.95-MeV (1971Sc21).
10690 26		<40 keV			E(level): See also 10.78-MeV (1971Sc21).
11040 @					
11240 @					
11820 13	7/2 <sup>+</sup>		5	0.96	Configuration= $^{12}\text{C}_{\text{g.s.}} + 1\text{p}_{1/2}^0, 1\text{d}_{5/2}^5$ -(5p-4h). E(level): See also 11.88-MeV (1971Sc21).
12000 26	9/2 <sup>+</sup>	<50 keV	3	0.56	Configuration= $^{12}\text{C}_{\text{g.s.}} + 1\text{p}_{1/2}^0, 1\text{d}_{5/2}^5$ -(5p-4h).
12220 26	7/2 <sup>-</sup>		2	2.16	Configuration= $^{12}\text{C}_{\text{g.s.}} + 1\text{p}_{1/2}^3, 1\text{d}_{5/2}^2$ -(2p-1h).
12420 26	9/2 <sup>+</sup>	<50 keV	5	0.77	Configuration= $^{12}\text{C}_{\text{g.s.}} + 1\text{p}_{1/2}^0, 1\text{d}_{5/2}^5$ -(5p-4h).
12760 26		<70 keV			Γ: Estimated value based on the FWHM of the peak in the $^{12}\text{C}(^7\text{Li,d})$ reaction (2008Cr03).
13060 26					
13580 26					
14550 26					
14720 26					
14880 26					
15070 26					
15620 26					
15800 26					

<sup>†</sup> From (1986Sm10) except where noted. Width measurement limited by detector resolution of the  $^{12}\text{C}(^6\text{Li,p})$  measurement (2008Cr03).

<sup>‡</sup> From (2008Cr03) except where noted. Some concern is raised over the small number of nodes used in the DWBA analysis for some cases (priv. comm. J. Millener).

# Assuming  $^5\text{He}$  cluster, assumed configurations are listed.

@ From Fig. 1 of (2008Cr03).

& Doublet.