

$^{18}\text{O}(\alpha, \alpha)$ 1969De06

Type	Author	Citation	History	Literature Cutoff Date
Full Evaluation	C. G. Sheu, J. H. Kelley, J. Purcell	ENSDF		5-Aug-2021

1969De06: An $E(^3\text{He})=16$ MeV beam from the Heidelberg E(n) Tandem Van de Graaff accelerator bombarded a target containing $10 \mu\text{g}/\text{cm}^2$ of ^{18}O and $\approx 6 \mu\text{g}/\text{cm}^2$ of ^{16}O . A broad range magnetic spectrograph was used to analyze α -particles. The α -particle spectrum was obtained at $\theta=5^\circ$ and the absolute cross sections were determined with an accuracy of 25%. Eight analogue T=3/2 excited states in ^{17}O were identified. The l -transfer values and spectroscopic factors were also deduced for four of these states.

1970Mc02: Branching ratios were measured for the decays of the lowest T=3/2 levels of ^{17}F and ^{17}O to the ground state and unresolved 6.05- and 6.13-MeV levels of ^{16}O . The experiment was performed by bombarding a nickel oxide target (98% ^{18}O enriched) with an $E=12$ ^3He ion beam. Alpha particles were detected at $\theta=10^\circ$ with a double-focusing magnetic spectrometer. The branching ratios for transition $^{17}\text{O}^*(11.08 \text{ MeV}) \rightarrow ^{16}\text{O}_{\text{g.s.}}$ and $^{17}\text{O}^*(11.08 \text{ MeV}) \rightarrow ^{16}\text{O}^*(6.05+6.13 \text{ MeV})$ are (0.91 15) and (0.05 2), respectively. The ratios of the reduced widths (θ^2) decaying to ^{16}O levels, $\Theta^2(\text{g.s.})/\theta^2(6.05)=3.4$ 14 and $\Theta^2(\text{g.s.})/\theta^2(6.13)=0.32$ 14 were also deduced. The width of $^{17}\text{O}^*(11.08 \text{ MeV})$ state is <20 keV (D.C. Hensly, Ph.D. thesis, Caltech (1969) unpublished).

1973Ad02: $^{18}\text{O}(\alpha, n\alpha)$, $E=12$ MeV; measured $\sigma(E_n, E_\alpha, \theta(\alpha), \theta(n))n\alpha$ -coin. ^{17}O deduced level-width(n).

 ^{17}O Levels

<u>E(level)[†]</u>	<u>J^π[‡]</u>	<u>Γ</u>	<u>L[‡]</u>	<u>C²S[#]</u>	<u>Comments</u>
11082 6	(1/2) ⁻	5 keV 1	1	0.49	Γ: from (McDonald et al., Bull. Amer. Phys. Soc. 16, 489 (1971) $^{13}\text{C}(\alpha, n)$). See also <20 keV (D.C. Hensly, Ph.D. thesis, Caltech (1969) unpublished). $\Gamma_{n0}/\Gamma=91$ 15 and $\Gamma_{n(1+2)}/\Gamma=0.05$ 2 were deduced in (1973Ad02). Also $\theta^2(\text{g.s.})/\theta^2(6.13)=0.31$ 14 (1973Ad02); these compare with $\theta^2(\text{g.s.})/\theta^2(6.05)=3.4$ 14 and $\theta^2(\text{g.s.})/\theta^2(6.13)=0.32$ 14 (1970Mc02). The value $\Gamma_{\alpha0}=0.3$ keV is deduced using the measured (1973Ad02) neutron branching ratios and the width from McDonald; however in the present evaluation we adopt a different $\Gamma=2.4$ keV 3 and $\Gamma_{n0}/\Gamma=81$ 6. This changes the interpretation.
12471 5	(3/2) ⁻		1	0.27	
12950 8	1/2 ⁺		0	0.096	
12994 8					
13640 5	(5/2) ⁺		2	0.39	
14219 8					
14282 12					
15101 8					

[†] From (1969De06); T=3/2 states.

[‡] From (1969De06).

[#] Calculated assuming $C^2S=4$ for $^{15}\text{O}^*(6.18 \text{ MeV})$ in $^{16}\text{O}(\alpha, \alpha)^{15}\text{O}$.