

Adopted Levels

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	Coral M. Baglin, E. A. Mccutchan		NDS 151, 334 (2018)	30-Jun-2018

$Q(\beta^-)=4330$  SY;  $S(n)=4600$  SY;  $S(p)=10750$  SY;  $Q(\alpha)=-1800$  SY    [2017Wa10](#)  
 $\Delta Q(\beta^-)=670$ ;  $\Delta S(n)=360$ ;  $\Delta S(p)=500$ ;  $\Delta Q(\alpha)=420$  ([2017Wa10](#)).  
 $S(2n)=10730$  (syst) 420;  $S(2p)=20610$  (syst) 590 ([2017Wa10](#)).  
[2010So03](#): possible production of  $^{171}\text{Dy}$  in  $^{170}\text{Er}(^{82}\text{Se},X)$ ,  $E=460$  MeV/nucleon. Measured yield of beam-like fragment  $^{81}\text{Kr}$  which implies production of complementary "target-like fragment"  $^{171}\text{Dy}$ .  
[2012Ku26](#):  $^{171}\text{Dy}$  produced and identified in  $^9\text{Be}(^{238}\text{U},F)$ , with  $E=1$  GeV/nucleon followed by separation using the Fragment Separator (FRS). Particle identification using event-by-event in-flight analysis of time-of-flight, energy loss and magnetic rigidity (tof- $\Delta E'$ - $B\rho$ ) measurements. Time-of-flight was measured using two plastic scintillation detectors, energy loss or deposit by ionization chambers (MUSIC), and magnetic rigidity by four time-projection chambers (TPC), which also provided energy deposit information. Measured production cross section.  
[2017Wu04](#):  $^{171}\text{Dy}$  produced using the  $^9\text{Be}(^{238}\text{U},F)$  reaction at  $E=345$  MeV/nucleon and identification made in the BigRIPS separator by measuring the atomic number and the mass-to-charge ratio of the ion using the tof- $B\rho$ - $\Delta E$  method. Reaction products were transported through the ZeroDegree Spectrometer and implanted into the beta-counting system WAS3ABi which was surrounded by the EURICA array consisting of 84 HPGe detectors. Measured implanted ion- $\beta^-(t)$ , implanted ion- $\beta^--\gamma(t)$  and implanted ions- $\gamma(t)$ ; deduced  $T_{1/2}$ .

 $^{171}\text{Dy}$  Levels

E(level)	$T_{1/2}$	Comments
0.0	4.1 s 4	$\% \beta^- = 100$ $\% \beta^-$ : Only $\beta^-$ decay mode is expected. $J^\pi$ : $7/2^-$ from systematics of known quasiparticle states in neighboring nuclei. $T_{1/2}$ : from <a href="#">2017Wu04</a> , using a fit to the implanted ion- $\beta^-(t)$ spectrum applying the least-squares and maximum-likelihood methods; data analysis included contributions from the parent, daughter and ground-daughter decays, as well as a constant background. Value is reported as 4.07 s 40 in <a href="#">2017Wu04</a> . Production $\sigma$ (at 1 GeV/nucleon)=441 nb 18 ( <a href="#">2012Ku26</a> ).