

$^{174}\text{Hf}(\text{n},\gamma) \text{E=th}$ 1971AJ01

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	M. Shamsuzzoha Basunia		NDS 102, 719 (2004)	1-Jun-2004

Enriched target (14%). Ge(Li) pair and anti-Compton spectrometers. $\approx 30\%$ of capture spectrum originates from ^{175}Hf .

 ^{175}Hf Levels

E(level)	J^π [†]	$T_{1/2}$	E(level)	J^π [†]	E(level)	J^π [†]
0.0 [‡]	5/2 ⁻	70 d 2	373.9 17		1455.7 9	(1/2,3/2)
81.2 [‡] 8	7/2 ⁻		866.9 [@] 7	(1/2 ⁻)	1642.8 13	(1/2,3/2)
125.9 [#] 7	1/2 ⁻		910.3 [@] 9	(3/2 ⁻)	(6708.9 7)	1/2 ⁺
196.5 [#] 7	3/2 ⁻		997.7 ^{&} 9	(1/2 ⁻ ,3/2 ⁻)		
361? 3			1375.1 11	(1/2,3/2)		

[†] From Adopted Levels.

[‡] 5/2(512) band.

[#] 1/2(521) band.

[@] 1/2(510)? band.

[&] 3/2(521)? band.

 $\gamma(^{175}\text{Hf})$

E_γ	I_γ [†]	$E_i(\text{level})$	J_i^π	E_f	J_f^π
81.5 ^{&}		81.2	7/2 ⁻	0.0	5/2 ⁻
115.7 20	0.87	196.5	3/2 ⁻	81.2	7/2 ⁻
125.7 10	18.6	125.9	1/2 ⁻	0.0	5/2 ⁻
177.4 15	1.2	373.9		196.5	3/2 ⁻
197.0 [‡] 20	3.9	196.5	3/2 ⁻	0.0	5/2 ⁻
^x 342.6 10	0.56				
^x 420.0 10	0.20				
^x 439.5 15	1.1				
^x 441.5 15	0.48				
588.0 15	0.55	1455.7	(1/2,3/2)	866.9	(1/2 ⁻)
^x 665.0 [#] 20	0.3				
670.5 10	5.2	866.9	(1/2 ⁻)	196.5	3/2 ⁻
714.1 10	0.89	910.3	(3/2 ⁻)	196.5	3/2 ⁻
^x 745.0 10	0.30				
784.1 10	0.98	910.3	(3/2 ⁻)	125.9	1/2 ⁻
^x 813.0 10	1.0				
^x 859.4 15	0.83				
866.7 10	2.2	866.9	(1/2 ⁻)	0.0	5/2 ⁻
997.7 10	1.1	997.7	(1/2 ⁻ ,3/2 ⁻)	0.0	5/2 ⁻
^x 1072.0 20	1.0				
^x 1095.0 20	1.2				
1374.9 [@] 15	0.67 [@]	1375.1	(1/2,3/2)	0.0	5/2 ⁻
1374.9 [@] 15	0.67 [@]	1455.7	(1/2,3/2)	81.2	7/2 ⁻
1446.0 15	0.44	1642.8	(1/2,3/2)	196.5	3/2 ⁻
1455.4 15	0.88	1455.7	(1/2,3/2)	0.0	5/2 ⁻
1644.0 25	1.0	1642.8	(1/2,3/2)	0.0	5/2 ⁻
5066.5 25	0.91	(6708.9)	1/2 ⁺	1642.8	(1/2,3/2)
5252.6 15	0.72	(6708.9)	1/2 ⁺	1455.7	(1/2,3/2)

Continued on next page (footnotes at end of table)

$^{174}\text{Hf}(\text{n},\gamma)$ E=th **1971AJ01** (continued) $\gamma(^{175}\text{Hf})$ (continued)

E_γ	$I_\gamma^\dagger$	$E_i(\text{level})$	J_i^π	E_f	J_f^π	E_γ	$I_\gamma^\dagger$	$E_i(\text{level})$	J_i^π	E_f	J_f^π
5333.6 15	0.39	(6708.9)	$1/2^+$	1375.1	(1/2,3/2)	6348 [#] 3		(6708.9)	$1/2^+$	361?	
5711.1 20	0.31	(6708.9)	$1/2^+$	997.7	(1/2 ⁻ ,3/2 ⁻)	6512.1 10	1.5	(6708.9)	$1/2^+$	196.5	3/2 ⁻
5799.1 20	0.48	(6708.9)	$1/2^+$	910.3	(3/2 ⁻)	6582.8 10	0.92	(6708.9)	$1/2^+$	125.9	1/2 ⁻
5842.2 10	5.5	(6708.9)	$1/2^+$	866.9	(1/2 ⁻)						

[†] Primary γ 's normalized to $I_\gamma(6512\gamma)=1.5$, secondary γ 's normalized to $I_\gamma(670.5\gamma)=5.2$ based on absolute intensity measurements for natural Hf from [1969Gr41](#).

[‡] Doublet.

[#] Uncertain transition.

@ Multiply placed with undivided intensity.

& Placement of transition in the level scheme is uncertain.

^x γ ray not placed in level scheme.

$^{174}\text{Hf}(n,\gamma) \text{E=th}$ 1971A101

Level Scheme

Intensities: Relative I_γ
& Multiply placed: undivided intensity given

Legend

- $\longrightarrow$ $I_\gamma < 2\% \times I_\gamma^{\max}$
 $\longrightarrow$ $I_\gamma < 10\% \times I_\gamma^{\max}$
 $\longrightarrow$ $I_\gamma > 10\% \times I_\gamma^{\max}$
 $\longrightarrow$ γ Decay (Uncertain)

