

(HI,xn γ) 1981Ho29,1981Ma28,2008An01

Type	Author	History Citation	Literature Cutoff Date
Full Evaluation	F. G. Kondev	NDS 201,346 (2025)	21-Jan-2025

1981Ho29: $^{197}\text{Au}(^{14}\text{N},5n\gamma)$ reaction with $E(^{14}\text{N})=80\text{--}94$ MeV and $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$ reaction on enriched (97%) target with $E(^{16}\text{O})=85\text{--}110$ MeV. The targets were backed with 8-10 mg/cm² thick ^{208}Pb layers; Detectors: Ge(Li); Measured: excitation functions, $E\gamma$, $I\gamma$, $\gamma\gamma(t)$ coin, $T_{1/2}$; Deduced: level scheme. No longer-lived isomers ($T_{1/2}<0.05$ s) were observed in a pulsed beam experiment.

1981Ma28: $^{198}\text{Hg}(^{12}\text{C},4n\gamma)$, $E(^{12}\text{C})=80$ MeV with 85.3% enriched ^{198}Hg target cooled at -30° C. The beam was pulsed with 1 ns width separated by 1.5 μs periods. Detectors: two Ge(Li); Measured: $E\gamma$, $I\gamma$, $\gamma(t)$ $I(t,\theta)$, angular distributions, g-factors, $T_{1/2}$; Deduced: μ .

2008An01,2008Kr01: $^{197}\text{Au}(^{14}\text{N},5n\gamma)$, $E(^{14}\text{N})=82$ MeV. Target: 3 mg/cm²; OSIRIS-II gamma-ray spectrometer consisting of 10 HPGe detectors and internal electron conversion spectrometer with two Si(Li) and three PIPS detectors; Deduced: $\alpha(K)\text{exp}$ and $\alpha(L)\text{exp}$.

Others: 1970InZZ, 1969Ru08.

 ^{206}Rn Levels

E(level) [†]	J^π [‡]	$T_{1/2}$	Comments
0.0	0 ⁺		
575.40 10	2 ⁺		
1134.50 15	4 ⁺		
1763.30 18	6 ⁺		
1818.50 18	6 ⁺		
1924.60 20	8 ⁺	19 ns 3	$T_{1/2}$: From 575.4 $\gamma(t)$ in 1981Ho29 and by taking into account the half-life of the 2476-keV isomer. Other: 13.5 ns 10 (1981Ma28) using 575.4 γ -559.1 γ -628.8 $\gamma(t)$. μ : g=0.83 5 (1981Ma28). This value was corrected for diamagnetic and Knight shift by +1.86% 2. $\mu=6.6$ 4. Configuration= $\pi(h_{9/2}^{+2})_{8+}$.
2270.4 3	9 ⁻		
2476.0 3	10 ⁻	65 ns 5	$T_{1/2}$: From 551.2 $\gamma(t)$ in 1981Ho29. The non-observation of a prompt component in the time spectrum suggests that this transition directly depopulates the isomer. Other: 75 ns 10 (1981Ma28) using 575.4 γ -559.1 γ -628.8 $\gamma(t)$. μ : g=1.120 10 (1981Ma28). This value was corrected for diamagnetic and Knight shift by +1.86% 2. $\mu=11.20$ 10. Configuration= $\pi(f_{7/2}^{+1}, i_{13/2}^{+1})_{10-}$.
2534.9 3	10 ⁺		
2585.8 4	11 ⁽⁺⁾		
2834.6 5	12		
3131.8 5	12 ⁺		
3362.4 5	13 ⁽⁺⁾		
3887.9? 6	14		
4130.2 9	15	11 ns 2	$T_{1/2}$: From 768 $\gamma(t)$ in 1981Ho29. Possible configuration= $\pi(h_{9/2}^{+3}, i_{13/2}^{+1})$. The assignment is tentative.

[†] From a least-squares fit to $E\gamma$.

[‡] From 1981Ho29, based on $\gamma(\theta)$ data and the apparent band structures.

(HI,xn γ) 1981Ho29,1981Ma28,2008An01 (continued)

$\gamma(^{206}\text{Rn})$							Comments
E_γ †	$E_i(\text{level})$	J_i^π	E_f	J_f^π	Mult.‡	$I_{(\gamma+ce)}^\#$	
109.8 3	2585.8	11(+)	2476.0	10 ⁻	(E1)	4	Mult.: $A_2=-0.20$ 13 in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$; $A_2=-0.11$ 37 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$. $I_{(\gamma+ce)}$: Other: 13 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$ at 95 MeV (1981Ho29).
161.3 1	1924.60	8 ⁺	1763.30	6 ⁺	E2	120	Mult.: $A_2=0.11$ 7 in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$; $A_2=0.06$ 2 and $A_4=-0.48$ 32 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$. $I_{(\gamma+ce)}$: Other: 90 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$ at 95 MeV (1981Ho29).
205.7 2	2476.0	10 ⁻	2270.4	9 ⁻	M1	32	Mult.: $A_2=-0.06$ 5 and $A_4=0.17$ 8 in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$; $A_2=-0.38$ 11 and $A_4=0.40$ 16 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$; using α deduced from intensity balance in $\gamma\gamma$ coin data. $I_{(\gamma+ce)}$: Other: 31 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$ at 95 MeV (1981Ho29).
242 1	4130.2	15	3887.9?	14			E_γ : γ is shown in the decay scheme, but it is not listed in γ table.
248.9 3	2834.6	12	2585.8	11(+)	D		Mult.: $A_2=-0.26$ 6 in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$; $A_2=-0.52$ 9 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$. $I_{(\gamma+ce)}$: 5 (E1) or 10 (M1) in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$ at 88 MeV (1981Ho29); $I_{(\gamma+ce)}=4$ (E1) or 8 (M1) in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$ at 95 MeV (1981Ho29).
346.0 2	2270.4	9 ⁻	1924.60	8 ⁺	E1	46	Mult.: $\alpha(\text{K})\text{exp}=0.010$ 3 and $\alpha(\text{L})\text{exp}=0.0097$ 23 (2008An01); $A_2=-0.08$ 2 in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$; $A_2=0.01$ 3 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$. $I_{(\gamma+ce)}$: Other: 55 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$ at 95 MeV (1981Ho29).
^x 427.9							
525.5 3	3887.9?	14	3362.4	13(+)	D		Mult.: $A_2=-0.56$ 40 in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$; $A_2=-0.13$ 30 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$. $I_{(\gamma+ce)}$: 4 (E1) or 5 (M1) in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$ at 88 MeV (1981Ho29); $I_{(\gamma+ce)}=6$ (E1) or 7 (M1) in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$ at 95 MeV (1981Ho29).
527.8 2	3362.4	13(+)	2834.6	12	D		Mult.: $A_2=-0.55$ 4 and $A_4=0.64$ 61 in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$; $A_2=-0.50$ 6 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$. $I_{(\gamma+ce)}$: 16 (E1) or 18 (M1) in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$ at 88 MeV (1981Ho29); $I_{(\gamma+ce)}=18$ (E1) or 20 (M1) in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$ at 95 MeV (1981Ho29).
551.2 2	2476.0	10 ⁻	1924.60	8 ⁺	M2	16	Mult.: $\alpha(\text{K})\text{exp}=0.20$ 5 (2008An01); $A_2=0.00$ 3 in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$; $A_2=0.03$ 6 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$. $I_{(\gamma+ce)}$: Other: 30 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$ at 95 MeV (1981Ho29).
559.1 1	1134.50	4 ⁺	575.40	2 ⁺	E2	98	Mult.: $\alpha(\text{K})\text{exp}=0.027$ 7 and $\alpha(\text{L})\text{exp}=0.0076$ 18 (2008An01); $A_2=0.13$ 1 in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$; $A_2=0.16$ 2 and $A_4=0.31$ 22 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$. $I_{(\gamma+ce)}$: Other: 104 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$ at 95 MeV (1981Ho29).
575.4 1	575.40	2 ⁺	0.0	0 ⁺	E2	100	Mult.: $\alpha(\text{K})\text{exp}=0.018$ 4 and $\alpha(\text{L})\text{exp}=0.0068$ 16 (2008An01); $A_2=0.13$ 2 in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$; $A_2=0.15$ 1 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$. $I_{(\gamma+ce)}$: Other: 100 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$ at 95 MeV (1981Ho29).
596.9 3	3131.8	12 ⁺	2534.9	10 ⁺	E2	9	Mult.: $\alpha(\text{K})\text{exp}=0.018$ 5 (2008An01); $A_2=0.04$ 3 and $A_4=0.25$ 6 in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$; $A_2=0.16$ 5 and $A_4=0.16$

Continued on next page (footnotes at end of table)

(HI,xn γ) 1981Ho29,1981Ma28,2008An01 (continued) $\gamma(^{206}\text{Rn})$ (continued)

E_γ [†]	$E_i(\text{level})$	J_i^π	E_f	J_f^π	Mult. [‡]	$I_{(\gamma+ce)}$ [#]	Comments
							7 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$. $I_{(\gamma+ce)}$: Other: 20 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$ at 95 MeV (1981Ho29).
610.3 2	2534.9	10 ⁺	1924.60	8 ⁺	E2	16	Mult.: $A_2=0.34$ 10 in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$; $A_2=0.21$ 10 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$. $I_{(\gamma+ce)}$: 26 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$ reaction at 95 MeV (1981Ho29).
628.8 1	1763.30	6 ⁺	1134.50	4 ⁺	E2	93	Mult.: $\alpha(K)\text{exp}=0.011$ 3 and $\alpha(L)\text{exp}=0.0016$ 4 (2008An01); $A_2=0.11$ 3 in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$; $A_2=0.12$ 1 and $A_4=-0.48$ 16 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$. $I_{(\gamma+ce)}$: Other: 97 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$ at 95 MeV (1981Ho29).
684.0 1	1818.50	6 ⁺	1134.50	4 ⁺	E2		E_γ : From adopted gammas. Mult.: $\alpha(K)\text{exp}=0.017$ 4 (2008An01). Note, that 684 γ is labeled as E1 in Table 1 in 2008An01.
768 1	4130.2	15	3362.4	13 ⁽⁺⁾			E_γ : γ is shown in the decay scheme, but it is not listed in γ table.
776.6 3	3362.4	13 ⁽⁺⁾	2585.8	11 ⁽⁺⁾	E2	13	Mult.: $A_2=0.22$ 3 and $A_4=-0.45$ 46 in $^{197}\text{Au}(^{14}\text{N},5n\gamma)$; $A_2=0.18$ 18 and $A_4=0.29$ 28 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$. $I_{(\gamma+ce)}$: Other: 15 in $^{194}\text{Pt}(^{16}\text{O},4n\gamma)$ at 95 MeV (1981Ho29).

[†] From 1981Ho29. Uncertainties were assigned by the evaluators from authors' general statement that they are 0.1 to 0.3 keV.

[‡] From $\gamma(\theta)$ data and the apparent band structures in 1981Ho29 and $\alpha(K)\text{exp}$ and $\alpha(L)\text{exp}$ in 2008An01.

[#] From $^{197}\text{Au}(^{14}\text{N},5n\gamma)$ reaction at 88 MeV (1981Ho29). The α from 1978Ro21 were used by the authors.

^x γ ray not placed in level scheme.

(HI,xn γ) 1981Ho29,1981Ma28,2008An01

Level Scheme

