

$^9\text{Be}(^{208}\text{Pb},\text{X}\gamma)$ 2011St21

Type	Author	History Citation	Literature Cutoff Date
Full Evaluation	F. G. Kondev	NDS 201,346 (2025)	21-Jan-2025

2011St21: ^{208}Pb beam at 1 GeV/A energy was provided by the SIS-18 accelerator at GSI, Darmstadt. Target consisting of 2.526 g/cm² ^9Be backed by ^{93}Nb foil of thickness=0.223 g/cm². Fragments were identified in flight by the Fragment Separator (FRS) operated in achromatic mode based on time of flight, $B\rho$ and energy loss. They were slowed in Al degraders and stopped in a plastic catcher. The stopper was surrounded by the RISING γ -ray spectrometer. Measured delayed $E\gamma$, $I\gamma$, $\gamma(t)$.

 ^{206}Tl Levels

$E(\text{level})^\dagger$	J^π^\ddagger	$T_{1/2}$	Comments
0.0	0^-	4.202 min 14	$T_{1/2}$: From Adopted Levels.
265.4 5	2^-		
951.2 7	4^-		
1404.1 9	$(5)^+$	71 ns 4	$T_{1/2}$: From sum of 266 $\gamma(t)$, 453 $\gamma(t)$ and 686 $\gamma(t)$ in 2011St21. Experimental isomeric state population ratio=14.2% +21-32.

† From least-squares fit to $E\gamma$.

‡ From Adopted Levels.

 $\gamma(^{206}\text{Tl})$

E_γ^\dagger	I_γ^\dagger	$E_i(\text{level})$	J_i^π	E_f	J_f^π
265.4 5	89 6	265.4	2^-	0.0	0^-
452.9 5	90 6	1404.1	$(5)^+$	951.2	4^-
685.8 5	100 7	951.2	4^-	265.4	2^-

† From 2011St21. $\Delta E_\gamma=0.5$ keV assigned in consultation with Zs. Podolyak (U. Surrey).

$^9\text{Be}(^{208}\text{Pb},\text{X}\gamma)$ 2011St21

Level Scheme

Intensities: Relative I_γ

Legend

- \longrightarrow $I_\gamma < 2\% \times I_\gamma^{\text{max}}$
- \longrightarrow $I_\gamma < 10\% \times I_\gamma^{\text{max}}$
- \longrightarrow $I_\gamma > 10\% \times I_\gamma^{\text{max}}$

