⁹Be(238 U,X γ) **2012Go19**

History

Type Author Citation Literature Cutoff Date
Full Evaluation Shaofei Zhu and E. A. Mccutchan NDS 175, 1 (2021)

1-May-2021

2012Go19: 214 Pb ions were produced in fragmentation reaction with 1 GeV/nucleon 238 U beam on 9 Be target, and were separated and identified by fragment separator (FRS) at GSI. The γ rays were detected by RISING γ -spectrometer of 105 Ge crystals in correlation with the 214 Pb ions. Measured E γ , level half-life by γ (t). Deduced levels, J, π .

²¹⁴Pb Levels

E(level) [†]	$J^{\pi \ddagger}$	$T_{1/2}$	Comments			
0#	0+					
835 [#] 1	(2^{+})					
1179 [#] 2	(4^{+})					
1365 [#] 2	(6^{+})					
1365+x? [#]	(8^{+})	6.2 μ s 3	%IT=100			
			E(level): x<88 keV (2012Go19).			
			$T_{1/2}$: from $\gamma(t)$ decay curve (2012Go19).			

[†] From E γ values, assuming 1 keV uncertainty for each γ ray.

γ (214Pb)


E_{γ}	$E_i(level)$	\mathbf{J}_i^{π}	\mathbf{E}_f	\mathbf{J}_f^{π}	Comments
х	1365+x?	(8+)	1365	(6 ⁺)	E_{γ} : transition to (6 ⁺) level was not seen in γ -ray spectra, x is assumed to be smaller than 88 keV based on the observation that the intensity of K_{α} x rays was only compatible with that from the internal conversion of the 6 ⁺ to 4 ⁺ transition (2012Go19).
186	1365	(6^+)	1179	(4^{+})	
344	1179	(4^{+})	835	(2^{+})	
835	835	(2^{+})	0	0_{+}	

[‡] The transitions in the yrast cascade are attributed to E2 transitions with the assumption that the isomer is predominantly a fully aligned pair of neutrons with a $(g_{9/2})^2$ configuration (2012Go19).

[#] Seq.(A): Yrast cascade.

9 Be(238 U,Χ γ) 2012Go19

Level Scheme



⁹Be(238 U,Χ γ) **2012Go19**

Seq.(A): Yrast cascade



 $^{214}_{\ 82}\mathrm{Pb}_{132}$