

Adopted Levels

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	Balraj Singh and Michael Birch		ENSDF	15-May-2011

$Q(\beta^-)=5.9\times 10^3$  syst;  $S(n)=3.9\times 10^3$  syst;  $S(p)=7.2\times 10^3$  syst;  $Q(\alpha)=3.5\times 10^3$  syst 2012Wa38

Note: Current evaluation has used the following Q record \$ 5420 syst 4070 syst 7410 syst 3330 syst 2011AuZZ.

Estimated uncertainties in 2011AuZZ: 500 for  $Q(\beta^-)$ , 640 for  $S(n)$  590 for  $S(p)$  and  $Q(\alpha)$ .

$S(2n)=9690$  510,  $S(2p)=16870$  640 (syst,2011AuZZ).

2010AI24:  $^{226}\text{At}$  nuclide identified in  $^9\text{Be}(^{238}\text{U},\text{X})$  reaction with a beam energy of 1 GeV/nucleon produced by the SIS synchrotron at GSI facility. Target=2500 mg/cm<sup>2</sup>. The fragment residues were analyzed with the high resolving power magnetic spectrometer Fragment separator (FRS). The identification of nuclei was made on the basis of magnetic rigidity, velocity, time-of-flight, energy loss and atomic number of the fragments using two plastic scintillators and two multisampling ionization chambers. The FRS magnet was tuned to center on  $^{210}\text{Au}$ ,  $^{216}\text{Pb}$ ,  $^{219}\text{Pb}$ ,  $^{227}\text{At}$  and  $^{229}\text{At}$  nuclei along the central trajectory of FRS.

Unambiguous identification of nuclides required the separation of different charge states of the nuclei passing through the FRS. At 1 GeV/nucleon incident energy of  $^{238}\text{U}$ , fraction of fully stripped  $^{226}\text{Po}$  nuclei was about 89%. Through the measurement of difference in magnetic rigidity in the two sections of the FRS and the difference in energy loss in the two ionization chambers, the charge state of the transmitted nuclei was determined, especially, that of the singly charged (hydrogen-like) nuclei which preserved their charge in the current experimental setup. Measured production cross sections with 10% statistical and 20% systematic uncertainties.

Criterion established in 2010AI24 for acceptance of identification of a new nuclide: 1. number of events should be compatible with the corresponding mass and atomic number located in the expected range of positions at both image planes of the FRS spectrometer; 2. number of events should be compatible with >95% probability that at least one of the counts does not correspond to a charge-state contaminant. Comparisons of measured  $\sigma$  with model predictions using the computer codes COFRA and EPAX.

 $^{226}\text{At}$  Levels

E(level)	$T_{1/2}$	Comments
0	>300 ns	<p><math>{}^{\%}\beta^-=?</math></p> <p>Production <math>\sigma=45.4</math> nb (from e-mail reply of Oct 29, 2010 from H. Alvarez-Pol, which also stated that further analysis was in progress).</p> <p>From A/Z plot (figure 1 in 2010AI24), a large number (certainly more than few hundreds) of events are assigned to <math>^{226}\text{At}</math>.</p> <p>E(level): the observed fragments are assumed to be in the ground state of <math>^{226}\text{At}</math> nuclei.</p> <p>The <math>\beta^-</math> decay is the only decay mode expected.</p> <p><math>T_{1/2}</math>: lower limit from time-of-flight as given in 2006Ca30 for a similar setup. Actual half-life is expected to be much larger as suggested by the calculated value of 4.2 s for <math>\beta</math> decay and <math>&gt;10^{20}</math> s for <math>\alpha</math> decay (1997Mo25), and systematic value of 20 s for <math>\beta</math> decay (2011AuZY).</p> <p><math>J^\pi</math>: <math>1/2^+</math> for proton and <math>5/2^-</math> neutron configuration predicted in 1997Mo25 calculations.</p> <p>Production cross section measured in 2010AI24, values are given in figure 2, plot of <math>\sigma</math> versus mass number for At isotopes. Statistical uncertainty=10%, systematic uncertainty=20%.</p>