

Be($^{32}\text{Mg}, \text{X}\gamma$) 2015Pe09

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	M. S. Basunia, A. Chakraborty		NDS 197,1 (2024)	31-May-2024

Adapted/Edited the XUNDL dataset compiled by J. Chen (NSCL, MSU), July 17, 2015.

One-proton and one-neutron knockout reaction.

2015Pe09: ^{32}Mg secondary beam, $E=87$ MeV/nucleon, was produced by fragmentation of ^{48}Ca primary beam, $E=140$ MeV/nucleon, at NSCL, bombarding a ^9Be target, thickness 888 mg/cm². The secondary target was also ^9Be , thickness 376 mg/cm². Fragments were separated by the A1900 separator. γ rays were detected with the SeGA array comprised of sixteen 32-fold segmented HPGe detectors. Measured $E\gamma$, $I\gamma$, $\gamma\gamma$ -coin, ^{30}Na - γ -coin. Deduced levels, J, π , band structures, configurations.

 ^{30}Na Levels

$E(\text{level})^\dagger$	$J^\pi^\ddagger$	$T_{1/2}$	Comments
0 [#]	(2 ⁺)		
151.0 [@] 10	(1 ⁺)	≈ 347 ps	$T_{1/2}$: from $\tau \approx 500$ ps, estimated in 2015Pe09 from low-energy tail of 151 γ in the γ spectra.
338.0 ^{&} 15			
424.0 [#] 20	(3 ⁺)		
510.0 ^{&} 25			
516.0 [@] 23	(2 ⁺)		
758 ^{&} 4			
925 [#] 3	(4 ⁺)		
926 ^a 3	(1 ⁺)		
1032 [@] 3			
1263 [@] 4			
1527 ^{&} 5			

[†] From a least-squares fit to γ -ray energies.

[‡] Tentative assignments in 2015Pe09 based on band structure.

[#] Band(A): $K^\pi=(2^+)$, 2p2h, g.s. band.

[@] Band(B): $K^\pi=(1^+)$, 2p2h band.

[&] Band(C): 1p1h/3p3h, $\pi=-$ band.

^a Band(D): $K^\pi=(1^+)$, 0p0h band.

 $\gamma(^{30}\text{Na})$

$E_\gamma^\dagger$	$E_i(\text{level})$	J_i^π	E_f	J_f^π	Comments
151 1	151.0	(1 ⁺)	0	(2 ⁺)	E_γ : listed in 2015Pe09 from 2007Tr08. The measured value is 148 keV in 2015Pe09. The E_γ shift is attributed to the longer mean lifetime ($\tau \approx 500$ ps) of this state by the authors of 2015Pe09.
172 [‡] 2	510.0		338.0		
187 [‡] 2	338.0		151.0	(1 ⁺)	
248 [‡] 2	758		510.0		
338 [‡] 2	338.0		0	(2 ⁺)	
365 2	516.0	(2 ⁺)	151.0	(1 ⁺)	
410 2	926	(1 ⁺)	516.0	(2 ⁺)	
424 2	424.0	(3 ⁺)	0	(2 ⁺)	
501 2	925	(4 ⁺)	424.0	(3 ⁺)	
516 ^{‡#} 2	1032		516.0	(2 ⁺)	

Continued on next page (footnotes at end of table)

Be(³²Mg,Xγ) 2015Pe09 (continued)

γ(³⁰Na) (continued)

<u>E_γ[†]</u>	<u>E_i(level)</u>	<u>J_i^π</u>	<u>E_f</u>	<u>J_f^π</u>
747 [‡] # 3	1263		516.0	(2 ⁺)
769 [‡] 3	1527		758	

[†] From 2015Pe09.

[‡] New transition (2015Pe09).


The γ was identified in singles spectrum, not expected to be present in γγ-coin spectrum. The placement in the level scheme is firm.

Be($^{32}\text{Mg}, X\gamma$) 2015Pe09


Legend

Level Scheme

● Coincidence



Be($^{32}\text{Mg},\text{X}\gamma$) 2015Pe09



$^{30}_{11}\text{Na}_{19}$