

$^{40}\text{Ca}(\mu^-, \nu\alpha\gamma)$ **2006Me08**

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	Jun Chen, John Cameron and Balraj Singh		NDS 112,2715 (2011)	20-Oct-2011

2006Me08: The μ^- beam obtained from decay of π^- beam at 90 MeV/c from M9B beamline at TRIUMF. Measured $I\gamma$, $\gamma\gamma$, γ -p using two HPGe detectors.

Muonic Lyman series for natural Calcium

μ x ray	Energy	Intensity in percent
2p-1s	783.659 25	83.8 10
3p-1s	940.63 10	6.2 2
4p-1s	995.48 10	2.0 1
5p-1s	1020.81 10	2.0 1
6p-1s	1034.62 10	1.8 1
7p-1s	1042.71 20	1.4 1
(8- ∞)p-1s	1046-1063	2.8 4

Muonic Balmer series for natural Calcium

μ x ray	Energy	Intensity in percent
3d-2p	157.35 13	64.5 9
4d-2p	212.03 10	8.85 20
5d-2p	237.31 10	4.34 20
6d-2p	251.06 10	3.29 20
7d-2p	259.45 10	1.37 20
(8- ∞)d-2p	261-277	1.4 3

 ^{35}Cl Levels

E(level) [†]	J [†]
0	3/2 ⁺
1219.3	1/2 ⁺
1763.0	5/2 ⁺
2645.7	7/2 ⁺
2693.8	3/2 ⁺

[†] From Adopted Levels.

 $\gamma(^{35}\text{Cl})$

E $_{\gamma}$ [†]	Percent γ -ray yield	E $_i$ (level)	J $^{\pi}_i$	E $_f$	J $^{\pi}_f$
1219.3	<0.3	1219.3	1/2 ⁺	0	3/2 ⁺
1763.1	0.5 3	1763.0	5/2 ⁺	0	3/2 ⁺
2645.7	<0.2	2645.7	7/2 ⁺	0	3/2 ⁺
2693.6	<0.2	2693.8	3/2 ⁺	0	3/2 ⁺

[†] Round off values from Adopted Gammas.

$^{40}\text{Ca}(\mu^-, \nu\alpha n\gamma)$ **2006Me08**Level SchemeIntensities: Percent γ -ray yield per muon capture

Legend

- $I_\gamma < 2\% \times I_\gamma^{max}$
- $I_\gamma < 10\% \times I_\gamma^{max}$
- $I_\gamma > 10\% \times I_\gamma^{max}$

