

$^9\text{Be}(^{44}\text{S},\text{X}\gamma)$ 2009Ri11

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	Balraj Singh and Jun Chen [#]		NDS 126, 1 (2015)	31-Mar-2015

2009Ri11: E=92 MeV/nucleon ^{44}S beam was produced by fragmentation of a 140 MeV/nucleon ^{48}Ca on a ^9Be fragmentation target and incident on a target of 376 mg/cm² thick ^9Be . Fragments (84% ^{44}S , 14% ^{45}Cl) were separated by the A1900 separator and identified by the time-of-flight and energy loss in the S800 ionization chamber; γ -rays were detected by the Segmented Germanium Array (SeGA). Measured E_γ , I_γ , $\gamma\gamma$ -coin. Deduced levels, J, π , branching ratios and rotational band. Comparisons with shell-model calculations.

This dataset shares the γ -energies with the dataset of $^9\text{Be}(^{45}\text{Cl},\text{X}\gamma)$.

 ^{43}S Levels

E(level) [†]	J π [‡]
0 [#]	3/2 ⁻
970 [#] 5	(5/2 ⁻ , 7/2 ⁻)
1153 [#] 5	(5/2 ⁻ , 7/2 ⁻)
2616 9	(7/2 ⁻)

[†] From least-squares fit to E_γ data.

[‡] From comparisons with shell-model calculations.

[#] Band(A): Ground state rotational band.

 $\gamma(^{43}\text{S})$

E_γ	I_γ	$E_i(\text{level})$	J π_i	E_f	J π_f
183 1	53 3	1153	(5/2 ⁻ , 7/2 ⁻)	970	(5/2 ⁻ , 7/2 ⁻)
^x 231 1	6 1				
^x 459 3	7 2				
^x 621 4	31 3				
^x 719 4	21 3				
^x 770 5	12 3				
^x 849 5	24 3				
971 6	56 4	970	(5/2 ⁻ , 7/2 ⁻)	0	3/2 ⁻
1154 7	100	1153	(5/2 ⁻ , 7/2 ⁻)	0	3/2 ⁻
^x 1203 7	21 3				
1468 9	5 3	2616	(7/2 ⁻)	1153	(5/2 ⁻ , 7/2 ⁻)
^x 1529 9	8 3				
^x 1855 11	5 3				
2600 16	98 7	2616	(7/2 ⁻)	0	3/2 ⁻

^x γ ray not placed in level scheme.


$^9\text{Be}(^{44}\text{S},\text{X}\gamma)$ 2009Ri11

Level Scheme

Intensities: Relative I_γ


Legend

- \longrightarrow $I_\gamma < 2\% \times I_\gamma^{max}$
- \longrightarrow $I_\gamma < 10\% \times I_\gamma^{max}$
- \longrightarrow $I_\gamma > 10\% \times I_\gamma^{max}$



$^9\text{Be}(^{44}\text{S},\text{X}\gamma)$ 2009Ri11

Band(A): Ground state rotational band



$^{43}_{16}\text{S}_{27}$