

Adopted Levels, Gammas

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	Jun Chen and Balraj Singh		NDS 157, 1 (2019)	15-Apr-2019

$Q(\beta^-)=4958$ 15; $S(n)=6360.8$ 16; $S(p)=17266.7$ 18; $Q(\alpha)=-12241.2$ 19 [2017Wa10](#)

$S(2n)=11507.2$ 16, $S(2p)=31890$ 310 ([2017Wa10](#)).

Mass measurement: [2012La05](#) (TOF-ICR resonance frequency ratios using TITAN Penning trap spectrometer at TRIUMF-ISAC facility).

Theory references: consult the NSR database (www.nndc.bnl.gov/nsr/) for 125 primary references for structure calculations.

[Additional information 1](#).

 ^{50}Ca LevelsCross Reference (XREF) Flags

A	^{50}K β^- decay (472 ms)	F	$^{48}\text{Ca}(\alpha,2p)$
B	^{51}K β^-n decay (365 ms)	G	$^{48}\text{Ca}(^{238}\text{U},X\gamma)$
C	^{52}K β^-2n decay (110 ms)	H	$^{208}\text{Pb}(^{48}\text{Ca},X\gamma)$
D	$^1\text{H}(^{50}\text{Ca},p'\gamma)$	I	$^{238}\text{U}(^{48}\text{Ca},X\gamma)$
E	$^{48}\text{Ca}(t,p),(pol\ t,p)$		

E(level) [†]	J^π [‡]	$T_{1/2}$ [§]	XREF	Comments
0.0	0 ⁺	13.45 s 5	ABCDEFGH	$\% \beta^- = 100$ $T_{1/2}$: measured by 2017Ga25 from fit to the decay curves of 1519- and 1591-keV γ transitions, ^{50}Ca beam produced in Ta(p,X),E=500 MeV at TRIUMF, and counted using GRIFFIN array of Ge detectors. Others: 13.9 s 6 (1970Wa29 , from decay curve for 257 γ); 14 s 3 (1968Ch11 , from decay curve for all γ rays); 9 s 2 (1964Sh14 , decay curves for 72 γ and 258 γ). Nuclear rms charge radius: $\langle r^2 \rangle^{1/2} = 3.517$ fm 7 (2013An02 , evaluation). Measured $\delta r^2(^{40}\text{Ca}, ^{50}\text{Ca}) = 0.291$ fm ² 3(stat) 12(syst) (2016Ga34 , using COLLAPS at ISOLDE-CERN; see also 2017Ne04 review article on measurements at this facility). Previous measurement: 0.276 fm ² 34 (1992Ve02 , online collinear laser spectroscopy). Measured isotope shift $\delta\nu(^{40}\text{Ca}, ^{50}\text{Ca}) = 1969.2$ MHz 9(stat) 47(syst) (2016Ga34 , using COLLAPS at ISOLDE-CERN). Previous measurement: 1951 MHz 9(stat) 20(syst) (1992Ve02 , online collinear laser spectroscopy). Measurement of isotope shift and rms radii: 1992Ve02 , 2017Ne04 .
1026.72 10	2 ⁺	66.5 ps 21	ABCDEFGH	J^π : E2 1026.7 γ to 0 ⁺ ; L(t,p)=2 from 0 ⁺ . $T_{1/2}$: recoil-distance Doppler-shift method (2009Va06) in $^{208}\text{Pb}(^{48}\text{Ca},X\gamma)$. Other: 68.6 ps 55 from DSAM in ($^{50}\text{Ca},p'\gamma$).
3002.1 5	(2 ⁺)	<0.69 ps	AB DEFGH	J^π : L(t,p)=(2). L(t,p)=(4) and L($\alpha,2p$)=(4) are also proposed but in the latter case L=2 does not seem ruled out in figure 32 of 1990Fi07 .
3531.7 4	(1,2 ⁺)		AB E	XREF: E(3519). J^π : 3531.7 γ to 0 ⁺ . Note that $J^\pi=0^+$ is suggested in 1968Br01 , 1967G108 and 1966Ve06 in theoretical analyses of (t,p) results for a 3519 level observed by 1967Bj06 . It is possible two separate levels are populated near this energy.
3997.22 21	(3 ⁻)	<0.69 ps	DEFGH	J^π : L(t,p)=(3). Inconsistent with L($\alpha,2p$)=4, but L=3 comparison of $\sigma(\theta)$ data was not shown in figure 32 of 1990Fi07 .
4035.7 4	(1,2 ⁺)		AB D	J^π : 4035.6 γ to 0 ⁺ .
4475.8 5	(0 ⁺)		A E	J^π : L(t,p)=(0).
4515.04 14	(4 ⁺)	<1.04 ps	DE GHI	J^π : strong population in $^{238}\text{U}(^{48}\text{Ca},X\gamma)$ suggests yrast 4 ⁺ level. L(t,p)=(3) for a 4517 15 group is inconsistent.
4830.6 3	(4)	<0.69 ps	E GHI	J^π : L(t,p)=(4); (4 ⁻) proposed in $^{238}\text{U}(^{48}\text{Ca},X\gamma)$ from γ to (3 ⁻).
4870 5	(2 ⁺)		G	J^π : 4870 γ to 0 ⁺ . J=1 less likely to be populated in high-spin reaction.
4886.3 5	(1 ⁻)		A E	J^π : L(t,p)=(1); 4886.0 γ to 0 ⁺ .

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{50}Ca Levels (continued)

E(level) [†]	J^π [‡]	$T_{1/2}$ ^a	XREF	Comments
4.97×10 ³ 5	(4 ⁺ &5 ⁻) [#]		F	$J^\pi, E(\text{level}): L(\alpha, 2p)=4+5$ for a possible doublet.
5043 15	(1 ⁻)		E	$J^\pi: L(t,p)=(1)$.
5084.56 25	(4 ⁻) [@]		I	
5109.88 20	(5 ⁻) [@]	<0.69 ps	DE GHI	
5147.34 17	(5 ⁺) [@]		I	
5168 20			E	
5281 20			E	
5362 20			E	
5434 20			E	
5516.92 20	(5 ⁻) [@]		E I	$J^\pi: L(t,p)=(4)$, but data were insufficient to get a reliable L value.
5576 20			E	$J^\pi: L(t,p)=(4)$, but data were insufficient to get a reliable L value.
6519 8			A	%n≈100 Additional information 2.
6869.27 25	(7 ⁻) [@]		I	
7039 36			A	%n≈100 Additional information 3.
7269 46			A	%n≈100 Additional information 4.
7309 51			A	%n≈100 Additional information 5.
7619 66			A	%n≈100 Additional information 6.
7999 87			A	%n≈100 Additional information 7.
8249 97			A	%n≈100 Additional information 8.
8.38×10 ³ 5	(7 ⁻) [#]		F	
8.81×10 ³ 12	(0 ⁻ , 1 ⁻) ^{&}		A	%n≈100 Additional information 9.
8.98×10 ³ 5	(7 ⁻) [#]		F	
9239 46	(0 ⁻ , 1 ⁻) ^{&}		A	%n≈100 Additional information 10.
9779 72	(0 ⁻ , 1 ⁻) ^{&}		A	%n≈100 Additional information 11.
9.80×10 ³ 5	(6 ⁺) [#]		F	$J^\pi: 8^+$ is not completely ruled out.
10.33×10 ³ 5	(8 ⁺) [#]		F	$J^\pi: 6^+$ is not completely ruled out.
10430 36	(0 ⁻ , 1 ⁻) ^{&}		A	%n≈100 Additional information 12.
1.055×10 ⁴ 11	(0 ⁻ , 1 ⁻) ^{&}		A	%n≈100 Additional information 13.
11059 36	(0 ⁻ , 1 ⁻) ^{&}		A	%n≈100 Additional information 14.
11479 52	(0 ⁻ , 1 ⁻) ^{&}		A	%n≈100 Additional information 15.

[†] For levels populated in γ -ray studies, values are from least-squares to γ -ray energies. For levels populated in particle-reaction studies, averages are taken when possible.

[‡] From DWBA analysis of $\sigma(\theta)$ in (t,p), except as noted. See (t,p) for additional tentative J^π assignments. For L(t,p) and L(α ,2p) transfer reactions, target $J^\pi=0^+$. Most L(t,p) are considered by the evaluators as tentative values due either to disagreements with

Adopted Levels, Gammas (continued) ^{50}Ca Levels (continued)

other reactions or to weak populations of levels.

From CCBA and DWBA analyses in $(\alpha,2p)$.

@ Tentative assignments in $^{238}\text{U}(^{48}\text{Ca},X\gamma)$ based on γ -decay pattern and possible model predictions. No supporting data are available for transition multipolarity assignments.

& From allowed β transition ($\log ft=4.1$ to 4.9) from $0^{(-)}$ parent state.

^a From RDDS method in $^{208}\text{Pb}(^{48}\text{Ca},X\gamma)$, unless otherwise stated.

$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult.#	$\gamma(^{50}\text{Ca})$ Comments
1026.72	2^+	1026.7 1	100	0.0	0^+	E2	B(E2)(W.u.)=0.68 2 E $_\gamma$: weighted average of 1027.0 5 from $^{50}\text{K} \beta^-$ decay (472 ms), 1026.2 3 from $(^{238}\text{U},X\gamma)$, and 1026.7 1 from $(^{48}\text{Ca},X\gamma)$. Others: 1027 1 from $^{51}\text{K} \beta^-n$ decay (365 ms), 1027 1 from $^{52}\text{K} \beta^-2n$ decay (110 ms), and 1028 2 from $(^{50}\text{Ca},p'\gamma)$. Mult.: $\gamma(\theta)$ and linear polarization in $^{208}\text{Pb}(^{48}\text{Ca},X\gamma)$.
3002.1	(2^+)	1975.3 5	100	1026.72	2^+	(D)	E $_\gamma$: from β^- decay. Other: 1976 1 from β^-n decay; 1978.2 6 in $^{48}\text{Ca}(^{238}\text{U},X\gamma)$ is discrepant, possibly due to Doppler-shift effects. It is possible that 1978,2 γ defined a level separate from 3002, (2^+) , but there is no strong evidence for its existence.
3531.7	$(1,2^+)$	2504.5 8	100 13	1026.72	2^+		E $_\gamma$: weighted average of 2504.9 5 from $^{50}\text{K} \beta^-$ decay (472 ms) and 2503 1 from $^{51}\text{K} \beta^-n$ decay (365 ms). I $_\gamma$: from $^{50}\text{K} \beta^-$ decay . Other: 100 17 from β^-n decay.
		3531.7 4	92 13	0.0	0^+		E $_\gamma$: weighted average of 3531.8 5 from $^{50}\text{K} \beta^-$ decay (472 ms) and 3530 2 from $^{51}\text{K} \beta^-n$ decay (365 ms). I $_\gamma$: from $^{50}\text{K} \beta^-$ decay . Other: 133 17 from β^-n decay.
3997.22	(3^-)	2970.3 3	100	1026.72	2^+	D	E $_\gamma$: weighted average of 2964 8 from $(^{50}\text{Ca},p'\gamma)$, 2971.4 6 from $(^{238}\text{U},X\gamma)$, and 2970.2 2 from $(^{48}\text{Ca},X\gamma)$.
4035.7	$(1,2^+)$	3008.8 5	60 4	1026.72	2^+		E $_\gamma$: weighted average of 3008.9 5 from $^{50}\text{K} \beta^-$ decay and 3008 2 from $^{51}\text{K} \beta^-n$ decay. I $_\gamma$: from $^{50}\text{K} \beta^-$ decay. Others: 60 20 from $^{51}\text{K} \beta^-n$ decay and 75 25 from $(^{50}\text{Ca},p'\gamma)$.
		4035.6 [‡] 5	100 6	0.0	0^+		E $_\gamma$: weighted average of 4035.6 5 from $^{50}\text{K} \beta^-$ decay and 4035 2 from $^{51}\text{K} \beta^-n$ decay. Other: 4030 18 from $(^{50}\text{Ca},p'\gamma)$. I $_\gamma$: from $^{50}\text{K} \beta^-$ decay. Others: 100 20 from $^{51}\text{K} \beta^-n$ decay, and 100 25 from $(^{50}\text{Ca},p'\gamma)$.
4475.8	(0^+)	3449.0 [‡] 5	100	1026.72	2^+		
4515.04	(4^+)	518.4 7	2 1	3997.22	(3^-)		
		3488.2 1	100	1026.72	2^+	(E2)	E $_\gamma$: others: 3482 14 from $(^{50}\text{Ca},p'\gamma)$ and 3488.4 8 from $(^{238}\text{U},X\gamma)$.
4830.6	(4)	833.4 2	100	3997.22	(3^-)	(D)	E $_\gamma$: weighted average of 833.9 5 from $(^{238}\text{U},X\gamma)$ and 833.3 2 from $(^{48}\text{Ca},X\gamma)$.
4870	(2^+)	4870 5		0.0	0^+		E $_\gamma$: from $(^{238}\text{U},X\gamma)$ only.
4886.3	(1^-)	4886.0 [‡] 5	100	0.0	0^+		
5084.56	(4^-)	1087.2 3	100	3997.22	(3^-)		
5109.88	(5^-)	594.9 2	100 6	4515.04	(4^+)	D	E $_\gamma$: weighted average of 603 11 from $(^{50}\text{Ca},p'\gamma)$, 595.5 3 from $(^{238}\text{U},X\gamma)$, and 594.8 1 from $(^{48}\text{Ca},X\gamma)$.
		1112.6 2	6.8 14	3997.22	(3^-)		
5147.34	(5^+)	632.3 1	100	4515.04	(4^+)		

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) $\gamma(^{50}\text{Ca})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π
5516.92	(5 ⁻)	407.3 2	75 13	5109.88	(5 ⁻)
		432.3 2	75 13	5084.56	(4 ⁻)
		1001.9 2	100 13	4515.04	(4 ⁺)
		1519.7 5	50 13	3997.22	(3 ⁻)
6869.27	(7 ⁻)	1352.9 3	91 18	5516.92	(5 ⁻)
		1759.1 2	100 9	5109.88	(5 ⁻)

† From $^{238}\text{U}(^{48}\text{Ca}, X\gamma)$, unless stated otherwise.

‡ From β^- decay.

From $\gamma(\theta)$ in $^{208}\text{Pb}(^{48}\text{Ca}, X\gamma)$, unless otherwise stated.

Adopted Levels, Gammas**Level Scheme**

Intensities: Relative photon branching from each level

