${}^{1}\text{H}({}^{59}\text{Sc,2p}\gamma)$ **2023Ch26**

Type Author Citation Literature Cutoff Date
Full Evaluation Balraj Singh ENSDF 08-Sept-2023

Quasi-free one-proton knockout reaction.

2023Ch26: ¹H(⁵⁹Sc,2p),E⁵⁹Sc=199 MeV/nucleon, secondary ⁵⁹Sc beam from ⁹Be(⁷⁰Zn,X),E(⁷⁰Zn)=345 MeV/nucleon, followed by separation and identification of ions of interest using the BigRIPS separator at RIBF-RIKEN facility. Measured reaction residues of ⁵⁸Ca through identification by the SAMURAI spectrometer, Doppler-corrected Eγ, Iγ, (particle)γ-coin spectra restricted to γ multiplicity of ≤5, using the DALI2⁺ array of NaI(Tl) detectors, and the MINOS liquid hydrogen (LH₂) target. Deduced energy of the first 2⁺ level, and production cross sections. Comparison with shell-model calculations with the GXPF1B and A3DA-t Hamiltonians in full *pf* model space, and the state-of-the-art ab initio approaches: valence-space in-medium similarity renormalization group (VS-IMSRG) method, and coupled-cluster (C-C) calculations.

⁵⁸Ca Levels

E(level)	J^{π}	Comments
0	0+	Measured partial cross section for the g.s.=0.66 mb 24.
1115 34	(2 ⁺)	Measured inclusive cross section for 58 Ca=1.14 mb <i>15</i> . J^{π} : from 2023Ch26, based on shell-model predictions. Measured partial cross section for the 1115,(2 ⁺) level=0.47 mb <i>19</i> .

 γ (58Ca)

$$\frac{E_{\gamma}}{1115 \ 34} \quad \frac{E_{i}(\text{level})}{1115} \quad \frac{J_{i}^{\pi}}{(2^{+})} \quad \frac{E_{f}}{0} \quad \frac{J_{f}^{\pi}}{0^{+}}$$

${}^{1}\text{H}({}^{59}\text{Sc,2p}\gamma)$ 2023Ch26

Level Scheme

