

**Adopted Levels, Gammas**

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	Caroline D. Nesaraja, Scott D. Geraedts and Balraj Singh		NDS 111,897 (2010)	12-Jan-2010

Q( $\beta^-$ )=-2307.9 12; S(n)=10044.60 18; S(p)=11957.3 16; Q( $\alpha$ )=-7645.7 5 2012Wa38  
 Note: Current evaluation has used the following Q record -2307.6 1210044.601811955.5 19-7645.8 4 2009AuZZ,2003Au03.  
 S(2n)=17690.69 19, S(2p)=21450.1 21 (2009AuZZ).  
 Structure calculations (levels, transition probabilities, etc.): 2009Su20, 2004Ho08, 2002Ca48, 1997Na04, 1990Ha16, 1979Mc03,  
 1978Jo01, 1976La06, 1974Pa13, 1973Ba12.  
 Additional information 1.

<sup>58</sup>Fe Levels

Cross Reference (XREF) Flags

<b>A</b>	<sup>58</sup> Mn $\beta^-$ decay (3.0 s)	<b>J</b>	<sup>57</sup> Fe(n, $\gamma$ ),(n,n):resonances	<b>S</b>	<sup>59</sup> Co( $\gamma$ ,p)
<b>B</b>	<sup>58</sup> Mn $\beta^-$ decay (65.4 s)	<b>K</b>	<sup>57</sup> Fe(d,p),(pol d,p)	<b>T</b>	<sup>59</sup> Co( $\mu^-$ ,n $\gamma$ )
<b>C</b>	<sup>58</sup> Co $\epsilon$ decay (70.86 d)	<b>L</b>	<sup>58</sup> Fe(e,e')	<b>U</b>	<sup>59</sup> Co(n,d)
<b>D</b>	<sup>48</sup> Ca( <sup>13</sup> C,3n $\gamma$ )	<b>M</b>	<sup>58</sup> Fe(n,n' $\gamma$ )	<b>V</b>	<sup>59</sup> Co(p,2p)
<b>E</b>	<sup>54</sup> Cr( <sup>6</sup> Li,d)	<b>N</b>	<sup>58</sup> Fe(p,p')	<b>W</b>	<sup>59</sup> Co(d, <sup>3</sup> He)
<b>F</b>	<sup>55</sup> Mn( $\alpha$ ,p $\gamma$ )	<b>O</b>	<sup>58</sup> Fe(d,d'),(pol d,d')	<b>X</b>	<sup>62</sup> Ni( <sup>3</sup> He, <sup>7</sup> Be)
<b>G</b>	<sup>56</sup> Fe(t,p),(pol t,p)	<b>P</b>	<sup>58</sup> Fe( <sup>3</sup> He, <sup>3</sup> He')	<b>Y</b>	<sup>60</sup> Ni( $\mu^-$ , $\nu$ p $\nu$ $\gamma$ )
<b>H</b>	<sup>56</sup> Fe( $\alpha$ , <sup>2</sup> He)	<b>Q</b>	<sup>58</sup> Fe( $\alpha$ , $\alpha'$ )	<b>Z</b>	Cu(K $^-$ , $\gamma$ )
<b>I</b>	<sup>57</sup> Fe(n, $\gamma$ ) E=th	<b>R</b>	Coulomb excitation		

E(level) <sup>†</sup>	J $^\pi$ <sup>‡</sup>	T <sub>1/2</sub> <sup>#</sup>	XREF	Comments
0.0 <sup>a</sup>	0 <sup>+</sup>	stable	ABCDEFGHIJKLMN OPQRSTUVWXYZ	<r <sup>2</sup> > <sup>1/2</sup> =3.7748 fm 14 (2004An14,evaluation).
810.7662 <sup>a</sup> 20	2 <sup>+</sup>	6.54 ps 19	ABCDEFGHI I KLMNOPQR TUVWXYZ	$\mu$ =+0.95 11 (2009Ea02) Q=-0.27 5 (1981Le02,1989Ra17) $\mu$ : g factor=+0.468 56 from measured g(811,2 <sup>+</sup> , <sup>58</sup> Ni)/g(847,2 <sup>+</sup> , <sup>56</sup> Fe)= 0.920 55 (2009Ea02) and measured g factor=+0.509 53 (2009Ea01) for the 847, 2 <sup>+</sup> state in <sup>56</sup> Fe. Using earlier measured ratio of 0.75 24 (1977Br23), 2009Ea02 recommend averaged ratio of 0.912 54 and g factor of +0.464 56. Further, 2009Ea02 recommend averaged g factor=+0.473 51 by considering earlier measured (1969Si13, IPAC method) g factor=+0.514 118. 1989Ra17 give +0.92 26 from 1977Br23 (transient- field integral PAC). See also 2005St24 compilation with quoted values from 1977Br23 and 1969Si13. Q: reorientation in Coulomb excitation (1981Le02). See also 2005St24 compilation. J $^\pi$ : E2 $\gamma$ to 0 <sup>+</sup> . T <sub>1/2</sub> : from B(E2)=0.1234 36 (1981Le02, Coul. ex.). 2001Ra27 evaluation gives 6.73 ps 22 based on adopted B(E2)=0.120 4 from Coulomb excitation and DSA methods. Values of 2.4 ps 7 from DSAM in ( $\alpha$ ,p $\gamma$ ) and 8.6 ps 7 from B(E2) in (e,e') are discrepant.
1674.731 <sup>b</sup> 6	2 <sup>+</sup>	1.6 ps 4	BCDEFG I K MNO Q TUVWX	J $^\pi$ : L(t,p)=2. Also M1+E2 $\gamma$ to 2 <sup>+</sup> . T <sub>1/2</sub> : from ( $\alpha$ ,p $\gamma$ ).
2076.52 <sup>a</sup> 3	4 <sup>+</sup>	0.28 ps 4	B D F MN Q T WX	J $^\pi$ : $\Delta$ J=2, E2 $\gamma$ to 2 <sup>+</sup> . T <sub>1/2</sub> : weighted average of 0.24 ps 4 in (n,n' $\gamma$ ), 0.24 ps 7 and 0.37 ps +6-5 in ( $\alpha$ ,p $\gamma$ ).

Continued on next page (footnotes at end of table)

**Adopted Levels, Gammas (continued)** $^{58}\text{Fe}$  Levels (continued)

E(level) <sup>†</sup>	J <sup>π</sup> <sup>‡</sup>	T <sub>1/2</sub> <sup>#</sup>	XREF	Comments
2133.895 <sup>b</sup> 21	3 <sup>+</sup>	2.2 ps 7	B DEFG I K MN T	XREF: N(2123). T <sub>1/2</sub> : from (α,pγ). J <sup>π</sup> : ΔJ=1, M1+E2 γ to 2 <sup>+</sup> , ΔJ=1, dipole γ from 4 <sup>+</sup> . Significant excitation in (p,p'), ( <sup>6</sup> Li,d), and (t,p) suggests important role of indirect two-step processes, L=4 in ( <sup>6</sup> Li,d) from 1977Fu03 contradicts J=3 <sup>+</sup> .
2257.95 21	0 <sup>+</sup> @	>2.5 ps	EFG I K MN U	T <sub>1/2</sub> : DSAM in (α,pγ).
2600.397 <sup>b</sup> 25	4 <sup>+</sup>	0.55 ps 18	B DEFG K MNO Q WX	XREF: X(2573). J <sup>π</sup> : ΔJ=2, E2 γ to 2 <sup>+</sup> ; M1+E2 γ to 4 <sup>+</sup> . T <sub>1/2</sub> : unweighted average of 0.37 ps +12-7 and 0.73 ps 14 in (α,pγ) Other: >0.28 ps in (n,n'γ). XREF: D(?).
2782.14 19	1 <sup>+</sup>	0.18 ps 3	DEFG I K MN	E(level): population in ( <sup>6</sup> Li,d) is questionable. T <sub>1/2</sub> : weighted average of 0.18 ps +3-2 in (α,pγ) and 0.20 ps +9-5 in (n,n'γ). Other: 0.062 ps 17 in (n,γ) is in disagreement. J <sup>π</sup> : 1 <sup>+</sup> ,2 <sup>+</sup> from M1+E2 to 2 <sup>+</sup> and primary γ D from 0 <sup>-</sup> ,1 <sup>-</sup> neutron resonance; γ(θ) in neutron capture excludes J=2.
2864.72 12	(5)	3.1 ps 14	De n	J <sup>π</sup> : ΔJ=1, dipole γ to 4 <sup>+</sup> ; γ from (7). T <sub>1/2</sub> : from ( <sup>13</sup> C,3nγ).
2876.34 13	2 <sup>+</sup> @	0.095 ps 14	AB DeFG I K Mn T	XREF: D(?). T <sub>1/2</sub> : weighted average of 0.094 ps 14 in (α,pγ) and 0.097 ps +21-14 in (n,n'γ). Other: 0.030 ps +17-8 in (n,γ) is in disagreement.
2970 30	(5 <sup>-</sup> )		N	J <sup>π</sup> : L=(5) in (p,p').
3083.69 19	2 <sup>+</sup> @	0.031 ps 6	AB FG I K MN WX	XREF: N(3072)X(3030). T <sub>1/2</sub> : weighted average of 0.025 ps +6-4 in (α,pγ), 0.033 ps +12-8 in (n,n'γ) and 0.047 ps 9 in (n,γ).
3134 5	4 <sup>+</sup>		G N	E(level): from (p,p') with 11 keV correction added. J <sup>π</sup> : L(p,p')=4.
3233.26 6	2 <sup>+</sup>	0.22 ps 5	B F K MN	XREF: N(3222). J <sup>π</sup> : L=1+3 in (d,p). T <sub>1/2</sub> : from (α,pγ).
3243.97 23	0 <sup>+</sup> @	31 fs +67-14	AB FG I M	T <sub>1/2</sub> : from (n,γ).
3389 30	2 <sup>+</sup>		N	J <sup>π</sup> : L(p,p')=2.
3449.7 3	(4 <sup>+</sup> )	0.36 ps +13-8	D F K MN	T <sub>1/2</sub> : from (α,pγ). J <sup>π</sup> : from σ analysis of (n,n'γ); ΔJ=1 γ to 3 <sup>+</sup> .
3537.97 15	1 <sup>+</sup>	8 fs 3	FG I K M	T <sub>1/2</sub> : weighted average of 6 fs 3 in (α,pγ) and 10 fs 3 in (n,n'γ). J <sup>π</sup> : L=1, L+1/2 in (pol d,p), γγ(θ) in (n,γ); γ(circ pol) in (n,γ).
3543 5	2 <sup>+</sup>		N	E(level): from (p,p') with 11 keV correction added. J <sup>π</sup> : L(p,p')=2.
3596.90 <sup>a</sup> 14	6 <sup>+</sup>	0.20 ps 7	D F M WX	J <sup>π</sup> : ΔJ=2, E2 γ to 4 <sup>+</sup> ; band assignment. T <sub>1/2</sub> : unweighted average of 0.34 ps 4 and 0.15 ps +3-2 in (α,pγ); 0.11 ps +8-4 in (n,n'γ). Other: <3 ps in ( <sup>13</sup> C,3nγ).
3629.60 23	2 <sup>+</sup> @	8 fs 4	B FG I K MNO	J <sup>π</sup> : σ(θ) in (p,p') inconsistent with L=2 which may imply a separate level near this energy. T <sub>1/2</sub> : unweighted average of 6 fs 2 in (α,pγ), 15 fs 3 in (n,n'γ) and 2.6 fs +29-11 in (n,γ).

Continued on next page (footnotes at end of table)

**Adopted Levels, Gammas (continued)**

$^{58}\text{Fe}$ Levels (continued)						
E(level) <sup>†</sup>	$J^\pi$ <sup>‡</sup>	$T_{1/2}$ <sup>#</sup>	XREF			Comments
3754.2 4	(4) <sup>+</sup>	<0.013 ps	FG	K	W	$J^\pi$ : L=(4) in (t,p), L=1+3 in (d, <sup>3</sup> He). $T_{1/2}$ : from ( $\alpha$ ,p $\gamma$ ).
3789.49 18	(5 <sup>-</sup> ) <sup>@</sup>	0.026 ps +6-4	FG	K M		$T_{1/2}$ : from ( $\alpha$ ,p $\gamma$ ).
3854 10	2 <sup>+</sup>			K		$J^\pi$ : L=1+3 in (d,p).
3860.9 7	3 <sup>-</sup>	0.090 ps +35-21	G	LMNOPQ	X	B(E3) $\uparrow$ =0.0139 I3 XREF: N(3845)P(3800). $J^\pi$ : L(p,p')=L( <sup>3</sup> He, <sup>3</sup> He')=3. $T_{1/2}$ : from (n,n' $\gamma$ ). B(E3) from (e,e'). See also 2002Ki06 evaluation.
3880.1 3	1 <sup>+</sup>	<4 fs	F	I K M		$T_{1/2}$ : from ( $\alpha$ ,p $\gamma$ ); other: 0.7 fs 7 (n, $\gamma$ ). $J^\pi$ : 0 <sup>+</sup> ,1 <sup>+</sup> from CP of $\gamma$ 's in polarized thermal (n, $\gamma$ ) and L(d,p)=1; $\gamma$ to 0 <sup>+</sup> excludes 0 <sup>+</sup> .
3886.40 <sup>e</sup> 15	6 <sup>+</sup>	0.48 ps 10	D F	M	W	$J^\pi$ : $\Delta J=2$ , E2 $\gamma$ to 4 <sup>+</sup> ; $\Delta J=0$ $\gamma$ to 6 <sup>+</sup> ; L=3 in (d, <sup>3</sup> He). $T_{1/2}$ : weighted average of 0.49 ps +15-7 (1977Ca28) and 0.47 ps +17-11 (1978Bo35) from ( $\alpha$ ,p $\gamma$ ). Other: 11.8 ps 14 in ( <sup>13</sup> C,3n $\gamma$ ) is in severe disagreement.
3901.62 7	(3) <sup>+</sup>	0.031 ps 7	B	K M		$T_{1/2}$ : from (n,n' $\gamma$ ). $J^\pi$ : L=3 in (d,p), analysis of $\sigma$ in (n,n' $\gamma$ ).
4010.8	2 <sup>+</sup> <sup>@</sup>		G	K M		
4015.01 24	1 <sup>+</sup>	0.008 ps +4-3	F	I		$T_{1/2}$ : from ( $\alpha$ ,p $\gamma$ ). $J^\pi$ : from circular polarization of $\gamma$ 's in polarized thermal (n, $\gamma$ ).
4088.49 17	4 <sup>+</sup> <sup>@</sup>	0.06 ps +8-3	B	G MN	W	E(level), $J^\pi$ : possibly a doublet in (p,p') with L=3,4. $T_{1/2}$ : DSAM in (n,n' $\gamma$ ).
4139.24 25	1 <sup>+</sup>	2.8 fs 21	F	I K		$T_{1/2}$ : from (n, $\gamma$ ); other: <0.7 fs in ( $\alpha$ ,p $\gamma$ ). $J^\pi$ : 0 <sup>+</sup> ,1 <sup>+</sup> from CP of polarized thermal (n, $\gamma$ ) and L(d,p)=1; $\gamma$ to 0 <sup>+</sup> excludes 0 <sup>+</sup> .
4158 10	0 <sup>+</sup> <sup>@</sup>		G	K		
4214.64 <sup>c</sup> 15	(5 <sup>+</sup> )	0.45 ps +14-10	D FG	K M	U	$J^\pi$ : $\Delta J=1$ $\gamma$ to 4 <sup>+</sup> . Positive parity is tentatively proposed in ( $\alpha$ ,p $\gamma$ ) and ( <sup>13</sup> C,3n $\gamma$ ) and from shell-model predictions (1978Na06,2000ApZW). The 1997 evaluation (1997Bh02) assigned negative parity, primarily based on L(p,p')=5+3 for a 4230 30 group, but this L value gives $J^\pi=4^-$ in contradiction to J=5 from angular distribution data in ( $\alpha$ ,p $\gamma$ ) and ( <sup>13</sup> C,3n $\gamma$ ) reactions. The L(t,p)=(6) and L(d,p)=(3) suggest positive parity but implied spins are in disagreement with 5 <sup>+</sup> . For (p,p'), a separate level is now proposed. $T_{1/2}$ : from ( $\alpha$ ,p $\gamma$ ). $J^\pi$ : L(p,p')=3+5. $J^\pi$ : L=(1+3) in (d,p). $T_{1/2}$ : from (n, $\gamma$ ). $J^\pi$ : L=2 in (p,p'). $T_{1/2}$ : from (n, $\gamma$ ). $J^\pi$ : L=1+3 in (d,p). $J^\pi$ : from CP of $\gamma$ 's in polarized thermal (n, $\gamma$ ).
4230 30	4 <sup>-</sup>			N		
4237 10	(2 <sup>+</sup> )			K		
4297.8 5	2 <sup>+</sup>	2.8 fs 21	G I	K n		
4312.92 9	2 <sup>+</sup>	11 fs 7	B	K Mn	W	
4322.5 3	1 <sup>+</sup>			I		
4340 20	(5 <sup>-</sup> ,4 <sup>+</sup> ) <sup>@</sup>		G			

Continued on next page (footnotes at end of table)

**Adopted Levels, Gammas (continued)**

<sup>58</sup>Fe Levels (continued)

E(level) <sup>†</sup>	J <sup>π</sup> <sup>‡</sup>	T <sub>1/2</sub> <sup>#</sup>	XREF				Comments
4348 10	2 <sup>+</sup>			K			J <sup>π</sup> : L=1+3 in (d,p).
4350 20	(0 <sup>+</sup> ) <sup>@</sup>		G				
4352.7 7	1 <sup>+</sup>			I			J=1 from CP of γ's in polarized thermal neutron capture.
4398 10				K			
4438 10	2 <sup>-</sup> ,3 <sup>-</sup>			K	n		J <sup>π</sup> : L=2 and L+1/2 transfer in (pol d,p).
4440 20	3 <sup>-</sup> ,4 <sup>-</sup>		G		n	W	J <sup>π</sup> : L(d, <sup>3</sup> He)=0. L(t,p)=(5,4) is inconsistent with J <sup>π</sup> =3 <sup>-</sup> ,4 <sup>-</sup> .
4444.3 5	1 <sup>+</sup>	6 fs +28-6		I	M		T <sub>1/2</sub> : from (n,γ). J <sup>π</sup> : from CP of γ's in (n,γ).
4450 20	(0 <sup>+</sup> ) <sup>@</sup>		G				
4468 10	3 <sup>-</sup>		G	K	n	Q	J <sup>π</sup> : L=3 in (α,α'); L(p,p')=3 for a 4441 30 group.
4493.1 3			B	K			
4514 10	(3 <sup>+</sup> ,2 <sup>+</sup> )			K			J <sup>π</sup> : L=(3) in (d,p), shell model.
4530.15 23	1,2		AB				J <sup>π</sup> : γ to 0 <sup>+</sup> .
4550.37 24	1 <sup>+</sup>	21 fs 7		I	K	T	T <sub>1/2</sub> : from (n,γ). J <sup>π</sup> : 0 <sup>+</sup> ,1 <sup>+</sup> from CP of γ's in polarized thermal neutron capture; L=1, L-1/2 in (pol d,p); γ to 0 <sup>+</sup> excludes 0 <sup>+</sup> .
4590.0 4	(2 <sup>+</sup> ,3,4 <sup>+</sup> )		B	K			J <sup>π</sup> : γ's to 2 <sup>+</sup> and 4 <sup>+</sup> .
4610 20	3 <sup>-</sup> ,4 <sup>-</sup>					VW	XREF: V(4700). J <sup>π</sup> : L=0 in (d, <sup>3</sup> He) and (p,2p).
4620 10	2 <sup>+</sup> <sup>@</sup>		G	K			
4661 10				K			
4669.38 <sup>C</sup> 14	(7 <sup>+</sup> )	0.38 ps +12-6	D	FGH			E(level): unresolved doublet in (α, <sup>2</sup> He) at 4650 50 with (7 <sup>-</sup> and 5 <sup>-</sup> ) from DWBA analysis. L(t,p)=2+8 for E=4670. J <sup>π</sup> : ΔJ=2 γ, E2 γ to (5 <sup>+</sup> ); ΔJ=1 γ to 6 <sup>+</sup> . T <sub>1/2</sub> : from (α,pγ). J <sup>π</sup> : L=(1+3) in (d,p), L=1+3 in (d, <sup>3</sup> He).
4711 10	(2 <sup>+</sup> )			K		W	
4720 20	1 <sup>-</sup> <sup>@</sup>		G				
4809 10	(5 <sup>-</sup> )		G	K			J <sup>π</sup> : L(t,p)=6,(5); L(d,p)=5 needed to give a J <sup>π</sup> =6 <sup>+</sup> is considered unlikely.
4833.89 25	1 <sup>+</sup> ,2 <sup>+</sup>		B	I	K	W	J <sup>π</sup> : primary γ from 0 <sup>-</sup> ,1 <sup>-</sup> neutron resonance. γ to 3 <sup>+</sup> . L(d, <sup>3</sup> He)=3.
4890 20	2 <sup>+</sup> <sup>@</sup>		G				
4937 10	2 <sup>+</sup> <sup>@</sup>		G	K		W	J <sup>π</sup> : L(t,p)=2 for E=4960 20.
4990	(2 <sup>+</sup> ,3 <sup>-</sup> )		G				J <sup>π</sup> : L(t,p)=2,(3).
5000.23 18	1 <sup>+</sup>	3.0 fs 10		I	K		XREF: K(4992). J <sup>π</sup> : 0 <sup>+</sup> ,1 <sup>+</sup> from CP of γ's in polarized thermal (n,γ); L=1, L-1/2 in (pol d,p); γ to 0 <sup>+</sup> excludes 0 <sup>+</sup> . T <sub>1/2</sub> : from (n,γ).
5020 20	5 <sup>-</sup> <sup>@</sup>		G				
5060 20	2 <sup>+</sup> <sup>@</sup>		G			W	
5138 10	0 <sup>+</sup> <sup>@</sup>		G	K			
5164 10				K		w	
5213 10	2 <sup>+</sup> <sup>@</sup>		G	K		w	
5220.9 5	1,2	<0.38 ps		I	M	w	T <sub>1/2</sub> : from (n,n'γ); other: <2.4 fs in (n,γ). J <sup>π</sup> : dipole γ from 0 <sup>-</sup> ,1 <sup>-</sup> (n,γ) resonance; γ to 0 <sup>+</sup> excludes J=0.

Continued on next page (footnotes at end of table)

**Adopted Levels, Gammas (continued)**

$^{58}\text{Fe}$ Levels (continued)					
E(level) <sup>†</sup>	$J^\pi$ <sup>‡</sup>	$T_{1/2}$ <sup>#</sup>	XREF		Comments
5236 10				K	w
5254 10	3 <sup>-</sup> @		G	K	
5294.8 6	(1 <sup>+</sup> ,2,3 <sup>+</sup> )	3.5 fs 28	G I	K	$T_{1/2}$ : from (n, $\gamma$ ). $J^\pi$ : $\gamma$ 's to 1 <sup>+</sup> and 3 <sup>+</sup> .
5315 10	3 <sup>-</sup> ,4 <sup>-</sup>			K	W
5343.33 <sup>e</sup> 22	8 <sup>+</sup>	0.42 ps +10-8	D F		$J^\pi$ : L=0+2 in (d, <sup>3</sup> He). $J^\pi$ : $\Delta J=2$ E2 $\gamma$ to 6 <sup>+</sup> ; band assignment. $T_{1/2}$ : from ( $\alpha$ ,p $\gamma$ ).
5370 10	(4 <sup>+</sup> ,5 <sup>-</sup> )@		G	K	x
5400 50	-				Wx
5406 10	0 <sup>+</sup> @		G	K	
5417.6 6	(1 <sup>+</sup> ,2,3 <sup>-</sup> )	<0.7 fs	I		$J^\pi$ : $\gamma$ 's to 2 <sup>+</sup> and 3 <sup>+</sup> ; primary $\gamma$ from 0 <sup>-</sup> ,1 <sup>-</sup> . E(level): 5418.1 keV obtained from the 4626.5 5 primary neutron capture $\gamma$ ray populating this level is discrepant with the level energy from a least-squares fit.
5462 10	(2 <sup>+</sup> )@		G	K	
5502.9 <sup>a</sup> 10	(8 <sup>+</sup> )	<0.14 ps	D F		$J^\pi$ : $\gamma$ to 6 <sup>+</sup> ; band assignment. $T_{1/2}$ : from ( <sup>13</sup> C,3n $\gamma$ ).
5506 10				K	
5523.0 22	0 <sup>+</sup> @		G I	K	
5620 10	0 <sup>+</sup> @		G	K	
5655 10	2 <sup>+</sup> @		G	K	W
5716 10	3 <sup>-</sup> ,4 <sup>-</sup>			K	W
5734 10	2 <sup>+</sup> @		G	K	
5763 10				K	
5788 10	(2 <sup>+</sup> ,3 <sup>-</sup> )@		G	K	
5817 10	(2 <sup>-</sup> ,3 <sup>-</sup> )			K	$J^\pi$ : L=(2) in (d,p), shell model.
5830 20	0 <sup>+</sup> @		G		
5832.08 <sup>c</sup> 23	(9 <sup>+</sup> )	0.40 ps +15-4	D F		$J^\pi$ : $\Delta J=2$ , E2 $\gamma$ to (7 <sup>+</sup> ). $T_{1/2}$ : from 1977Ca28 in ( $\alpha$ ,p $\gamma$ ). Other: 0.8 ps 3 from ( <sup>13</sup> C,3n $\gamma$ ). $J^\pi$ : L=(2) in (d,p).
5857 10	(2 <sup>-</sup> ,3 <sup>-</sup> )			K	
5880 20	(2 <sup>+</sup> ,3 <sup>-</sup> )@		G		
5887 10	(0 <sup>-</sup> ,1 <sup>-</sup> )			K	$J^\pi$ : L=(0) in (d,p).
5914 10				K	
5952 10	(2 <sup>+</sup> )@		G	K	
5989 10				K	
6030 10				K	
6032.9 <sup>d</sup> 5	(9 <sup>+</sup> )		D		$J^\pi$ : $\Delta J=2$ $\gamma$ to (7 <sup>+</sup> ); band assignment.
6054 10			G	K	E(level): possible doublet in (t,p), (pol t,p).
6100 50	3 <sup>-</sup> ,4 <sup>-</sup>				W
6146 10	2 <sup>+</sup> @		G	K	$J^\pi$ : L=0 in (d, <sup>3</sup> He).
6168 10	(0 <sup>+</sup> )@		G	K	
6202 10	3 <sup>-</sup> ,4 <sup>-</sup>			K	W
6238 10	(1 <sup>-</sup> ,2 <sup>+</sup> )@		G	K	
6279 10	(1 <sup>-</sup> ,2 <sup>+</sup> )@		G	K	
6282.7 <sup>e</sup> 5	(9 <sup>+</sup> )	<0.14 ps	D		$J^\pi$ : $\Delta J=1$ $\gamma$ to 8 <sup>+</sup> ; band assignment. $T_{1/2}$ : from ( <sup>13</sup> C,3n $\gamma$ ).

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) $^{58}\text{Fe}$  Levels (continued)

E(level) <sup>†</sup>	J <sup>π</sup> <sup>‡</sup>	T <sub>1/2</sub> <sup>#</sup>	XREF		Comments
6295 10	(5 <sup>-</sup> )		H	K	J <sup>π</sup> : from DWBA analysis in ( $\alpha$ , <sup>2</sup> He).
6328 10			G	K	E(level): possible doublet in (t,p), (pol t,p).
6348 10				K	
6370 10				K	
6400 10	(6 <sup>+</sup> ,7 <sup>-</sup> ) <sup>@</sup>		G	K	
6436 10	1 <sup>-</sup> <sup>@</sup>		G	K	
6450 10	0 <sup>+</sup> <sup>@</sup>		G	K	
6476 10				K	
6532 10				K	
6558 10				K	
6580 20	(6 <sup>+</sup> ) <sup>@</sup>		G		
6593 10				K	
6615 10				K	
6636 10				K	
6650 20	0 <sup>+</sup> <sup>@</sup>		G		
6679 10	(3 <sup>-</sup> ,2 <sup>-</sup> )			K	J <sup>π</sup> : L=(2) in (d,p), shell model.
6741 10				K	
6760 20	0 <sup>+</sup> <sup>@</sup>		G		
6771 10				K	
6789 10				K	
6842 10			G	K	
6870 20	(5 <sup>-</sup> ) <sup>@</sup>		G		
6909 10	1 <sup>-</sup> <sup>@</sup>		G	K	
6953 10	2 <sup>+</sup> <sup>@</sup>		G	K	
7023 10				K	
7028 10				K	
7048 10	(1 <sup>-</sup> ,2 <sup>+</sup> ) <sup>@</sup>		G	K	
7060 10				K	
7094 10				K	
7124 10	0 <sup>+</sup> <sup>@</sup>		G	K	
7166 10	1 <sup>-</sup> <sup>@</sup>		G	K	
7199 10				K	
7230 10				K	
7242.6 <sup>e</sup> 9	(10 <sup>+</sup> )	<0.14 ps	D		J <sup>π</sup> : $\Delta J=1$ $\gamma$ to (9 <sup>+</sup> ); band assignment. T <sub>1/2</sub> : from ( <sup>13</sup> C,3n $\gamma$ ).
7272 10				K	
7289 10				K	
7351 10				K	
7380 50	(8 <sup>+</sup> )		H		J <sup>π</sup> : from analysis of $\sigma$ in ( $\alpha$ , <sup>2</sup> He).
7429 10	(0 <sup>-</sup> ,1 <sup>-</sup> )			K	J <sup>π</sup> : L=(0) in (d,p).
7456.7 <sup>d</sup> 5	(10 <sup>+</sup> )		D		J <sup>π</sup> : $\Delta J=1$ $\gamma$ to (9 <sup>+</sup> ); band assignment.
7457 10				K	
7473 10				K	
7492 10				K	
7507 10				K	
7534 10				K	
7567 10				K	
7578 10				K	
7585 10				K	
7605 10				K	
7628 10				K	
7653 10				K	

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) $^{58}\text{Fe}$  Levels (continued)

E(level) <sup>†</sup>	J <sup>π</sup> <sup>‡</sup>	T <sub>1/2</sub> <sup>#</sup>	XREF	Comments
7680? 10			K	
7690? 10			K	
7731.3 <sup>c</sup> 5	(11 <sup>+</sup> )	<0.14 ps	D	J <sup>π</sup> : ΔJ=2, (E2) γ to (9 <sup>+</sup> ); band assignment. T <sub>1/2</sub> : from ( <sup>13</sup> C,3nγ).
7734 10			K	
7775 10			K	
7797 10			K	
7824 10			K	
7846 10			K	
7883 10			K	
7901 10			K	
7918 10			K	
7946 10			K	
7974 10			K	
7997 10			K	
8018 10			K	
8045 10			K	
8065 10			K	
8084 10			K	
8100 10			K	
8121 10			K	
8137 10			K	
8157 10			K	
8182 10			K	
8310 50	(6 <sup>+</sup> )		H	J <sup>π</sup> : from analysis of σ in (α, <sup>2</sup> He).
9444.8 <sup>d</sup> 6	(12 <sup>+</sup> )		D	J <sup>π</sup> : ΔJ=2 γ to (10 <sup>+</sup> ); band assignment.
9939.1 9			D	J <sup>π</sup> : γ to (11 <sup>+</sup> ) suggests (11,12,13 <sup>+</sup> ).
9984.5 7	(12)		D	J <sup>π</sup> : ΔJ=1 γ to (11 <sup>+</sup> ).
10041.05 18	1- <sup>&amp;</sup>		J	
(10044.31 19)	1 <sup>-</sup>		I	J <sup>π</sup> ,E(level): for s-wave capture on <sup>57</sup> Fe(J <sup>π</sup> =1/2 <sup>-</sup> ). S(n)=10044.60 18 (2009AuZZ).
10046.20 18	2+ <sup>&amp;</sup>		J	
10048.48 18	0- <sup>&amp;</sup>		J	
10049.26 18	+ <sup>&amp;</sup>		J	
10050.71 18	1- <sup>&amp;</sup>		J	
10051.69 18	(+) <sup>&amp;</sup>		J	
10052.40 18	(+) <sup>&amp;</sup>		J	
10052.97 18	(+) <sup>&amp;</sup>		J	
10053.64 18	(+) <sup>&amp;</sup>		J	
10056.48 18	(+) <sup>&amp;</sup>		J	
10057.17 18	(+) <sup>&amp;</sup>		J	
10057.68 18	(+) <sup>&amp;</sup>		J	
10058.30 18	1- <sup>&amp;</sup>		J	
10058.49 18			J	
10062.34 18	(+) <sup>&amp;</sup>		J	
10062.52 18	(-) <sup>&amp;</sup>		J	
10062.98 18	(+) <sup>&amp;</sup>		J	
10065.28 18	(+) <sup>&amp;</sup>		J	
10065.52 18	(-) <sup>&amp;</sup>		J	
10065.6 3			J	

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) $^{58}\text{Fe}$  Levels (continued)

E(level) <sup>†</sup>	J <sup>π</sup> <sup>‡</sup>	T <sub>1/2</sub> <sup>#</sup>	XREF	Comments
10069.94 18			J	
10071.33 18	(+)&		J	
10072.73 18	(-)&		J	
10073.14 19	1-&		J	
10075.4 <sup>c</sup> 9	(13 <sup>+</sup> )	<0.14 ps	D	J <sup>π</sup> : ΔJ=2, (E2) γ to (11 <sup>+</sup> ); band assignment. T <sub>1/2</sub> : from ( <sup>13</sup> C,3nγ).
10075.98 18	(+)&		J	
10079.18 18			J	
10081.07 18	(+)&		J	
10081.83 18	(+)&		J	
10083.30 18	(-)&		J	
10083.71 19	(+)&		J	
10085.28 20	1-&		J	
10085.8 18	(+)&		J	
10086.2 18			J	
10087.4 19			J	
10090.8 18	1-&		J	
10093.66 19	(+)&		J	
10094.64 19			J	
10096.39 19			J	
10096.56 19	(-)&		J	
10099.44 19	0-&		J	
10099.80 19	(+)&		J	
10102.40 19			J	
10102.51 18			J	
10104.59 22	1-&		J	
10105.53 19			J	
10105.77 19			J	
10106.49 19			J	
10107.44 19			J	
10107.71 19			J	
10110.19 19			J	
10111.48 19			J	
10114.80 19			J	
10116.03 19			J	
10117.60 18			J	
10120.16 23	1-&		J	
10123.50 20			J	
10126.30 20			J	
10127.60 20			J	
10130.32 20			J	
10131.34 20			J	
10133.01 20			J	
10134.35 20			J	
10136.36 20			J	
10136.67 20	1-&		J	
10136.93 20			J	
10137.65 20			J	
10139.22 20			J	
10140.99 20			J	
10141.73 20			J	

Continued on next page (footnotes at end of table)



Adopted Levels, Gammas (continued) $^{58}\text{Fe}$  Levels (continued)

E(level) <sup>†</sup>	J <sup>π</sup> <sup>‡</sup>	XREF	Comments
10142.73 20		J	
10143.94 20		J	
10144.63 20	(+)&	J	
10146.76 20		J	
10147.98 21		J	
10148.63 21		J	
10149.57 21		J	
10150.33 21		J	
10150.82 21		J	
10152.1 3	1- &	J	
10152.9 3	1- &	J	
10153.76 21		J	
10155.43 21		J	
10156.20 21		J	
10157.10 22		J	
10161.72 22	(+)&	J	
10163.98 22		J	
10166.42 22		J	
10167.4 3	1- &	J	
10168.4 3	0- &	J	
10169.09 22		J	
10171.84 22	1- &	J	
10172.53 22		J	
10174.10 22		J	
10176.36 22		J	
10176.8 3	0- &	J	
10177.52 22		J	
10182.9 3	0- &	J	
10190.81 23		J	
10192.23 23		J	
10192.68 23		J	
10196.87 23		J	
10200.15 24		J	
10201.72 24		J	
10206.53 25		J	
10208.23 25		J	
10208.7 4	1- &	J	
10208.99 25		J	
10210.46 25		J	
10210.66 25	1- &	J	
10210.97 23		J	
10217.83 25	0- &	J	
10221.37 25		J	
10227.1 4	1- &	J	
10228.15 3		J	
10230.8 3		J	
10234.9 3		J	
10238.4 3		J	
10240.0 3		J	
10241.2 3		J	
10353.8 9		D	J <sup>π</sup> : γ to (11 <sup>+</sup> ) suggests (11,12,13 <sup>+</sup> ).
11857.0 <sup>d</sup> 8	(14 <sup>+</sup> )	D	J <sup>π</sup> : γ to (12 <sup>+</sup> ); band assignment.

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) $^{58}\text{Fe}$  Levels (continued)

E(level) <sup>†</sup>	$J^\pi$ <sup>‡</sup>	XREF	Comments
11911.0 9		D	$J^\pi$ : $\gamma$ to (12) suggests (12,13,14).
12813.3 <sup>c</sup> 16	(15 <sup>+</sup> )	D	$J^\pi$ : $\gamma$ to (13 <sup>+</sup> ); band assignment.

<sup>†</sup> From a least-squares fit to  $E\gamma$ 's for levels populated in  $\gamma$ -ray studies. For levels populated in particle-transfer and/or inelastic scattering studies, the values are averaged over all available data. In addition poorly resolved groups are reported at 2.94, 3.24, 4.11, 4.75, 5.25, 5.68, 6.23 and 6.55 MeV with an uncertainty of 0.12 MeV in  $^{59}\text{Co}(\gamma,p)$ . These are not included in cross reference (XREF) table.

<sup>‡</sup> In in-beam  $\gamma$ -ray studies:  $^{55}\text{Mn}(\alpha,p\gamma)$  and  $^{48}\text{Ca}(^{13}\text{C},3n\gamma)$ , ascending order of spins are assumed as the excitation energy rises. When  $J^\pi$  is deduced from L-transfers in particle transfer reactions, the target  $J^\pi$ 's are as follows: 1/2<sup>-</sup> for  $^{57}\text{Fe}$  in (d,p) reaction; 7/2<sup>-</sup> for  $^{59}\text{Co}$  in (d, $^3\text{He}$ ); 0<sup>+</sup> in ( $^6\text{Li}$ ,d), (t,p) and ( $^3\text{He}$ , $^7\text{Be}$ ) reactions. The abbreviation CP in (n, $\gamma$ ) indicates circular polarization measurement.

<sup>#</sup> For excited states, most values are from DSAM in the following reactions:  $^{55}\text{Mn}(\alpha,p\gamma)$ ;  $^{57}\text{Fe}(n,\gamma)$  E=th and  $^{58}\text{Fe}(n,n'\gamma)$ . Selected values are also available from DSAM and recoil-distance method in  $^{48}\text{Ca}(^{13}\text{C},3n\gamma)$ .

@ From L(t,p).

& From L-value in neutron resonances. See  $^{57}\text{Fe}(n,\gamma),(n,n)$ :resonances.

<sup>a</sup> Band(A): yrast band.

<sup>b</sup> Band(B): Band based on 2<sup>+</sup>.

<sup>c</sup> Band(C): band based on 5<sup>+</sup>.

<sup>d</sup> Band(D): band based on 9<sup>(+)</sup>.

<sup>e</sup> Band(E): band based on 6<sup>+</sup>.

## Adopted Levels, Gammas (continued)

E <sub>i</sub> (level)	J <sub>i</sub> <sup>π</sup>	γ( <sup>58</sup> Fe)		E <sub>f</sub>	J <sub>f</sub> <sup>π</sup>	Mult. <sup>†</sup>	γ( <sup>58</sup> Fe)		Comments
		E <sub>γ</sub> <sup>‡</sup>	I <sub>γ</sub> <sup>‡</sup>				δ	α <sup>@</sup>	
810.7662	2 <sup>+</sup>	810.7593 20	100	0.0	0 <sup>+</sup>	E2		3.32×10 <sup>-4</sup>	B(E2)(W.u.)=18.5 6 Mult.: from γγ(θ), γ(θ), RUL and measured α(K)exp.
1674.731	2 <sup>+</sup>	863.951 6	100	810.7662	2 <sup>+</sup>	M1+E2	-0.69 5		B(M1)(W.u.)=0.0082 21; B(E2)(W.u.)=10 3 Mult.,δ: D+Q from γγ(θ), γ(θ), RUL. δ: from <sup>58</sup> Co ε decay. Others: -0.57 6 (n,γ), -0.50 5 (n,n'γ). B(E2)(W.u.)=0.87 22 B(E2)(W.u.)=47 7
2076.52	4 <sup>+</sup>	1674.725 10 1265.74 5	76.4 15 100	0.0 810.7662	0 <sup>+</sup> 2 <sup>+</sup>	[E2] E2			Mult.: from γ(θ) in (α,pγ) and ( <sup>13</sup> C,3nγ) and RUL. B(M1)(W.u.)=0.027 9 B(E2)(W.u.)=0.87 22 B(E2)(W.u.)=47 7
2133.895	3 <sup>+</sup>	459.160 25 1323.09 5	36 1 100 3	1674.731 810.7662	2 <sup>+</sup> 2 <sup>+</sup>	(M1) M1+E2	-0.40 5		Mult.: from γ(θ) in (n,n'γ). Other: -0.48 +12-10 (n,γ). B(M1)(W.u.)=0.0027 9; B(E2)(W.u.)=0.48 19 δ: from γ(θ) in (n,n'γ). Other: -0.48 +12-10 (n,γ). Mult.: from γ(θ) and RUL.
2257.95	0 <sup>+</sup>	1447.31 25	100	810.7662	2 <sup>+</sup>	[E2]			B(E2)(W.u.)<2.7
2600.397	4 <sup>+</sup>	466.48 3 523.86 3	34.3 12 100 3	2133.895 2076.52	3 <sup>+</sup> 4 <sup>+</sup>	(M1) M1+E2	-0.15 5		B(M1)(W.u.)=0.053 18 B(M1)(W.u.)=0.11 4; B(E2)(W.u.)=17 13 Mult.: from γ(θ) in (n,n'γ) and RUL. δ: from (n,n'γ). Other: +6.3 in (α,pγ); mult=Q in ( <sup>13</sup> C,3nγ). B(E2)(W.u.)=20 7
		925.68 5	45.5 17	1674.731	2 <sup>+</sup>	E2			Mult.: from γ(θ) in (α,pγ), ( <sup>13</sup> C,3nγ) and RUL. B(E2)(W.u.)=1.3 5
		1789.59 8	77.4 24	810.7662	2 <sup>+</sup>	E2			Mult.: from γ(θ) in (α,pγ), ( <sup>13</sup> C,3nγ) and RUL.
2782.14	1 <sup>+</sup>	524.4 3 1106.7 3	16.4 8 47 3	2257.95 1674.731	0 <sup>+</sup> 2 <sup>+</sup>	M1+E2	-0.18 3		Mult.: from γ(θ) in (α,pγ), ( <sup>13</sup> C,3nγ) and RUL. B(M1)(W.u.)=0.020 4; B(E2)(W.u.)=1.0 4 Mult.: from γ(θ) in (n,γ) and RUL. δ: from γ(θ) in (n,γ).
		1971.6 5	100 8	810.7662	2 <sup>+</sup>	M1+E2	-0.17 4		B(M1)(W.u.)=0.0074 15; B(E2)(W.u.)=0.11 6 Mult.: from γ(θ) in (n,γ) and RUL. δ: from γ(θ) in (n,γ).
2864.72	(5)	2781.9 9 264.36 12	47 5 100	0.0 2600.397	0 <sup>+</sup> 4 <sup>+</sup>	[M1] D			B(M1)(W.u.)=0.0013 3 Mult.: from γ(θ) in ( <sup>13</sup> C,3nγ).
2876.34	2 <sup>+</sup>	2065.59 14	100 8	810.7662	2 <sup>+</sup>	M1+E2	-0.33 +8-11		B(M1)(W.u.)=0.022 5; B(E2)(W.u.)=1.1 6 δ: from (n,γ); -0.13 3 in (n,n'γ). Mult.: from γ(θ) in (n,γ) and RUL.
3083.69	2 <sup>+</sup>	2876.3 <sup>#b</sup> 2272.99 23	≤17 <sup>#</sup> 100	0.0 810.7662	0 <sup>+</sup> 2 <sup>+</sup>	M1+E2	-0.05 1		Mult.: from γ(θ) in (n,γ) and RUL. B(M1)(W.u.)=0.052 13; B(E2)(W.u.)=0.048 23 Mult.: from γ(θ) in (n,n'γ) and RUL. δ: from (n,n'γ).
3233.26	2 <sup>+</sup>	3083.6 <sup>#b</sup> 632.71 10 1156.77 7	≤33 <sup>#</sup> 50 5 94 4	0.0 2600.397 2076.52	0 <sup>+</sup> 4 <sup>+</sup> 4 <sup>+</sup>				

Adopted Levels, Gammas (continued)γ(<sup>58</sup>Fe) (continued)

<u>E<sub>i</sub>(level)</u>	<u>J<sub>i</sub><sup>π</sup></u>	<u>E<sub>γ</sub><sup>‡</sup></u>	<u>I<sub>γ</sub><sup>‡</sup></u>	<u>E<sub>f</sub></u>	<u>J<sub>f</sub><sup>π</sup></u>	<u>Mult.<sup>†</sup></u>	<u>δ</u>	<u>Comments</u>
3233.26	2 <sup>+</sup>	1558.71 19	46 2	1674.731	2 <sup>+</sup>			
		2422.45 17	100 2	810.7662	2 <sup>+</sup>			
		3233.2 <sup>#b</sup>	≤2.4 <sup>#</sup>	0.0	0 <sup>+</sup>			
3243.97	0 <sup>+</sup>	2433.05 25	100	810.7662	2 <sup>+</sup>	[E2]		B(E2)(W.u.)=16 +8-16
3449.7	(4 <sup>+</sup> )	849.7 4	100 16	2600.397	4 <sup>+</sup>			
		1315.6 4	45 7	2133.895	3 <sup>+</sup>	(M1)		B(M1)(W.u.)=0.0083 +25-35
		1373 <sup>b</sup>		2076.52	4 <sup>+</sup>			Weak γ ray.
3537.97	1 <sup>+</sup>	1862.2 5	22 3	1674.731	2 <sup>+</sup>	M1+E2	-0.59 +14-11	B(M1)(W.u.)=0.047 20; B(E2)(W.u.)=9 5 Mult.: from γ(θ) in (n,γ) and RUL. δ: from (n,γ).
		2727.24 16	100 10	810.7662	2 <sup>+</sup>	M1+E2	-0.57 +7-5	B(M1)(W.u.)=0.07 3; B(E2)(W.u.)=6 3 Mult.: γ(θ) in (n,γ) and RUL. δ: from (n,γ).
3596.90	6 <sup>+</sup>	3540 3	25 4	0.0	0 <sup>+</sup>	[M1]		B(M1)(W.u.)=0.011 5
		1520.45 20	100 4	2076.52	4 <sup>+</sup>	E2		B(E2)(W.u.)=26 10 Mult.: from γ(θ) in (α,pγ) and ( <sup>13</sup> C,3nγ) and RUL.
3629.60	2 <sup>+</sup>	2818.5& 3	100& 3	810.7662	2 <sup>+</sup>			
		3629.8 4	8.3 21	0.0	0 <sup>+</sup>			
3754.2	(4 <sup>+</sup> )	1677.7 4	100	2076.52	4 <sup>+</sup>			
3789.49	(5 <sup>-</sup> )	1712.94 17	100	2076.52	4 <sup>+</sup>			
3860.9	3 <sup>-</sup>	2186.0	100	1674.731	2 <sup>+</sup>			
		(3860.8)		0.0	0 <sup>+</sup>	[E3]		B(E3)(W.u.)=9.9 9
3880.1	1 <sup>+</sup>	1097.4 3	25 6	2782.14	1 <sup>+</sup>			
		3071 2	100 19	810.7662	2 <sup>+</sup>	(M1+E2)	+0.15 9	B(M1)(W.u.)>0.085 δ: from (n,γ).
3886.40	6 <sup>+</sup>	3881.4 7	88 19	0.0	0 <sup>+</sup>	[M1]		B(M1)(W.u.)>0.039
		289.49 12	55 3	3596.90	6 <sup>+</sup>	(M1(+E2))	<0.14	B(M1)(W.u.)>0.46 δ: deduced by the evaluators by requiring RUL(E2)=300. Not given in ( <sup>13</sup> C,3nγ).
		437.9 11	9	3449.7	(4 <sup>+</sup> )			
		1285.4 3	10.0 11	2600.397	4 <sup>+</sup>			
		1810.3 7	100.0 18	2076.52	4 <sup>+</sup>	E2		B(E2)(W.u.)=2.6 6 Mult.: Q in ( <sup>13</sup> C,3nγ); E2 from RUL.
3901.62	(3 <sup>+</sup> )	1301.10 11	22 1	2600.397	4 <sup>+</sup>			
		1767.74 8	100 4	2133.895	3 <sup>+</sup>			
		1825.1 <sup>#b</sup>	≤1.1 <sup>#</sup>	2076.52	4 <sup>+</sup>			
		2226.88 18	9.4 22	1674.731	2 <sup>+</sup>			
		3090.7 4	3.1 6	810.7662	2 <sup>+</sup>			
		3901.5 <sup>#b</sup>	≤0.6 <sup>#</sup>	0.0	0 <sup>+</sup>			
4015.01	1 <sup>+</sup>	3204.10 26	100	810.7662	2 <sup>+</sup>			

## Adopted Levels, Gammas (continued)

$\gamma(^{58}\text{Fe})$ (continued)								
$E_i(\text{level})$	$J_i^\pi$	$E_\gamma^{\ddagger}$	$I_\gamma^{\ddagger}$	$E_f$	$J_f^\pi$	Mult. <sup>†</sup>	$\delta$	Comments
4088.49	4 <sup>+</sup>	458.5 <sup>b</sup> 3 1488.17 20 2011.7 3	100 22 <100	3629.60 2600.397 2076.52	2 <sup>+</sup> 4 <sup>+</sup> 4 <sup>+</sup>			
4139.24	1 <sup>+</sup>	4139.1 3	100	0.0	0 <sup>+</sup>	[M1]		B(M1)(W.u.)=0.11 9
4214.64	(5 <sup>+</sup> )	1614.16 21 2138.2 4	100 6 23 7	2600.397 2076.52	4 <sup>+</sup> 4 <sup>+</sup>	D D		
4297.8	2 <sup>+</sup>	3486 3 4298.1 6	100 20 100 20	810.7662 0.0	2 <sup>+</sup> 0 <sup>+</sup>	[E2]		B(E2)(W.u.)=5 4
4312.92	2 <sup>+</sup>	1436.5 <sup>#b</sup> 1712.21 26 2179.08 14 2236.33 15 2638.15 20 3501.9 8 4312.7 <sup>#b</sup>	$\leq 2.6^{\#}$ 5.3 20 36 3 26 1 100 3 1.3 13 $\leq 1.3^{\#}$	2876.34 2600.397 2133.895 2076.52 1674.731 810.7662 0.0	2 <sup>+</sup> 4 <sup>+</sup> 3 <sup>+</sup> 4 <sup>+</sup> 2 <sup>+</sup> 2 <sup>+</sup> 0 <sup>+</sup>			
4322.5	1 <sup>+</sup>	1238.7 7 1446.3 4 4322.1 6	5.7 29 100 9 60 11	3083.69 2876.34 0.0	2 <sup>+</sup> 2 <sup>+</sup> 0 <sup>+</sup>			
4444.3	1 <sup>+</sup>	1662.5 6 4443 2	100 22 78 22	2782.14 0.0	1 <sup>+</sup> 0 <sup>+</sup>			
4493.1		2818.5 <sup>&amp;</sup> 3 3681.7 5	100 <sup>&amp;</sup> 8.3	1674.731 810.7662	2 <sup>+</sup> 2 <sup>+</sup>			
4530.15	1,2	1446.53 27 2855.2 3 4531.0 15	100 18 64 9 36 18	3083.69 1674.731 0.0	2 <sup>+</sup> 2 <sup>+</sup> 0 <sup>+</sup>			
4550.37	1 <sup>+</sup>	410.9 5 1306.0 5 1674.2 3  2876 2	1.40 18 14.0 18 67 25 100 11	4139.24 3243.97 2876.34 1674.731	1 <sup>+</sup> 0 <sup>+</sup> 2 <sup>+</sup> 2 <sup>+</sup>	(M1+E2)	+0.17 +10-9	B(M1)(W.u.)=0.08 4; B(E2)(W.u.)=1.5 +20-15 I $\gamma$ : from (n, $\gamma$ ) where this $\gamma$ is multiply placed and undivided intensity is given. $\delta$ : from (n, $\gamma$ ). B(M1)(W.u.)=0.021 9; B(E2)(W.u.)=0.48 24 I $\gamma$ (2876)/I $\gamma$ (1674)=0.07 9 in ( $\mu^-n\gamma$ ) is in severe disagreement with adopted ratio of 1.5 6. $\delta$ : from (n, $\gamma$ ).
4590.0	(2 <sup>+</sup> ,3,4 <sup>+</sup> )	3740 <sup>a</sup> 3 2513.9 <sup>b</sup> 4 3778.1 6	$\approx 5^a$ 35 12 100 12	810.7662 2076.52 810.7662	2 <sup>+</sup> 4 <sup>+</sup> 2 <sup>+</sup>			
4669.38	(7 <sup>+</sup> )	454.73 14	33.2 19	4214.64	(5 <sup>+</sup> )	E2		B(E2)(W.u.)=1000 +107-330 is larger than RUL by a factor of 2 to 4. This suggests that either the reported $T_{1/2}$ is too small or branching is too large. Note that this $\gamma$ is not reported in ( $\alpha,\gamma$ ).

Adopted Levels, Gammas (continued)

$\gamma(^{58}\text{Fe})$ (continued)								
$E_i(\text{level})$	$J_i^\pi$	$E_\gamma^{\ddagger}$	$I_\gamma^{\ddagger}$	$E_f$	$J_f^\pi$	Mult. <sup>†</sup>	$\delta$	Comments
4669.38	(7 <sup>+</sup> )	782.84 16	100.0 25	3886.40	6 <sup>+</sup>	(M1(+E2))	-0.06 +16-10	B(M1)(W.u.)=0.063 +11-20; B(E2)(W.u.)=1 +4-1 $\delta$ : from ( $\alpha$ ,p $\gamma$ ).
		1072.55 17	48 3	3596.90	6 <sup>+</sup>	(M1+E2)	-0.10 +20-15	B(M1)(W.u.)=0.0117 +21-38; B(E2)(W.u.)=0.2 +8-2 $\delta$ : from ( $\alpha$ ,p $\gamma$ ).
		1219 <sup>b</sup>		3449.7	(4 <sup>+</sup> )			$I_\gamma$ : very weak, observed only in ( $\alpha$ ,p $\gamma$ ). Not reported in ( $\alpha$ ,p $\gamma$ ).
4833.89	1 <sup>+</sup> ,2 <sup>+</sup>	1804.9 3	9.6 9	2864.72	(5)			
5000.23	1 <sup>+</sup>	2699.94 25	100	2133.895	3 <sup>+</sup>			
5220.9	1,2	3326 2	100 9	1674.731	2 <sup>+</sup>	(M1+(E2))	-0.02 4	B(M1)(W.u.)=(0.15 6); B(E2)(W.u.)=(0.011 +43-11) $\delta$ : from (n, $\gamma$ ).
		4189.2 2	5.3 18	810.7662	2 <sup>+</sup>			
		5001.0 7	25 4	0.0	0 <sup>+</sup>			
		2137.6 7	6.6 19	3083.69	2 <sup>+</sup>			
		4411 3	4.7 19	810.7662	2 <sup>+</sup>			
5294.8	(1 <sup>+</sup> ,2,3 <sup>+</sup> )	5223 3	100 3	0.0	0 <sup>+</sup>			
		2513.5 10	100 19	2782.14	1 <sup>+</sup>			
		3162 3	88 19	2133.895	3 <sup>+</sup>			
5343.33	8 <sup>+</sup>	4483 2	31 13	810.7662	2 <sup>+</sup>			
		672 <sup>b</sup>	≈28	4669.38	(7 <sup>+</sup> )			
		1456.90 20	100 6	3886.40	6 <sup>+</sup>	E2		B(E2)(W.u.)=9.3 +21-25 Mult.: from $\gamma(\theta)$ in ( <sup>13</sup> C,3n $\gamma$ ) and RUL.
5417.6	(1 <sup>+</sup> ,2,3 <sup>-</sup> )	1746.4 3	37 3	3596.90	6 <sup>+</sup>	E2		B(E2)(W.u.)=1.4 4 Mult.: from $\gamma\gamma(\theta)$ in ( <sup>13</sup> C,3n $\gamma$ ) and RUL.
		3280 3	38 25	2133.895	3 <sup>+</sup>			
		3740 <sup>a</sup> 3	100 <sup>a</sup> 38	1674.731	2 <sup>+</sup>			
5502.9	(8 <sup>+</sup> )	1906.0 10	100	3596.90	6 <sup>+</sup>	[E2]		B(E2)(W.u.)>12
5523.0	0 <sup>+</sup>	4712 3	100	810.7662	2 <sup>+</sup>			
5832.08	(9 <sup>+</sup> )	1162.64 18	100	4669.38	(7 <sup>+</sup> )	E2		B(E2)(W.u.)=50 +5-19 Mult.: from $\gamma(\theta)$ in ( <sup>13</sup> C,3n $\gamma$ ) and RUL.
6032.9	(9 <sup>+</sup> )	1364.0 6	100	4669.38	(7 <sup>+</sup> )	Q		
6282.7	(9 <sup>+</sup> )	939.4 4	100	5343.33	8 <sup>+</sup>	D		Mult.: from $\gamma(\theta)$ in ( <sup>13</sup> C,3n $\gamma$ ).
7242.6	(10 <sup>+</sup> )	959.9 7	100	6282.7	(9 <sup>+</sup> )	D		Mult.: from $\gamma(\theta)$ in ( <sup>13</sup> C,3n $\gamma$ ).
7456.7	(10 <sup>+</sup> )	1424.1 4	100 7	6032.9	(9 <sup>+</sup> )	D		
7731.3	(11 <sup>+</sup> )	1625.7 5	62 5	5832.08	(9 <sup>+</sup> )	D		
		1898.3 4	100	5832.08	(9 <sup>+</sup> )	(E2)		B(E2)(W.u.)>12 Mult.: from $\gamma(\theta)$ in ( <sup>13</sup> C,3n $\gamma$ ) and RUL.
9444.8	(12 <sup>+</sup> )	1710.6 7	25 7	7731.3	(11 <sup>+</sup> )			$E_\gamma$ : poor fit, quoted energy may be a misprint. Level-energy difference=1716.7.
9939.1	(12)	1989.4 5	100 7	7456.7	(10 <sup>+</sup> )	Q		
		2207.7 7	100	7731.3	(11 <sup>+</sup> )			
9984.5	(12)	2253.1 5	100	7731.3	(11 <sup>+</sup> )	D		
(10044.31)	1 <sup>-</sup>	4521 3	6.8 17	5523.0	0 <sup>+</sup>			

Adopted Levels, Gammas (continued)

<u><math>\gamma(^{58}\text{Fe})</math> (continued)</u>							
$E_i(\text{level})$	$J_i^\pi$	$E_\gamma^\ddagger$	$I_\gamma^\ddagger$	$E_f$	$J_f^\pi$	Mult. <sup>†</sup>	Comments
(10044.31)	1 <sup>-</sup>	4626.3	5	28	2	5417.6	(1 <sup>+</sup> ,2,3 <sup>-</sup> )
		4749.6	6	25	2	5294.8	(1 <sup>+</sup> ,2,3 <sup>+</sup> )
		4823.7	6	20	2	5220.9	1,2
		5043.8	5	91	7	5000.23	1 <sup>+</sup>
		5212	3	5.1	25	4833.89	1 <sup>+</sup> ,2 <sup>+</sup>
		5493.6	6	92	8	4550.37	1 <sup>+</sup>
		5599.9	6	9.3	17	4444.3	1 <sup>+</sup>
		5691.3	6	21	2	4352.7	1 <sup>+</sup>
		5721.5	6	21	2	4322.5	1 <sup>+</sup>
		5746.7	6	20	2	4297.8	2 <sup>+</sup>
		5905.3	7	21	3	4139.24	1 <sup>+</sup>
		6028.7	6	10.2	17	4015.01	1 <sup>+</sup>
		6162.7	6	28	3	3880.1	1 <sup>+</sup>
		6413.9	7	13.6	17	3629.60	2 <sup>+</sup>
		6506.0	7	58	7	3537.97	1 <sup>+</sup>
		6960.3	7	89	9	3083.69	2 <sup>+</sup>
		7163	5	5.1	17	2876.34	2 <sup>+</sup>
		7261.7	8	97	11	2782.14	1 <sup>+</sup>
		8369.1	9	100	13	1674.731	2 <sup>+</sup>
		9232.9	10	19	3	810.7662	2 <sup>+</sup>
10043.2	12	23	4	0.0	0 <sup>+</sup>		
10075.4	(13 <sup>+</sup> )	2344.0	8	100	7731.3	(11 <sup>+</sup> )	(E2) B(E2)(W.u.)>4.3 Mult.: from $\gamma(\theta)$ in ( <sup>13</sup> C,3n $\gamma$ ) and RUL.
10353.8		2622.4	7	100	7731.3	(11 <sup>+</sup> )	
11857.0	(14 <sup>+</sup> )	2412.2	6	100	9444.8	(12 <sup>+</sup> )	
11911.0		1926.5	6	100	9984.5	(12)	
12813.3	(15 <sup>+</sup> )	2737.8	13	100	10075.4	(13 <sup>+</sup> )	

<sup>†</sup> The mult=Q and D correspond to  $\Delta J=2$  and  $\Delta J=1$ , respectively. The mult=D+Q correspond to  $\Delta J=1$ , but in some cases it may be  $\Delta J=0$ . When mult=E2, M1, M1+E2 or E1 is given, it follows from  $\Delta(J^\pi)$ . When given in square brackets, multipolarity is assumed from  $\Delta J^\pi$  in the present level scheme.

<sup>‡</sup> Values represent averages of all available data.

#  $\gamma$  looked for but not seen in <sup>58</sup>Co  $\varepsilon$  decay (1974Ti01), an upper limit of intensity is given.

@ Total theoretical internal conversion coefficients, calculated using the BrIcc code (2008Ki07) with Frozen orbital approximation based on  $\gamma$ -ray energies, assigned multipolarities, and mixing ratios, unless otherwise specified.

& Multiply placed with undivided intensity.


<sup>a</sup> Multiply placed with intensity suitably divided.

<sup>b</sup> Placement of transition in the level scheme is uncertain.

**Adopted Levels, Gammas**

Level Scheme

Intensities: Relative photon branching from each level




$^{58}_{26}\text{Fe}_{32}$



Adopted Levels, GammasLevel Scheme (continued)

Intensities: Relative photon branching from each level

 $^{58}_{26}\text{Fe}_{32}$


**Adopted Levels, Gammas**

Legend

**Level Scheme (continued)**

Intensities: Relative photon branching from each level  
 @ Multiplied: intensity suitably divided

-----▶  $\gamma$  Decay (Uncertain)



$^{58}\text{Fe}_{32}$


**Adopted Levels, Gammas**

Level Scheme (continued)

Legend

Intensities: Relative photon branching from each level  
& Multiply placed: undivided intensity given  
@ Multiply placed: intensity suitably divided

----->  $\gamma$  Decay (Uncertain)



$^{58}\text{Fe}_{32}$


**Adopted Levels, Gammas**

**Level Scheme (continued)**


**Legend**

Intensities: Relative photon branching from each level  
& Multiply placed: undivided intensity given  
@ Multiply placed: intensity suitably divided

-----▶  $\gamma$  Decay (Uncertain)



<sup>58</sup>Fe<sub>32</sub>

**Adopted Levels, Gammas** $^{58}_{26}\text{Fe}_{32}$