

Adopted Levels, Gammas

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	Caroline D. Nesaraja, Scott D. Geraeds and Balraj Singh		NDS 111,897 (2010)	12-Jan-2010

$Q(\beta^-) = -8561.0$ 5; $S(n) = 12216.3$ 5; $S(p) = 8172.2$ 5; $Q(\alpha) = -6399.2$ 4 [2012Wa38](#)

Note: Current evaluation has used the following Q record -8565.6 1412217.0 188172.4 5 -6400.0 6 [2009AuZZ](#), [2003Au03](#).
 $S(2n) = 22467$ 11, $S(2p) = 14200.2$ 6 ([2009AuZZ](#)).

Other reactions:

$^{56}\text{Fe}(^{12}\text{C}, ^{10}\text{Be})$: [1998Pa43](#): $E=60$ MeV, $\sigma(\theta)$.

Additional information 1.

$^{58}\text{Ni}(^6\text{Li}, ^6\text{Li}')$: [1994Sa33](#), [1989Na11](#); elastic and inelastic scattering. Extracted deformation parameters, optical model potential.

$^{58}\text{Ni}(^6\text{Li}, ^6\text{Li}')$: [2000Sc11](#): $E=600$ MeV; [2009Ag02](#): $E=9.9, 11.2, 12.1, 13.0, 14.0$ MeV; measured σ , $\sigma(\theta)$.

$^{58}\text{Ni}(^7\text{Li}, ^7\text{Li}')$: [2000Gu17](#), [1999Gu02](#): $E=42$ MeV, $\sigma(\theta)$.

$^{58}\text{Ni}(\text{pol } ^7\text{Li}, ^7\text{Li})$: [1995De06](#): $E=70.5$ MeV, $\sigma(\theta)$, $Ay(\theta)$.

$^{58}\text{Ni}(^7\text{Be}, ^7\text{Be})$: [2009Ag02](#): $E=15.1, 17.1, 18.5, 19.9, 21.4$ MeV; measured σ , $\sigma(\theta)$.

$^{58}\text{Ni}(^8\text{B}, ^8\text{B})$: [2009Ag02](#): $E=20.7, 23.4, 25.3, 27.2, 29.3$ MeV; measured $\sigma(\theta)$.

$^{58}\text{Ni}(^8\beta, p^7\text{Be})$: [2008Ag11](#): $E=25.0, 26.9, 28.4$ MeV; measured light fragment energy spectra, $\sigma(\theta)$, excitation functions.

$^{58}\text{Ni}(^{12}\text{C}, ^{12}\text{C})$, ($^{12}\text{C}, ^{12}\text{C}'$): [1987FeZX](#); DWBA analysis extracted deformation parameters.

$^{58}\text{Ni}(^{14}\text{N}, ^{14}\text{N}')$: [1990Ga07](#); studied relationship between centroid and Γ of giant quadrupole resonance and neutron binding energy.

$^{58}\text{Ni}(^{18}\text{O}, ^{18}\text{O})$: [1997Si13](#): $E=35.1-55.1$ MeV, $\sigma(\theta)$.

$^{58}\text{Ni}(^{28}\text{Si}, ^{28}\text{Si})$: [2003Ga18](#): $E=74-77$ MeV, $\sigma(\theta)$.

$^{58}\text{Ni}(^{46}\text{Ti}, ^{46}\text{Ti}')$, ($^{58}\text{Ni}, ^{58}\text{Ni}'$), ($^{62}\text{Ni}, ^{62}\text{Ni}'$): [2000Va28](#): $E=96.9-116.5$ MeV, $\sigma(\theta)$.

$^{58}\text{Ni}(^{58}\text{Ni}, ^{58}\text{Ni})$: [2007Hi06](#): $E=220-260$ MeV, $\sigma(\theta)$.

$^{58}\text{Ni}(^{58}\text{Ni}, ^{58}\text{Ni})$: [1996Va10](#): $E=220-240$ MeV, $\sigma(\theta)$.

$^{58}\text{Ni}(^{58}\text{Ni}, ^{58}\text{Ni})$: [1994Me24](#): $E=200$ MeV, $\sigma(\theta)$.

$^{58}\text{Ni}(\pi, \pi)$, $^{58}\text{Ni}(\pi^+, \pi^+)$, $^{58}\text{Ni}(\pi^-, \pi^-)$: [1996La04](#), [1991Ra22](#), [1989Oa01](#); strong absorption model analysis of elastic scattering to extract deformation.

^{58}Ni mesic atoms, pionic x-rays: [1990Ku08](#), interaction shifts, and widths.

$^{58}\text{Ni}(\text{d}, \text{n})$: [1993InZZ](#); deduced occupation probabilities of proton orbits.

$^{58}\text{Ni}(\alpha, \alpha)$: [1992Du08](#); analysis of total cross section data to extract mean square radii of matter distribution.

$\text{Cu}(\text{K}-\gamma)$: [1972Ba55](#).

Structure calculations (levels, transition probabilities, etc.): [2009Be24](#), [2007Sv01](#), [2006Va21](#), [2004Ho08](#), [1999Ha21](#), [1977Ko02](#), [1975Va08](#), [1974Pa13](#), [1972Gl05](#), [1972Ob02](#).

58Ni Levels

Individual values of τ in ps for first 2^+ state at 1454 keV that were used in averaging are given below:

- Deduced from B(E2) measurement in Coulomb excitation: 0.82 16 ([1960An07](#)), earlier value of 0.67 17 in [1959Al95](#)), 0.95 6 and 1.04 22 ([1960Go08](#)), 0.88 9 ([1962St02](#)), 0.860 20 ([1970Le17](#)), 0.924 28 ([1971ChZF](#)), 0.83 17 ([2004Yu10](#)).
- From Γ in (γ, γ') : 0.62 20 ([1964Bo22](#)), 0.98 9 ([1970Me18](#)), 1.07 8 ([1972ArZD](#)), 0.90 11 ([1981Ca10](#)).
- From B(E2) in (e, e') : 0.956 16 ([1967Du07](#)), 1.14 6 ([1969Af01](#)), 1.07 9 ([1983Kl09](#)). Uncertainties in B(E2) are statistical, 15% for systematic uncertainty as suggested in [1967Du07](#) is added in quadrature. Other: 0.65 9 ([1961Cr01](#)), not included in the averaging procedure.
- From DSAM in $(p, p'\gamma)$: 0.94 12 ([1969Be48](#)), 0.92 17 ([1973BeYD](#)).
- From DSAM in $(n, n'\gamma)$: 1.00 ps +15–10 ([2008Or02](#)), weighted average of two measurements at $E(n)=1.6$ and 1.8 MeV). Value of 42 fs 12 from [1989Ge09](#) is discrepant and highly suspect.
- From DSAM in Coulomb excitation: 1.27 2 ([2001Ke08](#)), uncertainty is increased to 0.07 to take into account 5% systematic uncertainty due to stopping powers, as suggested by one of the authors of [2001Ke08](#) in an e-mail reply in December 2007. It should be pointed that this value stands as the highest amongst all the others and is higher by $\approx 35\%$ from the precise values deduced from B(E2) values in Coulomb excitation. In the e-mail reply, the author of [2001Ke08](#) claimed that their measurements are reliable for two main reasons: a) the γ rays were detected in coincidence with scattered ^{12}C ions thus giving clean γ -ray spectra, b) high ion velocities in inverse kinematics used for the first time. In a thesis by [2005NiZS](#) where lifetime of first 2^+ state in ^{22}Ne was measured using Coulomb excitation technique and ^{nat}Ni and ^{107}Ag as targets, the results for first 2^+ state in ^{22}Ne were found

Adopted Levels, Gammas (continued)⁵⁸Ni Levels (continued)

to be consistent in the two measurements only when B(E2) value for first 2⁺ state from Coulomb excitation data was used. Use of the lifetime from [2001Ke08](#) gave inconsistent results.

Cross Reference (XREF) Flags

A	⁵⁸ Cu ε decay (3.204 s)	J	⁵⁴ Fe(¹² C, ⁸ Be)	S	⁵⁸ Ni(³ He, ³ He')
B	⁵⁹ Zn εp decay (182.0 ms)	K	⁵⁴ Fe(¹⁶ O, ¹² C)	T	⁵⁸ Ni(α,α')
C	²⁸ Si(³⁶ Ar, α 2p γ)	L	⁵⁶ Fe(³ He,n)	U	⁵⁸ Ni(⁶ Li, ⁶ Li')
D	²⁸ Si(³⁶ Ar, α p γ):prompt p decay	M	⁵⁶ Fe(α ,2n γ)	V	Coulomb excitation
E	⁴⁰ Ca(²⁴ Mg, α 2p γ)	N	⁵⁸ Ni(γ,γ'),(pol γ,γ')	W	⁶⁰ Ni(p,t)
F	⁴⁵ Sc(¹⁶ O,p2n γ)	O	⁵⁸ Ni(e,e')	X	Ni(K ⁻ ,x ray γ)
G	⁴⁸ Ti(¹² C,2n γ)	P	⁵⁸ Ni(n,n'),(n,n' γ)	Y	⁵⁸ Ni(¹⁶ O, ¹⁶ O')
H	⁵⁴ Fe(⁶ Li,d)	Q	⁵⁸ Ni(p,p'),(pol p,p'),(p,p' γ)	Z	⁵⁸ Ni(¹⁸ O, ¹⁸ O')
I	⁵⁴ Fe(⁷ Li,t)	R	⁵⁸ Ni(d,d')		

E(level) [†]	J ^{π‡}	T _{1/2} [#]	XREF	Comments
0.0 I	0 ⁺	stable	ABCDEFGHIJKLMNPQRSTUVWXYZ	T _{1/2} : >7.0×10 ²⁰ y for decay by double $\varepsilon\beta^+$ channel to the 0 ⁺ g.s. of ⁵⁸ Fe, and >4.0×10 ²⁰ y for decay by the same mode to the 2 ⁺ , 811-keV level of ⁵⁸ Fe (1993Va19). Others: 1984No09 , 1982Be20 .
1454.21 I 9	2 ⁺	0.652 ps 21	ABCDEFGHIJK MNOPQRST VWXYZ	$\mu=+0.076$ 18 (2001Ke02,2005St24) $Q=-0.10$ 6 (1974Le13,1989Ra17) $\langle r^2 \rangle^{1/2}=3.7748$ fm 14 (2004An14 evaluation). J ^π : E2 γ to 0 ⁺ . T _{1/2} : different averaging methods were employed to 20 independent values given in header comments above but minimum uncertainty was assigned as 5%, which required increasing the uncertainty by a factor of ≈2 to values in 1970Le17 and 1971ChZF . Average values of τ in ps obtained are: 0.95 3 by weighted average, normalized $\chi^2=2.0$; 0.95 9 by limitation of statistical weights method (the uncertainties is increased to overlap the most precise value of 0.86 4); 0.931 21 by normalized residuals method (NRM), normalized $\chi^2=1.2$; 0.926 21 by Rajeval's technique, normalized $\chi^2=1.0$. The evaluators adopt 0.94 3 as in 2008Or02 ; this value overlaps all the averaging methods used. 2001Ra27 evaluation (which did not include 2008Or02 , 2004Yu10 and 2001Ke08) lists $\tau=0.904$ ps 26. μ : transient field integral perturbed angular correlation (2001Ke02). Other: -0.12 24 (1978Ha13,1989Ra17). Q: reorientation in Coulomb excitation (1974Le13,1970Le17). See also 2005St24 compilation.
2459.21 I 14	4 ⁺	3.7 ps 4	BCDEFGH JK M OPQRST VWX	J ^π : ΔJ=2, E2 γ to 2 ⁺ ; L(α,α')=4.
2775.42 I 14	2 ⁺	0.38 ps +12-9	AB HI K NOPQR T W	T _{1/2} : from 2001Ke08 , Coulomb excitation. T _{1/2} : from DSA in (p,p' γ). T _{1/2} =57 fs +25-13 from

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued)**⁵⁸Ni Levels (continued)**

E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF						Comments			
2902.15 21	1 ⁺ ^h	69 fs +15-14	A	H	N	PQ	t		B(E2) in (e,e') is in disagreement. J ^π : L(α, α')=2.			
2942.56 18	0 ⁺	1.46 ns 14	A	H	NOPQ	St	W		Configuration= $\nu(p_{3/2}f_{5/2})$ ((p,p'), 1986Ho15). T _{1/2} : from $\gamma\gamma(t)$ in ⁵⁸ Cu decay (1970Ra34). Other: 2.01 ns 7 in (p,p' γ) (1971St02) is in disagreement. Value from decay work is preferred due to cleaner γ -ray spectra such studies than in reaction data.			
3037.86 16	2 ⁺	57 fs 8	AB	H	NOPQRST VW							
3263.66 22	2 ⁺	37 fs 5	AB	H K	NOPQRST VW							
3269.1 8	(2) ⁱ	>57 [@] fs			N							
3273.7 7	(2) ⁱ	>50 [@] fs			N							
3420.55 ^q 18	3 ⁺	0.26 ps +22-10	C		K	PQ	T	W	J ^π : L(p,p')=2+4 gives 3 ⁺ uniquely.			
3450.9 5		>11 [@] fs			N							
3524 5	4 ⁺			I	R T							
3531.1 3	0 ⁺	0.19 ps 6	A	H KL	OPQ							
3593.71 25	1,2 ⁺	33 fs 9	A		N	PQ	T		J ^π : γ to 0 ⁺ ; population in (γ, γ'). T _{1/2} : DSA in (p,p' γ). Other: 39 fs 7 from (γ, γ').			
3620.09 ^q 22	4 ⁺	0.11 ps +8-5	BCDEFGH	K M	OPQ	ST	W		J ^π : L(α, α')=4; $\Delta J=2$, E2 γ to 2 ⁺ .			
3775.0 3	3 ⁺	0.28 ps +14-7		K	PQ	T	W		J ^π : L(p,p')=2+4 gives 3 ⁺ uniquely.			
3870					R							
3898.8 4	2 ⁺	23 fs 6	AB	H K	NOPQR T W							
3943.6 12		>24 [@] fs			N							
4020				I	O Q							
4105.9 3	(4 ⁺)		C E						J ^π : $\Delta J=(2)$, Q γ to 2 ⁺ . E(level): this level is different from 4107.7, 2 ⁺ .			
4108.4 3	2 ⁺	128 fs 55		H K	NOPQRST W							
4260 80	(2 ⁺)			L								
4294.7 4	4 ⁺	24 fs +22-18	C	K	OPQ		W		E(level): broad peak, from energy matching it could correspond to 4295 level, but J^{π} 's are different. J ^π : L(³ He,n)=2.			
4347.9 12		17 fs +15-13		k	PQ	T	W		J ^π : 4 from (e,e'); γ to 2 ⁺ and RUL.			
4358.7 7	(2 ^{+,3,4⁺)}		E	k	Q				J ^π : γ 's to 2 ⁺ and 4 ⁺ .			
4383.0 ^q 3	(5 ⁺)		CDEFG	M	Q	T			J ^π : $\Delta J=1$, (M1+E2) γ to 4 ⁺ .			
4404.3 4	4 ⁺	43 fs +17-14	C		PQ	T	W		J ^π : L(α, α')=4.			
4449.6 4	1 ^{+,2⁺}		A		QR	T			J ^π : log ft=5.1 from 1 ⁺ .			

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) **^{58}Ni Levels (continued)**

E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF											Comments
			B	H	I	K	O	P	Q	R	S			
4474.6 5	3 ^{-a}	22 fs 6												T _{1/2} : weighted average of 19 fs 8 from (p,p'γ), and 24 fs 8 from (n,n'γ). B(E3)(↑)=0.0176 16 (2002Ki06 evaluation, adopted from (e,e')).
4518.3 8									Q					
4538.0 6	0 ^{+a}	31 fs 11	A					OPQ	T					
4574.1 5	1 ⁱ	21 @ fs 3			J		N	Q					XREF: N(?)	
4752.2 8	4 ⁺		B				O	QRST	W				XREF: S(4720) J ^π : L(α, α')=4. E(level): weighted average of 4750 5 (e,e'), 4750 7 (p,t), 4750 8 (α, α'), 4755 5 (p,p').	
4920.0 6								Q						
4954.0 8	1 ⁱ	14 @ fs 2					N							
4964.7 3	(5 ⁺)		C E				Q						J ^π : ΔJ=1 γ to 4 ⁺ .	
5064.3 10				I			Q							
5084.5							Q							
5127.5 ^l 4	6 ^{+ah}		CDEFGH		M	O	Q	T					J ^π : L(p,t)=2.	
5156 11	2 ⁺							W						
5166 10	1 ^h						Q						Configuration= $v(p_{3/2}p_{1/2}) ((p,p'), 1986Ho15)$.	
5170.3 10				K			Q	S						
5359.3? 16	(2) ⁱ	>29 @ fs					N							
5384.5 ^q 4	6 ⁺		CDE H				Q						J ^π : ΔJ=1, E1 γ from 7 ⁻ ; ΔJ=2, Q γ's to 4 ⁺ .	
5394.0 9		41 @ fs 8		H			N	T						
5436.3 10	4 ⁺						O	Q					J ^π : L(p,p')=4.	
5452.2 4	1 ⁱ	>13 @ fs					N	Q						
5472.3 8	4 ⁺						Q		W				J ^π : L(p,p')=4 for 5470 5, and L(p,t)=4 for 5488 11.	
5503.5 10							Q							
5528.0 4	(1) ⁱ	>7 @ fs					N							
5589.0 7	(5 ⁻)		CD	H				t					J ^π : L(α, α')=4+5 for a doublet; γ from 7 ⁻ ; L(⁶ Li,d)=(5,6).	
5590.3 10	2 ⁺						Q	ST	W				J ^π : L(α, α')=2.	
5594.2 6	4 ^{+a}			K	O	Q	t						XREF: O(5585)	
5706.3 8							Q							
5744.7 5	(6 ⁺)		C										J ^π : ΔJ=2, Q γ to 4 ⁺ .	
5748.5 8	2 ⁺						Q						J ^π : L(p,p')=2.	
5766.3 8	4 ⁺						Q						J ^π : L(p,p')=4.	
5803.3 7							Q							
5824.6 11							Q							
5896.4 7			H		o	Q	S							
5905.3 7	1 ⁺ⁱ	25 @ fs 4			No								E(level): in (e,e'), level is at 5909 8 with J=2 in one study and 5903 15 in another with J=1, with assumed natural-parity states. It is possible that it is doublet in (e,e') representing 1 ⁺ and 2 ⁺ levels.	
5906 5	2 ⁺				o	Q							J ^π : L(p,p')=2. E(level): see comment for 5905.3 level.	
5924 10	(0 ⁺) ^b				o	Q		W						
5942.4 10	(0 ⁺) ^b				o	Q		W						
5963 10	(0 ⁺) ^b					Q		W						
5967 8	2 ^{+,3-} ^a					O								
5982 10	(0 ⁺) ^b		I		Q		W							
6018.4 10	3 ^{-a}				O	Q								

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) **^{58}Ni Levels (continued)**

E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF	Comments
6027.3 7	1 ⁻	0.85@ fs 5	H k NO Q T	J ^π : L(α, α')=1; J=1 from (γ, γ'); J ^π =2 ⁺ favored in (e,e'), but 1 ⁻ is not ruled out.
6067.5 ^q 4	(7 ⁺)		CDEF	J ^π : ΔJ=2, Q γ to (5 ⁺); ΔJ=1, (M1+E2) γ to 6 ⁺ .
6084.7 5	7 ⁻		CDE	J ^π : ΔJ=1, E1 γ to 6 ⁺ ; γ to 4 ⁺ not ΔJ=1, D+Q.
6116 10				
6145 15	3[−] ^a		O	
6174.3 8	2 ^{+,3⁻^a}		O Q	XREF: O(6182)
6199 10			H	
6220 10			Q	
6220.0 4	(7 ⁺)		C E	J ^π : ΔJ=1 γ to 6 ⁺ ; ΔJ=(2) γ to (5 ⁺).
6228.3 6	(2 ⁺) ^a		O Q	J ^π : 2 ^{+,1⁻ in (e,e') for a 6235 8 level.}
6248 10	(2 ⁺) ^a		O Q	J ^π : 2 ^{+,1⁻ in (e,e') for a 6235 8 level.}
6271 10			Q	
6274.3 10	4[+] ^a		O Q	
6308.5 6	3 ⁻		Q T	J ^π : L(p,p')=3.
6316 10	1 ⁻ ,2 ⁺ ^a		O Q	
6360.6 11			Q	
6389 10			W	
6402.4 10			i	
6417 5	2 ⁺		i O Q	J ^π : L(p,p')=2, also (e,e').
6424.9? 9	1 ⁱ	9.3@ fs 13	N	
6430.7? 10	1 ⁱ	6.9@ fs 7	N	
6437 10			hi Q S	
6447 10			hi K Q	E(level): there is a level at 6450 50 in (¹⁶ O, ¹² C).
6460 5	4 ⁺		Q T	J ^π : L(p,p')=4.
6468.4 7	(1 ⁺) ^a		O Q	J ^π : 1 ^{+,2⁻ from (e,e').}
6478.4 7	2 ⁺		Q	J ^π : L(p,p')=2.
6500 10			Q	
6507.2 11			Q	
6549 10	(4 ⁺) ^c		Q	
6571.4 10	2 ⁺		Q	J ^π : L(p,p')=2.
6598 10	(4 ⁺) ^c		F Q	
6601.3 8			W	
6604.6 ^l 4	(8 ⁺)		CDEFG M	XREF: G(?)M(?) J ^π : ΔJ=2, Q γ to 6 ⁺ ; ΔJ=1, (M1+E2) γ to 7 ⁺ ; band assignment.
6665.4 7			Q	
6674 10			Q	
6685.0? 9	1 ⁱ	3.6@ fs 4	N	
6714 10			Q W	
6717.4 7			Q W	
6735 8	3 ⁻		O T W	J ^π : L(α, α')=3; also (e,e').
6752 5	2 ⁺		Q	J ^π : L(p,p')=2.
6763.5 10	3 ^{-f}		hi O Q s	E(level): level at 6780 in (³ He, ³ He') where ΔE=25 keV for strong and 50 keV for weak levels.
6793 10	3 ^{-f}		iJk Q s	
6805.5 10	3 ^{-f}		hiJk Q s	E(level): level at 6780 30 in (⁷ Li,t), level at 6800 50 in (¹⁶ O, ¹² C).
6816 8	(2 ⁺) ^a		O Q	J ^π : 2 ^{+,1⁻ in (e,e').}
6845.7 7	(7 ⁺)			J ^π : γ 's to (5 ⁺) and 6 ⁺ .
6854.5 10	3 ^{-a}		H O Q T W	XREF: Q(6844) E(level): level at 6860 40 in (p,t).

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) **^{58}Ni Levels (continued)**

E(level) [†]	J [‡]	T _{1/2} [#]	XREF	Comments
6863.1 6	(6)		C	J ^π : ΔJ=1, D γ to (5^+). Negative parity assigned by 2009Jo03 in ($^{36}\text{Ar},\alpha 2\text{p}\gamma$).
6886 10	($2^+, 3^-$) ^d		1 Q	
6892.9? 15	(1) ⁱ	11 @ fs 5	N	
6912 10	($2^+, 3^-$) ^d		1 Q	
6925 10	4 ^{+e}		o Q	
6935 10	4 ^{+e}		o Q	
6960 10			Q	
6983 5	2 ⁺		0 Q	J ^π : L(p,p')=2; also (e,e').
6992.5 10			Q	
7017 10			i Q	
7042 10			i o Q	
7048.2 9	1 ⁻ⁱ	0.83 @ fs 3	i No	
7051 5	4 ⁺		i o Q	J ^π : L(p,p')=4.
7054.5 10			i o Q T	
7068 5	4 ⁺		Q	J ^π : L(p,p')=4.
7089 10			o Q	
7109 8	(2^+) ^a		i 0	J ^π : 4 ⁽⁺⁾ is also suggested in (e,e').
7111 5	3 ⁻		i Q	J ^π : L(p,p')=3.
7113.5& 7	(1,2 ⁺)		i Q	
7131.5 10			i Q	
7141 5	4 ⁺		i Q	J ^π : L(p,p')=4.
7180 25	3 ⁻		S w	J ^π : L($^3\text{He}, ^3\text{He}'$)=3.
7210.4 10	3 ⁻		H K 0 Q T w	J ^π : L(p,p')=3; also (e,e'); L(α, α')=4 is inconsistent.
7249.6 11	(1) ⁱ	9 @ fs 3	N	J ^π : L(p,p')=2; also (e,e').
7255 5	2 ⁺		0 Q	
7271.7 7	1 ⁱ	0.99 @ fs 11	N Q	J ^π : possible negative parity since no analog > state seen in ^{58}Cu from ($^3\text{He}, t$) (2002Fu07).
7273.7 6	7 ⁻		C	J ^π : ΔJ=1, E1+M2 γ to 6 ⁺ ; ΔJ=0, D+Q γ to 7 ⁻ .
7300.5 10	3 ⁻		0 Q	J ^π : L(p,p')=3.
7314.8 ^g 5	(8^+)		C	J ^π : ΔJ=1, D+Q γ to (7 ⁺); ΔJ=2, Q γ to 6 ⁺ .
7380.5 10	(1,2 ⁺)		Q	J ^π : γ to 0 ⁺ .
7388.8 4	1 ⁺ⁱ	1.00 @ fs 5	NO	
7420 5	3 ⁻		Q	J ^π : L(p,p')=3.
7446.2 ^q 5	(9^+)		C EF	J ^π : ΔJ=1, (M1+E2) γ to (8 ⁺); ΔJ=2, Q γ to (7 ⁺).
7462 8	(1 ⁺) ^a		0 Q	J ^π : 1 ⁺ ,(2 ⁻) in (e,e').
7514.5 10	3 ^{-a}		hi l 0 Q ST	
7560 8	1 ^{+a}		Hi kl 0	
7570.5 10	2 ⁺		kl Q	J ^π : L(p,p')=2.
7585.1 6			N	
7595.9 6	(2) ⁱ	5.2 @ fs 8	N	
7603 8	(1 ⁻) ^a		0	
7616.0? 10	(1) ⁱ	9.5 @ fs 40	N	
7618 5	4 ⁺		Q	
7680.6 10	1 ^{-a}		0 Q	J ^π : L(p,p')=4.
7709.7 6	1 ⁺ⁱ	0.72 @ fs 3	J NO Q	Configuration= $\nu(f_{7/2}^{-1}f_{5/2})$ ((p,p'), 1986Ho15).
7721 10			I Q T	
7724.3 ^r 5	(8^+)		C	J ^π : ΔJ=2 γ to 6 ⁺ ; ΔJ=0, D+Q γ to (8 ⁺).
7748 8	(1 ^{+,2-}) ^a		0 Q	
7766.0 7	(1) ⁱ	3.7 @ fs 6	N	
7807.3 5	1 ⁻ⁱ	0.81 @ fs 10	K N Q	

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) **^{58}Ni Levels (continued)**

E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF	Comments
7820 8	4[+] ^a		H O	
7858 5	3 ⁻		Q	
7860 5	4 ⁺		Q	J ^π : L(p,p')=3.
7862.6 ^{&} 7	(1,2 ⁺)		Q	J ^π : L(p,p')=4.
7876.7 26	1 ⁱ	0.9@ fs 5	N	J ^π : γ to 0 ⁺ .
7937 25	8 ^{-a}		H O	
7973.6 6	(8 ⁺)		C	J ^π : ΔJ=2 γ to (6 ⁺); ΔJ=1, D+Q γ to 7 ⁺ .
7982.8 6	(8 ⁻)		C	J ^π : ΔJ=1, D+Q γ to (7 ⁺); ΔJ=1, (M1+E2) γ to 7 ⁻ .
8068.6? 12	(1 ⁻ⁱ)	1.38@ fs 17	N	
8074.5 7	(8 ⁺)		C	J ^π : ΔJ=1, D+Q γ to (7 ⁺).
8096.3 6	1 ⁱ	1.6@ fs 3	N	
8100 15	4[+] ^a		K O T	XREF: K(8060)
8110.6 10	(1,2 ⁺)		Q	J ^π : γ to 0 ⁺ .
8115.1 6	(8 ⁻)		C	J ^π : γ to 7 ⁻ .
8120.8 ^r 5	(9 ⁺)		C E	J ^π : ΔJ=1, D+Q γ's to (8 ⁺).
8134 5	3 ⁻		Q	J ^π : L(p,p')=3.
8143 10			Q	
8203 20	(1 ⁺) ^g		K Q	
8237.3 4	1 ⁻ⁱ	0.15@ fs +3-2	NO Q	J ^π : 1 ⁺ suggested in (e,e') is in disagreement, if the level is the same as in (γ,γ'). J ^π : 1 ^{+,} (2 ⁻) in (e,e').
8276 8	1 ^{+ag}		O Q	
8317.1 17	1 ⁱ	1.9@ fs 3	N Q	
8372 20	(1 ⁺) ^g		Q	
8395 8	2 ^{+a}		O	
8395.1 12	1 ⁻ⁱ	0.40@ fs 8	N Q	
8419 10	1 ^{+gh}		Q	Configuration=γ(f _{7/2} ⁻¹ f _{5/2}) ((p,p'), 1986Ho15).
8461.0 7	1 ^{+gi}	0.51@ fs 3	N Q	
8475 8	2 ^{-a}		O	
8493 15	(3 ⁻ ,1 ⁻)		T	J ^π : L=(3,1) in (α,α').
8514.1 4	1 ⁻ⁱ	0.66@ fs 5	NO Q	J ^π : 1 ⁺ suggested in (e,e') is in disagreement, if the level is the same as in (γ,γ').
8552.7 13	1 ⁽⁺⁾	0.97@ fs 8	N Q	J ^π : from analysis of σ(θ) in (p,p'), J=1 from (γ,γ'). Configuration=γ(p _{3/2} p _{1/2}) ((p,p'), 1986Ho15).
8600.5 7	1 ^{+agi}	0.57@ fs 6	NO Q	
8654 9	(3 ⁻ ,1 ⁻)		Q T	E(level): unweighted average of 8645 10 (p,p'), and 8662 15 (α,α'). J ^π : L=(3,1) in (α,α').
8679.3 8	1 ^{+agi}	0.223@ fs 11	NO Q	
8692			Q	
8716 10			Q	
8718.1 6	(9 ⁻)		C	J ^π : ΔJ=1, (M1+E2) γ to (8 ⁻); γ to 7 ⁻ .
8780 8	2 ^{-a}		O	
8797 5	3 ⁻		Q	J ^π : L(p,p')=3.
8808 25	8 ^{-a}		O	
8817 8	(1 ⁺) ^a		O	J ^π : 1 ^{+,} (2 ⁻) in (e,e').
8830 40	2 ⁺		W	J ^π : L(p,t)=2.
8845 5	3 ⁻		O Q	J ^π : L(p,p')=3; 2 ^{+,} 3 ⁻ from (e,e').
8857.4 6	1 ^{(+)gi}	0.61@ fs 12	N Q	
8880.2 6	1 ⁻ⁱ	0.390@ fs 17	N Q	
8896.4? 10			C	J ^π : γ to (8 ⁺) suggests (8,9,10 ⁺).

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) **^{58}Ni Levels (continued)**

E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF		Comments
			J	Q	
8902 5	4 ⁺				J ^π : L(p,p')=4.
8934.6 5	1 ⁽⁻⁾ <i>i</i>	0.310@ fs 11	NO		J ^π : parity from (e,e').
8961.3 7	1 ⁺ <i>agi</i>	1.20@ fs 13	NO Q		
9012 5	3 ⁻		Q		J ^π : L(p,p')=3.
9027.2 7	(9 ⁻)		C		J ^π : ΔJ=2 γ to 7 ⁻ ; γ to (8 ⁻).
9037 8	(1 ⁺) <i>a</i>		O		J ^π : 1 ⁺ ,(2 ⁻) in (e,e').
9062.7 ^r 6	(10 ⁺)		C		J ^π : ΔJ=2 γ to (8 ⁺); ΔJ=1, (M1+E2) γ to (9 ⁺).
9073.4 6	1 ⁺ <i>agi</i>	0.51@ fs 3	NO Q		
9113 10			O		
9156.9 7	1 ⁺ <i>agi</i>	0.77@ fs 10	NO Q		
9190.7 5	1 ⁻ <i>i</i>	0.58@ fs 6	N Q		J ^π : (1 ⁺) suggested in (p,p') is in disagreement, if the level is the same as in (γ,γ').
9251 10	(1 ⁺) <i>g</i>		O Q		
9295 10	1 ⁺ <i>h</i>		O Q T		Configuration=γ(f _{7/2} ⁻¹ f _{5/2}) ((p,p'), 1986Ho15).
9304 5	3 ⁻		O Q		J ^π : L(p,p')=3.
9310 40	4 ⁺			W	J ^π : L(p,t)=4.
9322.1 ^q 9	(11 ⁺)		C		J ^π : ΔJ=2 γ to (9 ⁺).
9326.4 6	1 <i>i</i>	0.33@ fs 5	N Q		
9336 20	(1 ⁺) <i>g</i>		Q		
9345.5 6	(10 ⁻)		C E		XREF: E(?)
9368.5 6	1 ⁽⁺⁾ <i>ai</i>	0.37@ fs 4	NO		J ^π : ΔJ=2 γ to (8 ⁻); ΔJ=1, (M1+E2) γ to (9 ⁻).
9379 5	3 ⁻		Q		J ^π : 1 ⁺ ,(2 ⁻) in (e,e'); 1 ⁽⁻⁾ in (γ,γ').
9407 10	(2 ⁻) <i>a</i>		O		J ^π : L(p,p')=3.
9436 5	4 ⁺		Q		J ^π : 2 ⁻ ,(1 ⁺) in (e,e').
9455.4 18	1 <i>i</i>	2.1@ fs 4	N		J ^π : L(p,p')=4.
9458 5	3 ⁻		O Q		J ^π : L(p,p')=3.
9523.3 13	1 ⁻ <i>i</i>	0.118@ fs 13	NO Q		J ^π : 1 ⁺ ,(2 ⁻) suggested in (e,e') and (1 ⁺) in (p,p') are in disagreement, if the levels in these two reactions are the same as in (γ,γ').
9554.0 21	1 <i>i</i>	0.335@ fs 20	K NO		XREF: K(9500)
9585.2 ^o 8	(9 ⁻)		C		J ^π : (2 ⁻) in (e,e').
9588 5	4 ⁺			Q	J ^π : ΔJ=2 γ to 7 ⁻ ; ΔJ=1, D γ to (8 ⁺).
9630.5 24	1 <i>i</i>	0.15@ fs 3	N		J ^π : L(p,p')=4.
9632 5	4 ⁺		Q		J ^π : 2 ⁻ ,(1 ⁺) in (e,e').
9643 10	(2 ⁻) <i>a</i>		O		J ^π : ΔJ=2 γ to (8 ⁺); ΔJ=1, D+Q γ to (9 ⁺).
9666.9 8	(10 ⁺)		C		
9667 10	2 ⁻ <i>a</i>		O		
9667.8 15	1 <i>i</i>	0.38@ fs 13	N		
9672 5	3 ⁻		Q		J ^π : L(p,p')=3.
9723.0 9	1 ⁽⁻⁾ <i>i</i>	0.109@ fs 16	N		
9750 10	1 ⁺ <i>ag</i>		O Q		J ^π : 1 ⁺ ,(2 ⁻) in (e,e').
9790.6 10	(10 ⁺)		C		J ^π : ΔJ=1, D+Q γ to (9 ⁺).
9799 10			O		
9835 5	3 ⁻		Q		E(level): doublet in (p,p'), 3 ⁻ and 1 ⁺ levels, the latter corresponds to 9842 level here. J ^π : L(p,p')=3.
9843 5	1 ⁺ <i>gh</i>	0.26@ fs +27-10	NO Q		E(level): possible IAS of 1050,1 ⁺ in ^{58}Co . T _{1/2} : from (γ,γ') for J=1. Configuration=γ(p _{3/2} p _{1/2}) ((p,p'), 1986Ho15).

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) **^{58}Ni Levels (continued)**

E(level) [†]	J [‡]	XREF	Comments
9870 5	3 ⁻	O Q	
9886.8 7	(10 ⁺)	C	J ^π : L(p,p')=3. J ^π =(2 ⁻) in (e,e').
9890 40	2 ⁺	K	J ^π : ΔJ=2 γ to (8 ⁺). J ^π : L(p,t)=2.
9929 5	3 ⁻	1 Q	J ^π : L(p,p')=3.
9941 10	(2 ⁺) ^a	0	J ^π : 2 ^{+,} (1 ⁻) in (e,e').
9956 5	3 ⁻	1 Q	J ^π : L(p,p')=3.
10029 5	3 ⁻	0 Q	J ^π : L(p,p')=3. J ^π =(2 ⁻) in (e,e').
10059 5	4 ⁺	Q	J ^π : L(p,p')=4.
10073 10	1 ^{+a}	0	
10107 10	1 ^{+ag}	0 Q	
10120 5	4 ⁺	K Q	XREF: K(10100) J ^π : L(p,p')=4.
10137.2 12	(10 ⁺)	C	
10144.7 6	(10 ⁻)	C	
10157 10	1 ^{+eg}	O Q	E(level): possible IAS of 1377,1 ⁺ in ^{58}Co .
10180.8 6	(11 ⁻)	C	
10190 25	8 ^{-a}	0	
10192.5 ^r 7	(11 ⁺)	C	
10209 5	3 ⁻	Q	J ^π : L(p,p')=3. E(level): note that 10209 and 10214 are two different levels in (p,p').
10214 10	1 ^{+a}	O Q	J ^π : also from 0° data in (p,p') (2007Fu04) in (p,p'). E(level): possible IAS of 1435, 1 ⁺ in ^{58}Co . J ^π : L(p,p')=4.
10249 5	4 ⁺	Q	
10266 10	1 ^{+a}	0	
10293.5 11	(9 ⁻)	C	
10304 10		Q	
10355 10	1 ^{+a}	0	
10365 5	4 ⁺	Q	J ^π : L(p,p')=4.
10385 10	(1 ⁺) ^a	0	J ^π : 1 ^{+,} (2 ⁻) in (e,e').
10394.1 13	(10 ⁺)	C	
10404.8 7	(9 ⁻)	C	
10434 10	(2 ⁺) ^a	O Q	J ^π : 2 ^{+,} (1 ⁻) in (e,e').
10460 5	4 ⁺	Q	J ^π : L(p,p')=4.
10510 10	1 ^{+ag}	O Q	XREF: Q(10492) E(level): possible IAS of 1729,1 ⁺ in ^{58}Co . J ^π : L(p,p')=4.
10523 5	4 ⁺	Q	
10550 10	(1 ⁺ ,2 ⁻) ^a	O	
10582 10	(1 ⁺) ^a	O	J ^π : 1 ^{+,} (2 ⁻) in (e,e').
10586 5	3 ⁻	Q	J ^π : L(p,p')=3.
10590 50	0 ⁺	L	J ^π : L($^3\text{He},n$)=0.
10590.9 6	(11 ⁻)	C	
10630 40	4 ⁺	W	J ^π : L(p,t)=4.
10633 10	1 ^{+a}	O	
10638 5	3 ⁻	Q	J ^π : L(p,p')=3.
10667 10	1 ^{+agh}	O Q	E(level): possible IAS of 1868,1 ⁺ in ^{58}Co . Configuration= $\nu(f_{7/2}^{-1}f_{5/2})$ ((p,p'), 1986Ho15). J ^π : L(p,p')=4.
10694.7 ^p 7	(10 ⁻)	C	
10720 10	(3 ⁻ ,4 ⁺) ^a	O	
10744 5	4 ⁺	O Q	J ^π : L(p,p')=4.
10781.7 9	(11 ⁺)	C	
10805 10	1 ^{+,} 2 ⁻ ^a	O Q	
10823 5	4 ⁺	Q	J ^π : L(p,p')=4.
10856 10	(1 ⁻ ,2 ⁺) ^a	O	

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) **^{58}Ni Levels (continued)**

E(level) [†]	J [‡]	XREF			Comments
10882.0 14	(11 ⁺)	C			
10891 10	2 ^{+a}		O		
10902 5	4 ⁺		Q		J ^π : L(p,p')=4.
10950 10	1 ^{+a}		O		XREF: K(10950)
10967 5	4 ⁺	K	Q		J ^π : L(p,p')=4.
11005.6 8	(11 ⁻)	C			
11008 10	1 ^{+ag}		O Q		E(level): possible IAS of 2249,1 ⁺ in ^{58}Co .
11052 10	(1 ⁺) ^g		O Q		XREF: Q(11063)
					J ^π : (2 ⁺) in (e,e').
11080 10	(1 ⁺) ^a		O		J ^π : 1 ^{+,} (2 ⁻) in (e,e').
11117.0 ^o 8	(11 ⁻)	C			
11135 10	(3 ⁻ ,4 ^{+)a}		O		
11158 5	3 ⁻		O Q		E(level): note that 11158 and 11165 are two different levels in (p,p').
11165 10	1 ⁺		Q		J ^π : L(p,p')=3; 2 ^{+,} 3 ⁻ in (e,e').
11203 5	4 ⁺		Q		J ^π : from 0° data in (p,p') (2007Fu04) in (p,p').
11240 25	8 ^{-a}		O		J ^π : L(p,p')=4.
11255.2 ^p 7	(11 ⁻)	C			
11266 10	(1 ⁺) ^a		O Q		J ^π : 1 ^{+,} (2 ⁻) in (e,e').
11297 10	2 ^{+a}		O		
11297.7 7	(12 ⁻)	C			
11300 5	4 ⁺		Q		J ^π : L(p,p')=4.
11335 10	1 ⁻ ,2 ^{+a}		O Q		
11363 10	(2 ⁻) ^a		O		J ^π : 2 ⁻ ,(1 ⁺) in (e,e').
11410 10	(2 ^{+,} 3 ⁻) ^a		O		
11413.1 9	(11 ⁺)	C			
11434 5	4 ⁺		Q		J ^π : L(p,p')=4.
11450 25	(6 ^{+)a}		O		
11470 10	(2 ⁻) ^a		O		J ^π : 2 ⁻ ,(1 ⁺) in (e,e').
11474.5 ^r 7	(12 ⁺)	C			
11497 10	(3 ⁻)		Q	W	J ^π : L(p,t)=(3).
11536 10	(2 ⁻) ^a		O		J ^π : 2 ⁻ ,(1 ⁺) in (e,e').
11579.3 8	(12 ⁺)	C			
11593 10	2 ^{+a}		O Q		
11639 10	2 ^{+,} 3 ^{-a}		O		
11678 10	1 ^{+ag}		Q		
11728 5	4 ⁺		Q		J ^π : L(p,p')=4.
11734 10	2 ^{+a}		O		
11792 10	(2 ^{+)a}		O Q		
11814.3 8	(12 ⁻)	C			
11824.7 11	(12 ⁺)	C			
11850 40	(3 ⁻)		W		J ^π : L(p,t)=(3).
11860 10	1 ^{+a}		O		
11887 10	1 ^{+a}		O Q		J ^π : 1 ⁺ in (p,p') (2007Fu04); 2 ⁻ ,(1 ⁺) in (e,e').
11933 10	(3 ⁻ ,4 ^{+)a}		O		
11990 10	(1 ⁺) ^a		O		J ^π : 1 ^{+,} (2 ⁻) in (e,e').
11996.4 ^p 7	(12 ⁻)	C			
12040 10	2 ^{+a}		O		
12090 10			O		
12141 10	1 ⁻ ,2 ^{+a}		O		
12155.1 10	(12 ⁻)	C			
12197 10	(1 ^{+,} 2 ^{+)g}		O Q		J ^π : (2 ⁺) in (e,e').

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) **^{58}Ni Levels (continued)**

E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF	Comments
12249 10			0	
12283 10	(1) ^g		0 Q	
12330 10	(2 ⁻) ^a		0	J ^π : (1 ⁻) in (e,e').
12356.8 9	(12 ⁻)	C		J ^π : 2 ⁻ ,(1 ⁺) in (e,e').
12364.6 7	(12 ⁺)	C		
12386 10	(1 ⁺) ^g		0 Q	J ^π : (2 ⁺) in (e,e').
12447 10	(2 ⁺) ^a		0	
12482 10	(2 ^{+,4⁺}) ^a		0	
12500 25	8 ⁻ ^a		0	
12570.1 7	(12 ⁺)	C		%p=3.7 14 (2009Jo03) E(p)(c.m.)=1.83 MeV 5 (2009Jo03). Prompt p decay populates 2524, 13/2 ⁻ level in ^{57}Co which deexcites through 834-466-1224 cascade to ^{57}Co g.s.
12573 10	2 ^{+,3⁻^a}		0	
12613 10	2 ^{+a}		0	
12643 10	(1 ^{+,2⁺}) ^g		0 Q	J ^π : 2 ^{+,4⁺} in (e,e').
12700 10	(2 ⁻) ^a		0	J ^π : 2 ⁻ ,(1 ⁺) in (e,e').
12719.2 7	(12 ⁺)	C		
12744 10	(1 ^{+,2⁺}) ^g		0 Q	J ^π : (2 ⁺) in (e,e').
12796 10	(1 ⁺) ^a		0	J ^π : 1 ^{+,2⁻} in (e,e').
12831.6 ^o 9	(13 ⁻)	C		
12837 10	(2 ⁺) ^a		0	
12858 10	2 ^{+a}		0	
12912.1 ^p 9	(13 ⁻)	C		
12928 4		C		J ^π : possible γ to (11 ⁺) suggests (11,12,13).
12931 10	2 ^{+,3⁻^a}		0	
12971 10	2 ^{+a}		0	
13016.6 10	(13 ⁻)	C		
13022 10	2 ^{+,4⁺^a}		0	
13048.2 10	(13 ⁻)	C		
13057 10	2 ^{+a}		0	
13095.1 19	(12 ⁺)	C		
13125 10			0	
13129.2 18	(12 ⁺)	C		
13176 10	(1 ⁺) ^a		0	J ^π : 1 ^{+,2⁻} in (e,e').
13233 10	2 ^{+a}		0	
13238.1 7	(13 ⁺)	C		
13260 10	2 ^{+a}		0	
13305 10	(1 ^{+,2⁺}) ^g		0 Q	J ^π : (2 ⁺) in (e,e').
13345 10	2 ^{+a}		0	
13356.6 ^r 9	(13 ⁺)	C		
13411 10	1 ^{+a}		0	
13448 10	2 ^{+a}		0	
13492 10			0	
13556 10	(2 ⁺) ^a		0	
13590 10	(1 ^{+,2⁻}) ^a		0	J ^π : 2 ^{+,1⁻} in (e,e').
13606.8 ^w 13	(12 ⁺)	C		
13632 4		C		J ^π : γ to (11 ⁺) suggests (11,12,13).
13649 10	2 ^{+a}		0	
13685 10	(2 ⁺) ^a		0	
13.7×10 ³ 3	4.7 MeV 3		0	E(level): GQR.

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) **^{58}Ni Levels (continued)**

E(level) [†]	J ^π [‡]	XREF	Comments
13716 10	1 ⁺ <i>a</i>	0	
13765 10	(1 ⁺) <i>a</i>	0	
13814 10	2 ⁺ <i>a</i>	0	
13850.1 <i>p</i> 10	(14 ⁻)	C	
13884.2 17	(13 ⁺)	C	
13902 10	(2 ⁺)	0	J ^π : 2 ⁺ ,(3 ⁻) in (e,e').
13929 10	(2 ⁺) <i>a</i>	0	
13943 3		C	J ^π : γ to (11 ⁺) suggests (11,12,13).
13955 10	(2 ⁺) <i>a</i>	0	
14000 10	2 ⁺ <i>a</i>	0	
14045 10	(2 ⁺) <i>a</i>	0	
14081 10	1 ⁺ <i>a</i>	0	
14127.8 8	(14 ⁺)	C	
14138 10		0	
14180 10	(1 ⁺) <i>a</i>	0	J ^π : 1 ⁺ ,(2 ⁻) in (e,e').
14213 10	(2 ⁺) <i>a</i>	0	
14217.5 13	(14 ⁻)	C	
14272 10	1 ⁻ ,2 ⁺ ,3 ⁻ <i>a</i>	0	
14303 10	1 ⁻ ,2 ⁺ ,3 ⁻ <i>a</i>	0	
14337 10	2 ⁺ <i>a</i>	0	
14383 10	2 ⁺ <i>a</i>	0	
14441 10	(2 ⁺) <i>a</i>	0	J ^π : 2 ⁺ ,(3 ⁻) in (e,e').
14455.8 <i>n</i> 16	(13 ⁺)	C	
14470 40	(0 ⁺)		W J ^π : L(p,t)=(0).
14504 10	2 ⁺ <i>a</i>	0	
14542 10	(2 ⁺) <i>a</i>	0	J ^π : 2 ⁺ ,(1 ⁻ ,3 ⁻) in (e,e').
14598 10		0	
14630 10	2 ⁺ ,3 ⁻ <i>a</i>	0	
14692 10		0	
14736 10	(2 ⁺) <i>a</i>	0	
14823 10	2 ⁺ <i>a</i>	0	
14852 10	(1 ⁺) <i>a</i>	0	J ^π : 1 ⁺ ,(2 ⁻) in (e,e').
14853.1 <i>o</i> 11	(15 ⁻)	C	
14894 10	1 ⁻ ,2 ⁺ <i>a</i>	0	
14920.9 <i>r</i> 11	(14 ⁺)	C	
14934.7 <i>p</i> 12	(15 ⁻)	C	
14940 10	(2 ⁺) <i>a</i>	0	
15010.6 8	(14 ⁺)	C	
15031.0 10	(14 ⁺)	C	
15105.2 19		C	J ^π : γ 's to (12 ⁺) and (13 ⁺) suggest (12,13,14).
15187.0 23	(13 ⁺)	C	%p=43 6 (2009Jo03)
15241.9 14	(13 ⁻)	C	E(p)(c.m.)=2.15 MeV 5 (2009Jo03).
15242.0 18		C	Prompt p decay populates 4814, 17/2 ⁻ level in ^{57}Co which deexcites through 2290-834-466-1224 cascade to ^{57}Co g.s.
15266.3 10	(14 ⁺)	C	J ^π : γ to (13 ⁻) suggests (13,14,15).
15294.3 <i>w</i> 10	(14 ⁺)	C	
15324.1 <i>m</i> 12	(14 ⁺)	C	
≈15400	(13 ⁻)	C	%p=?
			E(p)(c.m.)≈2.35 MeV (2009Jo03).
			Prompt p decay populates 4814, 17/2 ⁻ level in ^{57}Co which deexcites through 2290-834-466-1224 cascade to ^{57}Co g.s.

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{58}Ni Levels (continued)

E(level) [†]	J [‡]	T _{1/2} [#]	XREF	Comments
15412.6 14	(13 ⁻)	C		J ^π : from 2005Ru06 .
15434.1 14	(13 ⁻)	C		
15709.3 9	(15 ⁺)	C		
15736.9 8	(15 ⁺)	C		
15858.2 9	(15 ⁺)	C		
16167.2 20		C		J ^π : γ's to (13 ⁺) and (14 ⁺) suggest (13,14,15).
16171.0 ⁿ 13	(15 ⁺)	C		
16246.6 ^p 14	(16 ⁻)	C		
16496.6 23	(16 ⁻)	C		
16567.0 9	(16 ⁺)	C		
16.64×10 ³ 12		5.81 MeV +16-11	T	E(level): L=2, isoscalar giant-quadrupole resonance (ISGQR).
16673 3	(14 ⁻)	C		%p=?
16676.4 8	(16 ⁺)	C		J ^π : from 2005Ru06 , decays by protons to
16707 3	(14 ⁻)	C		5918, 19/2 ⁻ level in ^{57}Co ; the decay mode not shown in 2009Jo03 .
16745 3	(14 ⁻)	C		%p=?
16758 3	(14 ⁻)	C		E(p)(c.m.)=2.61 MeV 12 (2009Jo03). Prompt p decay populates 5918, 19/2 ⁻ level in ^{57}Co which deexcites through 1104-2290-834-466-1224 cascade to ^{57}Co g.s.
16798.0 ^v 10	(15 ⁻)	17 ps 11	C	%p=41 6 (2009Jo03). E(p)(c.m.)=2.59 MeV 8 (2009Jo03). Prompt p decay populates 5918, 19/2 ⁻ level in ^{57}Co which deexcites through 1104-2290-834-466-1224 cascade to ^{57}Co g.s. %p=7 2 (2009Jo03); %α=2.6 3 (2009Jo03). T _{1/2} : from estimated T _{1/2} =7-28 ps (2001Ru03) from average Q(transition) in the band=2.4 3, assuming that 1364γ and 1385γ are part of the continuation of the band and that Q(transition) does not change at lower spins. E(p)(c.m.)=1.62 MeV 6, E(α)(c.m.)=6.90 MeV 6 (2009Jo03). Prompt p decay populates 6976, 21/2 ⁻ level in ^{57}Co which deexcites through 1058-1104-2290-834-466-1224 cascade to ^{57}Co g.s. Prompt α decay populates 2949, 6 ⁺ level in ^{54}Fe which deexcites through 411(6 ⁺ to 4 ⁺⁾ -1130(4 ⁺ to 2 ⁺)-1408(2 ⁺ to g.s.) cascade.
17019.6 ^o 19		C		J ^π : γ to (15 ⁻) suggests (15,16,17), (17 ⁻)

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) **^{58}Ni Levels (continued)**

E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF	O	T	Comments
17163.1 ^{<i>m</i>} 13	(16 ⁺)		C			from possible band assignment.
17197 3			C			
17290.0 ^{<i>w</i>} 11	(16 ⁺)		C			
17.42×10 ³ 25		3.9 MeV 4		O	T	E(level): L=1, giant-dipole resonance. Γ is from (α, α') . Other: 5.0 MeV 3 in (e, e'). %p=11 3 (2009Jo03) E(p)(c.m.)=2.35 MeV 6 (2009Jo03). Prompt p decay populates 6976, 21/2 ⁻ level in ^{57}Co which deexcites through 1058-1104-2290-834-466-1224 cascade to ^{57}Co g.s.
17482 3	(15 ⁻)		C			
17530.0 9	(17 ⁺)		C			%p=66 5 (2009Jo03); % α <10 (2009Jo03) E(p)(c.m.)=2.43 MeV 4, E(α)(c.m.)=7.71 MeV 8 (2009Jo03).
17582 3	(15 ⁻)		C			Prompt p decay populates 6976, 21/2 ⁻ level in ^{57}Co which deexcites through 1058-1104-2290-834-466-1224 cascade to ^{57}Co g.s.
17607 3	(15 ⁻)		C			Prompt α decay populates 2949, 6 ⁺ level in ^{54}Fe which deexcites through 411(6 ⁺ to 4 ⁺)-1130(4 ⁺ to 2 ⁺)-1408(2 ⁺ to g.s.) cascade. %p=43 4 (2009Jo03) E(p)(c.m.)=2.47 MeV 7 (2009Jo03). Prompt p decay populates 6976, 21/2 ⁻ level in ^{57}Co which deexcites through 1058-1104-2290-834-466-1224 cascade to ^{57}Co g.s.
17681.4 9	(17 ⁺)		C			
18261.1 ^{<i>n</i>} 14	(17 ⁺)		C			
18341.5 ^{<i>j</i>} 24	(16 ⁻)		C			
18.43×10 ³ 15		7.41 MeV 13		T		E(level): L=0, giant-monopole resonance.
18461.0 ^{<i>v</i>} 12	(17 ⁻)		C			
18638.9 ^{<i>s</i>} 10	(18 ⁺)		C			
19196 ^{<i>p</i>} 4			C			J ^π : γ to (16 ⁻) suggests (16,17,18), (18 ⁻) from possible band assignment.
19205.4 ^{<i>k</i>} 25	(17 ⁻)		C			
19482.5 ^{<i>m</i>} 16	(18 ⁺)		C			
19566.9 ^{<i>w</i>} 19	(18 ⁺)		C			
19945.7 ^{<i>t</i>} 11	(19 ⁺)		C			
20135.4 ^{<i>j</i>} 25	(18 ⁻)		C			%p<10 (2009Jo03) E(p)(c.m.)=1.94 MeV 7 (2009Jo03). Prompt p decay populates 10075, 25/2 ⁺ level in ^{57}Co .
20450.1 ^{<i>v</i>} 18	(19 ⁻)		C			
20826.2 ^{<i>n</i>} 23	(19 ⁺)		C			
21106.3 ^{<i>k</i>} 25	(19 ⁻)		C			%p<10 (2009Jo03) E(p)(c.m.)=1.89 MeV 7 (2009Jo03). Prompt p decay populates 11069, 27/2 ⁺ level in ^{57}Co .
21248.0 ^{<i>s</i>} 13	(20 ⁺)		C			

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) **^{58}Ni Levels (continued)**

E(level) [†]	J ^π [‡]	T _{1/2} [#]	XREF	Comments
22138 ^w 3	(20 ⁺)	C		
22211.3 ^j 25	(20 ⁻)	C		
22239.6 ^m 25	(20 ⁺)	C		
22767.9 ^t 15	(21 ⁺)	C		
22800.4 ^v 22	(21 ⁻)	C		
23331 ^k 3	(21 ⁻)	C		
23741 ⁿ 4	(21 ⁺)	C		
24.03×10 ³ 19		4.3 MeV 26	T	E(level): isoscalar giant-dipole resonance (ISGDR).
24211.9 ^s 17	(22 ⁺)	C		
24611 ^j 3	(22 ⁻)	C		
25141 ^w 4	(22 ⁺)	C		
25552 ^v 3	(23 ⁻)	C		
25918 ^k 3	(23 ⁻)	C		
26059.7 ^t 20	(23 ⁺)	C		
27366 ^j 4	(24 ⁻)	C		
28709 ^v 3	(25 ⁻)	C		
28931 ^k 3	(25 ⁻)	C		
30491 ^j 4	(26 ⁻)	C		
31.13×10 ³ 14		7.8 MeV 27	O T	E(level): isoscalar giant-dipole resonance (ISGDR). In (e,e'), composite energy is 28.3 MeV 3.
32175 3	(27 ⁻)	C		
32495 ^k 3	(27 ⁻)	C		
33972 ^j 4	(28 ⁻)	C		
36045 4	(29 ⁻)	C		
36535 ^k 3	(29 ⁻)	C		
37810 ^j 4	(30 ⁻)	C		
40333 4	(31 ⁻)	C		
40931 ^k 3	(31 ⁻)	C		
42007 ^j 4	(32 ⁻)	C		
x ^u		C		Additional information 2.
2868.1+x ^u 10		C		
6083.2+x ^u 15		C		
9667.3+x ^u 18		C		

[†] From a least-squares fit to Eγ's for levels populated in γ-ray studies. For levels populated in particle-transfer and inelastic scattering studies, the values are averaged over all available data. For levels populated in (γ,γ'), values are as given in the $^{58}\text{Ni}(\gamma,\gamma')$ dataset.

[‡] For high-spin (J>6) levels, all assignments are from γ-ray cascades observed in in-beam γ-ray studies: $^{28}\text{Si}(^{36}\text{Ar},\alpha 2\gamma)$, $^{28}\text{Si}(^{32}\text{S},2\gamma)$; $^{40}\text{Ca}(^{24}\text{Mg},\alpha 2\gamma)$; $^{45}\text{Sc}(^{16}\text{O},\text{p}2\gamma)$; $^{48}\text{Ti}(^{12}\text{C},2\text{n}\gamma)$ and $^{56}\text{Fe}(\alpha,2\text{n}\gamma)$. For many transitions angular distribution/correlation data support these assignments. For a few transitions, supporting γ(lin pol) data are available. In addition, ascending spins are assumed in these reactions as the excitation energy rises. Arguments for individual are given for levels below 10 MeV. Above this energy, all assignments are as proposed in $^{28}\text{Si}(^{36}\text{Ar},\alpha 2\gamma)$ reaction by [2009Jo03](#) and their previous papers, based on DCO data for selected transitions and γ cascades. The parentheses have been added by the evaluators since strong supporting arguments from polarization or other parity-sensitive seem to be lacking. When J^π is deduced from L-transfers, target J^π=0⁺ in all reactions.

Adopted Levels, Gammas (continued) **^{58}Ni Levels (continued)**

- [#] From DSA in ($p,p'\gamma$) ([1969Be48](#)), except where noted otherwise. Weighted or unweighted averages are taken when values are available from different reactions. Values from (γ,γ') are deduced from measured Γ_0^2/Γ and branching ratios.
- [@] From Γ_0^2/Γ or Γ_0 in (γ,γ') and adopted branching ratios, assuming $\Gamma(0)/\Gamma=1$ when there is only the ground-state transition listed from a level. See (γ,γ') dataset for details.
- [&] A level assumed by the evaluators to assign γ transitions to 1454 and g.s. in ($p,p'\gamma$). These transitions could not be assigned to levels in (p,p') because their J^π was 3^- or 4^+ .
- ^a From analysis of form factor in (e,e').
- ^b (0^+) from L(p,t)=(0) for 5960 40.
- ^c (4^+) from L(p,t)=(4) for 6560 40.
- ^d ($2^+,3^-$) from L(${}^3\text{He},n$)=(2,3) for 6900 50.
- ^e 4^+ from (e,e') for 6930 15.
- ^f 3^- from L(${}^3\text{He},{}^3\text{He}'$)=3 for 6780 25.
- ^g (1^+) from strong population of L(p,p')=0 state in near 0° data ([2007Fu04](#)), and interpretation as $>$ transition.
- ^h From analysis of $\sigma(\theta)$ and analyzing power data in (pol,p').
- ⁱ From $\gamma(\theta)$ and/or asymmetry measurement in (pol γ,γ').
- ^j Band(A): Band based on (16^-), $\alpha=0$. Parity from [2009Jo03](#) and [2006Ru02](#).
- ^k Band(a): Band based on (17^-), $\alpha=1$. Parity from [2009Jo03](#) and [2006Ru02](#).
- ^l Band(B): yrast structure.
- ^m Band(C): Band based on 15323,14 $^+$.
- ⁿ Band(D): Band based on 14455,13 $^+$.
- ^o Band(E): Band based on 9585,9 $^-$.
- ^p Band(F): $\Delta J=1$ band based on 10694,10 $^-$.
- ^q Band(G): $\Delta J=1$ band based on 3422,3 $^+$.
- ^r Band(H): $\Delta J=1$ band based on 7724,8 $^+$.
- ^s Band(I): Band based on 18638,18 $^+$.
- ^t Band(i): Band based on 19945,19 $^+$.
- ^u Band(J): γ cascade.
- ^v Band(K): SD-1 Band. BASED ON (15^-); from [2009Jo03](#), [2006Ru02](#) and [2001Ru03](#). This band has been assigned ([2001Ru03](#)) in the secondary minimum of the potential well. Population intensity $\approx 2\%$, relative to the total ${}^{58}\text{Ni}$ channel. The (13^-) states at 15410 and 15431 are possibly continuation of this band towards low-lying states. The (15^-) member of this band decays by prompt α emission to ${}^{54}\text{Fe}$. Average Q(transition)=2.4 3 ([2001Ru03](#)), from residual Doppler-shift method.
- ^w Band(L): SD-2 band. Based on (12^+); from [2009Jo03](#) and [2001Ru03](#). This band has been assigned ([2001Ru03](#)) in the secondary minimum of the potential well. Population intensity $\approx 1\%$, relative to the total ${}^{58}\text{Ni}$ channel.

Adopted Levels, Gammas (continued)

 $\gamma(^{58}\text{Ni})$

$E_i(\text{level})$	J_i^π	E_γ^\ddagger	I_γ^\ddagger	E_f	J_f^π	Mult. [†]	δ^\ddagger	$\alpha^\#$	$I_{(\gamma+ce)}$	Comments
1454.21	2 ⁺	1454.28 10	100	0.0	0 ⁺	E2				B(E2)(W.u.)=10.0 4
2459.21	4 ⁺	1004.80 15	100	1454.21	2 ⁺	E2				B(E2)(W.u.)=11.2 12
		2459.1	≤ 0.5	0.0	0 ⁺					
2775.42	2 ⁺	316.1	≤ 0.06	2459.21	4 ⁺					B(M1)(W.u.)=0.011 +3-4; B(E2)(W.u.)=15 +4-5
		1321.2 2	100.0 3	1454.21	2 ⁺	E2+M1	-1.1 1			Mult.: large $\delta(Q+D)$ from $\gamma(\theta)$.
2902.15	1 ⁺	2775.5 4	4.5 3	0.0	0 ⁺	E2				B(E2)(W.u.)=0.029 +8-10
		442.7	≤ 0.15	2459.21	4 ⁺					
		1448.2 4	100.0 6	1454.21	2 ⁺					
		2901.3 5	6.4 6	0.0	0 ⁺	(M1)				
2942.56	0 ⁺	40.3 4	100 4	2902.15	1 ⁺	[M1]		0.581 19		B(M1)(W.u.)=0.00079 +18-19
		167.2 2	18.4 19	2775.42	2 ⁺	[E2]		0.0809		Mult.: $\Delta J=1$, dipole from $\gamma(\theta)$; ΔJ^π requires M1.
		483.3 ^a	≤ 0.4	2459.21	4 ⁺					$\alpha(K)= 0.519 \ 17$; $\alpha(L)= 0.0541 \ 18$; $\alpha(M)=0.000762 \ 25$
		1488.3 3	19.9 19	1454.21	2 ⁺	[E2]				B(M1)(W.u.)=0.116 14
		2942.3		0.0	0 ⁺	E0				$\alpha(K)= 0.0722$; $\alpha(L)=0.00761$; $\alpha(M)=0.001063$
										B(E2)(W.u.)=21 3
3037.86	2 ⁺	95.2	≤ 0.5	2942.56	0 ⁺					
		135.8	≤ 0.2	2902.15	1 ⁺					
		262.6 3	1.7 3	2775.42	2 ⁺	M1+(E2)	-0.03 5			B(M1)(W.u.)=0.21 5; B(E2)(W.u.)=5 +18-5
		578.5	≤ 0.5	2459.21	4 ⁺					If M1, B(M1)(W.u.)=0.3 1.
		1583.8 3	100.0 17	1454.21	2 ⁺	M1+E2	+0.21 3			B(M1)(W.u.)=0.055 8; B(E2)(W.u.)=1.8 6
		3037.7 3	68.4 19	0.0	0 ⁺	[E2]				B(E2)(W.u.)=1.15 17
3263.66	2 ⁺	321	≤ 0.3	2942.56	0 ⁺					
		361.6	≤ 0.3	2902.15	1 ⁺					
		488.2	≤ 0.3	2775.42	2 ⁺					
		804.3	≤ 1.7	2459.21	4 ⁺					
		1809.5 3	65.8 18	1454.21	2 ⁺	M1+E2	+0.7 4			B(M1)(W.u.)=0.027 11; B(E2)(W.u.)=8 6
		3263.4 4	100.0 18	0.0	0 ⁺	[E2]				B(E2)(W.u.)=1.9 3
3269.1	(2)	3269.1 8		0.0	0 ⁺					
3273.7	(2)	3273.7 7		0.0	0 ⁺					
3420.55	3 ⁺	382.9 3	5.7 3	3037.86	2 ⁺	(M1+(E2))	+0.08 9			B(M1)(W.u.)=(0.08 +3-7); B(E2)(W.u.)=(7 +15-7)
		477.9	≤ 0.6	2942.56	0 ⁺					
		518.5	≤ 0.7	2902.15	1 ⁺					
		645.1	≤ 1.2	2775.42	2 ⁺					
		961.0 2	100.0 3	2459.21	4 ⁺	(M1+(E2))	-0.02 3			B(M1)(W.u.)=(0.09 +4-8); B(E2)(W.u.)=(0.07 +23-7)
		1966.3	≤ 2.1	1454.21	2 ⁺					

Adopted Levels, Gammas (continued)

 $\gamma(^{58}\text{Ni})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\ddagger	I_γ^\ddagger	E_f	J_f^π	Mult. [†]	δ^\ddagger	$I_{(\gamma+ce)}$	Comments
3420.55	3 ⁺	3420.3	≤ 0.4	0.0	0 ⁺				
3450.9		3450.9	5		0.0	0 ⁺			
3531.1	0 ⁺	493.3	≤ 1.0	3037.86	2 ⁺				
		588.5 ^a		2942.56	0 ⁺				
		629.1	≤ 1.1	2902.15	1 ⁺				
		755.7	≤ 1.8	2775.42	2 ⁺				
		1071.8 ^a	≤ 2.9	2459.21	4 ⁺				
		2076.9	3	100	1454.21	2 ⁺	[E2]		
		3530.9			0.0	0 ⁺	E0	0.068 11	$B(E2)(\text{W.u.})=5.6$ 18 $q_k^2(E0/E2)=0.27$ 4, $X(E0/E2)=0.47$ 7, $\rho^2(E0)=0.08$ 3 (2005Ki02 evaluation).
3593.71	1,2 ⁺	330.0	≤ 1.1	3263.66	2 ⁺				
		555.8	≤ 2.4	3037.86	2 ⁺				
		652.8	10	8.3	10	2942.56	0 ⁺		
		691.6	<1.5	2902.15	1 ⁺				
		818.4	4	27.2	15	2775.42	2 ⁺		
		1134.3		≤ 3.5	2459.21	4 ⁺			
		2139.2	5	18.0	10	1454.21	2 ⁺		
		3593.3	6	100.0	22	0.0	0 ⁺		
									I_γ : other: branching=24% 3 in (γ, γ') work of 2000Ba63 seems in error, the level in (γ, γ') is considered as the same as in other reactions and decays.
3620.09	4 ⁺	582.4	<3.8	3037.86	2 ⁺				
		844.8	<3.8	2775.42	2 ⁺				
		1161.2	3	100.0	24	2459.21	4 ⁺	(M1(+E2))	$+0.6 +3-6$
		2166.3	5	20.5	24	1454.21	2 ⁺	E2	$B(M1)(\text{W.u.})=(0.07 +4-6); B(E2)(\text{W.u.})=(4.E+1 4)$ $B(E2)(\text{W.u.})=1.3 +7-10$
		3620.0 ^a		<4.4		0.0	0 ⁺		
3775.0	3 ⁺	354.5	3	33	3	3420.55	3 ⁺	(M1(+E2))	$+0.05 +21-12$
		736	2	16	3	3037.86	2 ⁺		$B(M1)(\text{W.u.})=(0.33 +9-17); B(E2)(\text{W.u.})=(1.E+1 +11-1)$ If M1, $B(M1)(\text{W.u.})=0.019$ 9.
		872.6		<4.3	2902.15	1 ⁺			
		999.2			2775.42	2 ⁺			
		1316.4	15	100	7	2459.21	4 ⁺	M1(+E2)	$+0.19 15$
		2320.5	8	24	3	1454.21	2 ⁺		$B(M1)(\text{W.u.})=(0.019 +5-10); B(E2)(\text{W.u.})=(0.8 +12-8)$ If M1, $B(M1)(\text{W.u.})=9\times 10^{-4}$ 5.
		3774.4		<5		0.0	0 ⁺		
3898.8	2 ⁺	2444.7	4	100.0	16	1454.21	2 ⁺	[M1]	$B(M1)(\text{W.u.})=0.050$ 13 $B(E2)(\text{W.u.})=0.50$ 14
		3898.0	7	31.9	16	0.0	0 ⁺	[E2]	I_γ : from (p,p'γ).
3943.6		3943.6	12			0.0	0 ⁺		
4105.9	(4 ⁺)	486.0	3	6	3	3620.09	4 ⁺		
		683.7	5	3	3	3420.55	3 ⁺		
		1646.4	12	9	3	2459.21	4 ⁺		
		2653.1	12	100	3	1454.21	2 ⁺	(Q)	
4108.4	2 ⁺	687.4		4.3	22	3420.55	3 ⁺		

Adopted Levels, Gammas (continued)

 $\gamma(^{58}\text{Ni})$ (continued)

19

E _i (level)	J _i ^π	E _γ [‡]	I _γ [‡]	E _f	J _f ^π	Mult. [†]	δ [‡]	Comments
4108.4	2 ⁺	1205.9	11 4	2902.15	1 ⁺			
		1332.5	13 4	2775.42	2 ⁺	[M1]		B(M1)(W.u.)=0.0044 24
		2654.6 4	87 5	1454.21	2 ⁺	M1+E2	-0.58 +8-9	B(M1)(W.u.)=0.0028 13; B(E2)(W.u.)=0.26 13
		4107.4 7	100 5	0.0	0 ⁺	[E2]		B(E2)(W.u.)=0.13 6
4294.7	4 ⁺	1835.3 4	67 33	2459.21	4 ⁺			
		2840.8 10	100 33	1454.21	2 ⁺	[E2]		B(E2)(W.u.)=6 +5-6
4347.9		2893.6 12	100	1454.21	2 ⁺			If E2, B(E2)(W.u.)=12 11.
4358.7	(2 ^{+,3,4})	1584		2775.42	2 ⁺			
		1901.7 12		2459.21	4 ⁺			
		2902		1454.21	2 ⁺			
		276.7 6	3.8 7	4105.9 (4 ⁺)	D+Q			
4383.0	(5 ⁺)	763.0 3	100.0 20	3620.09	4 ⁺	(M1+E2)	-0.38 5	
		962 1	0.2 2	3420.55	3 ⁺			
		1923.9 7	31.2 20	2459.21	4 ⁺	D+Q	+0.27 10	
		2951.3 11	100	1454.21	2 ⁺	E2		B(E2)(W.u.)=4.4 +15-18
4404.3	4 ⁺	855.0 4	100 10	3593.71	1,2 ⁺			
		1547.0 7	11 3	2902.15	1 ⁺			
4449.6	1 ^{+,2+}	1673.8 6	12.4 17	2775.42	2 ⁺			
		1697.5 9	25 8	2775.42	2 ⁺	[E1]		B(E1)(W.u.)=0.0008 4
		3021.1 6	100 17	1454.21	2 ⁺	[E1]		B(E1)(W.u.)=0.00060 22
		2059		2459.21	4 ⁺			
4518.3		3064		1454.21	2 ⁺			
		3083.7 6	100	1454.21	2 ⁺	[E2]		B(E2)(W.u.)=4.9 18
4538.0	0 ⁺	4574.1 ^{@a}		0.0	0 ⁺			
4574.1	1	1132	100	3620.09	4 ⁺			
		2293	25	2459.21	4 ⁺			
4752.2	4 ⁺	1300		3620.09	4 ⁺			
		1656		3263.66	2 ⁺			
		2461		2459.21	4 ⁺			
		4954.0 8		0.0	0 ⁺			
4964.7	(5 ⁺)	1344.7 2	23 3	3620.09	4 ⁺			
		2503.8 13	100 3	2459.21	4 ⁺	D+Q	-0.42 4	
5064.3		2605	100	2459.21	4 ⁺			
		723.2 2	0.6 2	4404.3	4 ⁺			
5127.5	6 ⁺	744.6 3	100.0 21	4383.0 (5 ⁺)	(M1+E2)	-0.42 4		δ: from (³⁶ Ar,α2pγ). Other: -2.5 +6-8 or -0.20 +10-15 in (¹² C,2nγ).
		832.0 ^a 7	0.4 2	4294.7	4 ⁺			
		1020.3 7	1.46 21	4105.9 (4 ⁺)				
		2668.6 10	45.8 21	2459.21	4 ⁺	Q		
5170.3		2711		2459.21	4 ⁺			
		5359.3 ^{@a} 16		0.0	0 ⁺			
5359.3?	(2)	1000.8 8	100 10	4383.0 (5 ⁺)	D+Q			
5384.5	6 ⁺							

Adopted Levels, Gammas (continued)

 $\gamma(^{58}\text{Ni})$ (continued)

E _i (level)	J _i [¶]	E _γ [‡]	I _γ [‡]	E _f	J _f ^π	Mult. [†]	δ [‡]	Comments
5384.5	6 ⁺	1088.9 10 1764.5 11 2926.6 15	4 2 100 10 66 8	4294.7 3620.09 2459.21	4 ⁺ 4 ⁺ 4 ⁺	Q Q		
5394.0		5394.0 9		0.0	0 ⁺			
5436.3	4 ⁺	2977	100	2459.21	4 ⁺			
5452.2	1	5452.2 @ 4		0.0	0 ⁺			
5472.3	4 ⁺	3013		2459.21	4 ⁺			
		4018		1454.21	2 ⁺			
5503.5		2728	100	2775.42	2 ⁺			
5528.0	(1)	5528.0 4		0.0	0 ⁺			
5589.0	(5 ⁻)	3129.0 15	100	2459.21	4 ⁺			
5590.3	2 ⁺	5590	100	0.0	0 ⁺			
5594.2	4 ⁺	1819 3135 4140	53 100 3	3775.0 2459.21 1454.21	3 ⁺ 4 ⁺ 2 ⁺			E _γ : from (p,p'γ). E _γ : all γ's from (p,p'γ).
5706.3		2931 ^a 3247 4252		2775.42 2459.21 1454.21	2 ⁺ 4 ⁺ 2 ⁺			
5744.7	(6 ⁺)	3286.0 18	100	2459.21	4 ⁺	Q		
5748.5	2 ⁺	2155 3289 4294 ^a		3593.71 2459.21 1454.21	1,2 ⁺ 4 ⁺ 2 ⁺			
5766.3	4 ⁺	3307 4312		2459.21 1454.21	4 ⁺ 2 ⁺			
5803.3		4349 5803		1454.21 0.0	2 ⁺ 0 ⁺			
5824.6		2404		3420.55	3 ⁺			
5896.4		4442 5896		1454.21 0.0	2 ⁺ 0 ⁺			
5905.3	1 ⁺	5905.3 7		0.0	0 ⁺	M1		B(M1)(W.u.)=0.0043 7
5942.4	(0 ⁺)	4488	100	1454.21	2 ⁺			
6018.4	3 ⁻	4564	100	1454.21	2 ⁺			
6027.3	1 ⁻	3565 4574.1 @ 5		2459.21 1454.21	4 ⁺ 2 ⁺			
		6027.3 7	100 4	0.0	0 ⁺	E1		B(E1)(W.u.)=0.00197 17
6067.5	(7 ⁺)	322.8 2 682.7 5 940.1 4 1684.6 10	1.6 3 16.1 13 100 5 42 3	5744.7 (6 ⁺) 5384.5 6 ⁺ 5127.5 6 ⁺ 4383.0 (5 ⁺)	(M1+E2) (M1+E2) (M1+E2) (M1+E2)	-0.18 10 -0.11 8 -0.36 4		
6084.7	7 ⁻	495.6 6 699.6 8	2.4 12 23.5 24	5589.0 (5 ⁻) 5384.5 6 ⁺	Q E1(+M2)	-0.06 13		

Adopted Levels, Gammas (continued)

 $\gamma(^{58}\text{Ni})$ (continued)

E _i (level)	J _i ^π	E _γ [‡]	I _γ [‡]	E _f	J _f ^π	Mult. [†]	δ [†]	Comments
6084.7	7 ⁻	957.1 7 3625.1 13	100 4 10.6 12	5127.5 2459.21	6 ⁺ 4 ⁺	E1(+M2) (E3)	-0.06 5	Mult.: DCO in (³⁶ Ar, α 2p γ) consistent with pure octupole or $\Delta J=2,Q$, not with $\Delta J=1$, dipole.
6174.3	2 ^{+,3⁻}	3715 4720 ^a 6174		2459.21 1454.21 0.0	4 ⁺ 2 ⁺ 0 ⁺			
6220.0	(7 ⁺)	835.5 6 1092.7 5 1256.4 9	100 9 88 9 40 3	5384.5 5127.5 4964.7 (5 ⁺)	6 ⁺ 6 ⁺ (Q)	D+Q D+Q	-0.08 4	
6228.3	(2 ⁺)	3326 4774 6228		2902.15 1454.21 0.0	1 ⁺ 2 ⁺ 0 ⁺			
6274.3	4[+]	3815	100	2459.21	4 ⁺			
6308.5	3 ⁻	3366 3533 4854		2942.56 2775.42 1454.21	0 ⁺ 2 ⁺ 2 ⁺			
6360.6		2940	100	3420.55	3 ⁺			
6402.4		6402	100	0.0	0 ⁺			
6424.9?	1	6424.9 @ ^a 9		0.0	0 ⁺			
6430.7?	1	6430.7 ^a 10		0.0	0 ⁺			
6468.4	(1 ⁺)	5014 6468		1454.21 0.0	2 ⁺ 0 ⁺			
6478.4	2 ⁺	5024 6478		1454.21 0.0	2 ⁺ 0 ⁺			
6507.2		2887	100	3620.09	4 ⁺			
6571.4	2 ⁺	5117	100	1454.21	2 ⁺			
6601.3		2981 4142		3620.09 2459.21	4 ⁺ 4 ⁺			
6604.6	(8 ⁺)	384.8 3 519.5 4 537.0 3 1476.8 10	2.6 3 2.1 3 100 3 70 3	6220.0 (7 ⁺) 6084.7 7 ⁻ 6067.5 (7 ⁺) 5127.5 6 ⁺		(M1+E2) Q	-0.18 3	δ: from (³⁶ Ar, α 2p γ). Other: -0.20 +5-9 in (¹² C,2n γ).
6665.4		5211 6665		1454.21 0.0	2 ⁺ 0 ⁺			
6685.0?	1	6685.0 @ ^a 9		0.0	0 ⁺			
6717.4		5263 6717		1454.21 0.0	2 ⁺ 0 ⁺			
6763.5	3 ⁻	5309	100	1454.21	2 ⁺			
6805.5	3 ⁻	5351	100	1454.21	2 ⁺			
6845.7	(7 ⁺)	1718.0 10 2463.0 19	50 50 100 50	5127.5 4383.0 (5 ⁺)	6 ⁺			

Adopted Levels, Gammas (continued)

 $\gamma(^{58}\text{Ni})$ (continued)

E _i (level)	J _i ^π	E _γ [†]	I _γ [‡]	E _f	J _f ^π	Mult. [†]	δ [†]	Comments
6854.5	3 ⁻	5400	100	1454.21	2 ⁺			
6863.1	(6)	2478.9 18	100	4383.0	(5 ⁺)	D		
6892.9?	(1)	6892.9 ^a 15		0.0	0 ⁺			
6992.5		5538	100	1454.21	2 ⁺			
7048.2	1 ⁻	7048.2 9		0.0	0 ⁺	E1		B(E1)(W.u.)=0.00155 6
7054.5		7054	100	0.0	0 ⁺			
7113.5	(1,2 ⁺)	5659		1454.21	2 ⁺			
		7113		0.0	0 ⁺			
7131.5		7131	100	0.0	0 ⁺			
7210.4	3 ⁻	4751	100	2459.21	4 ⁺			
7249.6	(1)	7249.6 11		0.0	0 ⁺			
7271.7	1	5817		1454.21	2 ⁺			
		7271.7 7		0.0	0 ⁺			
7273.7	7 ⁻	410.5 3	8.5 21	6863.1	(6)			
		1189.9 8	57 4	6084.7	7 ⁻	D+Q		
		2146.4 15	100 6	5127.5	6 ⁺	E1+M2	-0.19 6	
7300.5	3 ⁻	5846	100	1454.21	2 ⁺			
7314.8	(8 ⁺)	709.7 5	48 4	6604.6	(8 ⁺)			
		1095.7 9	78 7	6220.0	(7 ⁺)			
		1245.9 9	100 7	6067.5	(7 ⁺)	D+Q	-0.15 5	
		1930.3 14	81 7	5384.5	6 ⁺	Q		
7380.5	(1,2 ⁺)	7380	100	0.0	0 ⁺			
7388.8	1 ⁺	7388.8 4		0.0	0 ⁺	M1		
7446.2	(9 ⁺)	841.6 4	100 3	6604.6	(8 ⁺)	(M1+E2)	-0.18 3	B(M1)(W.u.)=0.055 3
		1226.1 9	14.1 17	6220.0	(7 ⁺)	Q		
		1378.6 10	5.5 3	6067.5	(7 ⁺)			
7514.5	3 ⁻	6060	100	1454.21	2 ⁺			
7570.5	2 ⁺	7570	100	0.0	0 ⁺			
7585.1		7585.1 6		0.0	0 ⁺			
7595.9	(2)	7595.9 6		0.0	0 ⁺			
7616.0?	(1)	7616.0 ^a 10		0.0	0 ⁺			
7680.6	1 ⁻	6226	100	1454.21	2 ⁺			
7709.7	1 ⁺	7709.7 6		0.0	0 ⁺	M1		B(M1)(W.u.)=0.067 3
7724.3	(8 ⁺)	878.4 9	50 17	6845.7	(7 ⁺)			
		1119.6 4	100 17	6604.6	(8 ⁺)	D+Q		
		1639.0 10	50 17	6084.7	7 ⁻			
		1657.0 10	50 17	6067.5	(7 ⁺)			
		2343.0 20	67 17	5384.5	6 ⁺	Q		
7766.0	(1)	7766.0 7		0.0	0 ⁺			
7807.3	1 ⁻	6356		1454.21	2 ⁺			
		7807.3 5	100	0.0	0 ⁺	E1		B(E1)(W.u.)=0.00117 15

Adopted Levels, Gammas (continued)

 $\gamma(^{58}\text{Ni})$ (continued)

E _i (level)	J _i ^π	E _γ [‡]	I _γ [‡]	E _f	J _f ^π	Mult. [†]	δ [†]	Comments
7862.6	(1,2 ⁺)	6408 7862		1454.21 0.0	2 ⁺ 0 ⁺			
7876.7	1	6424.9 @ 9 7876.7 26	45 36 100 36	1454.21 0.0	2 ⁺ 0 ⁺			
7973.6	(8 ⁺)	1370.0 10 1752.0 11 2229.6 16	100 5 76 10 43 5	6604.6 (8 ⁺) 5744.7 (6 ⁺)		D+Q Q	-0.37 8	
7982.8	(8 ⁻)	709.2 5 1120.2 8 1915.6 13	100 4 6.5 22 65.2 22	7273.7 (6) 6863.1 (6)	7 ⁻ (M1+E2)	-0.15 3		
8068.6?	(1 ⁻)	8068.6 @a 12		0.0	0 ⁺	(E1)		B(E1)(W.u.)=0.00062 8
8074.5	(8 ⁺)	1470 1 1854.3 13	55 5 100 5	6604.6 (8 ⁺) 6220.0 (7 ⁺)	0 ⁺ 0 ⁺	D+Q	-0.21 8	
8096.3	1	8096.3 6		0.0	0 ⁺			
8110.6	(1,2 ⁺)	8110	100	0.0	0 ⁺			
8115.1	(8 ⁻)	2031.0 14	100	6084.7	7 ⁻			
8120.8	(9 ⁺)	396.5 1 805.5 5 1516.6 7	6.4 9 6.4 9 100 5	7724.3 (8 ⁺) 7314.8 (8 ⁺) 6604.6 (8 ⁺)	D+Q D+Q D+Q		-0.13 4	
8237.3	1 ⁻	8237.3 4		0.0	0 ⁺	E1		B(E1)(W.u.)=0.0054 +8-11
8317.1	1	8317.1 17		0.0	0 ⁺			
8395.1	1 ⁻	5359.3 @ 16		3037.86	2 ⁺			I _γ : 35 16 for 5359.3 γ +5452.2 γ , assuming the main placements of these γ rays are from 8395 level.
		5452.2 @ 4 8395.1 12		2942.56 0.0	0 ⁺ 0 ⁺	E1		B(E1)(W.u.)=0.0019 6
8461.0	1 ⁺	8461.0 7		0.0	0 ⁺	M1		B(M1)(W.u.)=0.071 5
8514.1	1 ⁻	8514.1 4		0.0	0 ⁺	E1		B(E1)(W.u.)=0.00111 9
8552.7	1 ⁽⁺⁾	8552.7 13		0.0	0 ⁺	[M1]		B(M1)(W.u.)=0.036 3
8600.5	1 ⁺	8600.5 7		0.0	0 ⁺	M1		B(M1)(W.u.)=0.061 7
8679.3	1 ⁺	8679.3 8		0.0	0 ⁺	M1		B(M1)(W.u.)=0.151 8
8718.1	(9 ⁻)	603.0 4 735.4 5 1403.2 10 1444.4 10 2114.0 15	4.4 11 51.6 22 20.9 22 23.1 11 100 3	8115.1 (8 ⁻) 7982.8 (8 ⁺) 7314.8 (8 ⁺) 6604.6 (8 ⁺)	D+Q (M1+E2) D+Q D+Q D(+Q)	-0.16 3 -0.13 10 -0.03 4		
8857.4	1 ⁽⁺⁾	8857.4 6	100	0.0	0 ⁺	[M1]		B(M1)(W.u.)=0.052 11
8880.2	1 ⁻	8880.2 6		0.0	0 ⁺	E1		B(E1)(W.u.)=0.00165 8
8896.4?		1581.6 11	100	7314.8	(8 ⁺)			
8934.6	1 ⁽⁻⁾	8934.6 5		0.0	0 ⁺	(E1)		B(E1)(W.u.)=0.00204 8
8961.3	1 ⁺	8961.3 7	100	0.0	0 ⁺	[M1]		B(M1)(W.u.)=0.026 3
9027.2	(9 ⁻)	912.3 6	17 6	8115.1	(8 ⁻)			

Adopted Levels, Gammas (continued)

 $\gamma(^{58}\text{Ni})$ (continued)

E _i (level)	J _i ^π	E _γ [‡]	I _γ [‡]	E _f	J _f ^π	Mult. [†]	δ [†]	Comments
9027.2	(9 ⁻)	2942.2 21	100 6	6084.7	7 ⁻	Q		
9062.7	(10 ⁺)	941.1 7	100 4	8120.8	(9 ⁺)	M1+E2	-0.24 6	
		1336.5 28	4.3 14	7724.3	(8 ⁺)			
		1617.0 11	14.3 14	7446.2	(9 ⁺)			
		2459.9 17	40 3	6604.6	(8 ⁺)	Q		
9073.4	1 ⁺	9073.4 6		0.0	0 ⁺	(M1)		B(M1)(W.u.)=0.058 4
9156.9	1 ⁺	9156.9 7		0.0	0 ⁺	M1		B(M1)(W.u.)=0.037 5
9190.7	1 ⁻	9190.7 5		0.0	0 ⁺	E1		B(E1)(W.u.)=0.00100 11
9322.1	(11 ⁺)	1876.4 13	100	7446.2	(9 ⁺)	Q		
9326.4	1	6424.9 @ 9	39 10	2902.15	1 ⁺			
		9326.4 8	100 10	0.0	0 ⁺			
9345.5	(10 ⁻)	627.5 5	100 4	8718.1	(9 ⁻)	(M1+E2)	-0.15 3	
		1363.1 10	22.7 14	7982.8	(8 ⁻)	Q		
		1899.9 13	74 4	7446.2	(9 ⁺)	D+Q	-0.16 3	
9368.5	1 ⁽⁺⁾	9368.5 6		0.0	0 ⁺	[M1]		B(M1)(W.u.)=0.072 8 If E1, B(E1)(W.u.)=0.00148 16.
9455.4	1	9455.4 18		0.0	0 ⁺			
9523.3	1 ⁻	8068.6 @ 12	72 9	1454.21	2 ⁺			
		9523.3 13	100 9	0.0	0 ⁺	E1		B(E1)(W.u.)=0.0026 5
9554.0	1	9554.0 21		0.0	0 ⁺			
9585.2	(9 ⁻)	1511.5 11	67 17	8074.5	(8 ⁺)	D		
		1610.6 11	100 17	7973.6	(8 ⁺)	D		
		3498.7 24	67 17	6084.7	7 ⁻	Q		
9630.5	1	9630.5 24		0.0	0 ⁺			
9666.9	(10 ⁺)	1592.2 11	53 6	8074.5	(8 ⁺)			
		1694.2 12	100 12	7973.6	(8 ⁺)	Q		
		2219.5 16	47 6	7446.2	(9 ⁺)	D+Q		
		3062.0 21	41 6	6604.6	(8 ⁺)			
9667.8	1	6892.9 @ 15	49 27	2775.42	2 ⁺			
		9667.8 15	100 27	0.0	0 ⁺			
9723.0	1 ⁽⁻⁾	6685.0 @ 9	139 13	3037.86	2 ⁺			
		9723.0 9	100 13	0.0	0 ⁺	(E1)		B(E1)(W.u.)=0.0019 4
9790.6	(10 ⁺)	2344.0 16	100	7446.2	(9 ⁺)	D+Q		
9843	1 ⁺	9842 5	100	0.0	0 ⁺	[M1]		B(M1)(W.u.)=0.09 +4-9
9886.8	(10 ⁺)	1811.4 13	69 6	8074.5	(8 ⁺)	Q		
		1913.2 4	100 6	7973.6	(8 ⁺)	Q		
10137.2	(10 ⁺)	2688.4 19	100 8	7446.2	(9 ⁺)			
		3533.0 20	92 8	6604.6	(8 ⁺)	Q		
10144.7	(10 ⁻)	799.1 6	100 7	9345.5	(10 ⁻)	D+Q		

Adopted Levels, Gammas (continued)

 $\gamma(^{58}\text{Ni})$ (continued)

E _i (level)	J ^{π} _i	E _{γ} [†]	I _{γ} [†]	E _f	J ^{π} _f	Mult. [†]	δ^{\dagger}
10144.7	(10 ⁻)	1117.8 8	21 7	9027.2 (9 ⁻)	D+Q		
		1426.1 10	21 7	8718.1 (9 ⁻)			
		2029.0 10	21 7	8115.1 (8 ⁻)			
		2162.9 9	14 7	7982.8 (8 ⁻)			
10180.8	(11 ⁻)	835.6 6	100 3	9345.5 (10 ⁻)	(M1+E2)	-0.09 4	
		1153.7 10	1.6 5	9027.2 (9 ⁻)			
		1463.9 10	12.1 16	8718.1 (9 ⁻)	Q		
10192.5	(11 ⁺)	1129.4 8	100 4	9062.7 (10 ⁺)	D+Q	-0.45 6	
		2072.7 15	22.4 15	8120.8 (9 ⁺)	Q		
		2746.6 19	39 3	7446.2 (9 ⁺)	Q		
10293.5	(9 ⁻)	3688.3 28	100 33	6604.6 (8 ⁺)			
		4207.7 30	33 33	6084.7 7 ⁻			
10394.1	(10 ⁺)	3078.0 22	50 50	7314.8 (8 ⁺)			
		3788.0 27	100 50	6604.6 (8 ⁺)			
10404.8	(9 ⁻)	4320.0 30	100	6084.7 7 ⁻			
10590.9	(11 ⁻)	410.3 3	31 3	10180.8 (11 ⁻)	D+Q		
		446.3 3	39 3	10144.7 (10 ⁻)	D+Q		
		1245.2 9	100 6	9345.5 (10 ⁻)	D+Q		
		1563.5 11	6 3	9027.2 (9 ⁻)			
10694.7	(10 ⁻)	1872.5 13	42 3	8718.1 (9 ⁻)	Q		
		289.9 2	10 10	10404.8 (9 ⁻)			
		401.0 10	20 10	10293.5 (9 ⁻)	D+Q		
		1350.0 10	10 10	9345.5 (10 ⁻)			
		1632.2 11	20 10	9062.7 (10 ⁺)			
		1798.2 13	20 10	8896.4?			
		2710.2 19	40 10	7982.8 (8 ⁻)	Q		
		3249.7 23	100 10	7446.2 (9 ⁺)	D		
10781.7	(11 ⁺)	991.1 7	50 10	9790.6 (10 ⁺)	D+Q		
		3336.0 23	100 10	7446.2 (9 ⁺)	Q		
10882.0	(11 ⁺)	1559.9 11	100	9322.1 (11 ⁺)	D+Q		
11005.6	(11 ⁻)	825.1 6	36 7	10180.8 (11 ⁻)	D+Q		
		1683.5 12	100 7	9322.1 (11 ⁺)			
11117.0	(11 ⁻)	1229.9 9	100 13	9886.8 (10 ⁺)	D+Q	-0.09 7	
		1531.2 11	88 13	9585.2 (9 ⁻)			
		2090.0 15	38 13	9027.2 (9 ⁻)	Q		
11255.2	(11 ⁻)	560.6 4	100 5	10694.7 (10 ⁻)	(M1+E2)	-0.26 5	
		1074.1 8	42 5	10180.8 (11 ⁻)			
11297.7	(12 ⁻)	707.0 5	38.8 11	10590.9 (11 ⁻)	(M1+E2)	-0.15 5	
		1116.3 8	100 3	10180.8 (11 ⁻)	D+Q	-0.22 3	
11413.1	(11 ⁺)	3966.2 28	100	7446.2 (9 ⁺)	Q		
11474.5	(12 ⁺)	1281.8 9	100 5	10192.5 (11 ⁺)	D+Q	-0.55 8	
		1807.5 13	35 3	9666.9 (10 ⁺)	Q		

Adopted Levels, Gammas (continued) $\gamma(^{58}\text{Ni})$ (continued)

E _i (level)	J ^π _i	E _γ [‡]	I _γ [‡]	E _f	J ^π _f	Mult. [†]	δ [‡]
11474.5	(12 ⁺)	2152.4 15	32 3	9322.1 (11 ⁺)	D+Q	-0.39 7	
		2410.9 17	30 3	9062.7 (10 ⁺)	Q		
11579.3	(12 ⁺)	1386.7 10	100 7	10192.5 (11 ⁺)	D+Q	-0.35 8	
		1692.7 10	27 7	9886.8 (10 ⁺)			
11814.3	(12 ⁻)	1223.8 9	52 4	10590.9 (11 ⁻)	D+Q	-0.08 2	
		1633.8 11	100 4	10180.8 (11 ⁻)	D+Q	-0.07 2	
		2467.9 17	56 4	9345.5 (10 ⁻)	Q		
11824.7	(12 ⁺)	1632.0 10	100	10192.5 (11 ⁺)	D+Q	-0.61 14	
11996.4	(12 ⁻)	741.4 5	100 6	11255.2 (11 ⁻)	D+Q		
		1301.0 10	12 6	10694.7 (10 ⁻)	Q		
		1406.2 10	12 6	10590.9 (11 ⁻)			
		1813.8 13	18 6	10180.8 (11 ⁻)			
12155.1	(12 ⁻)	1564.0 12	67 11	10590.9 (11 ⁻)	D+Q	+0.15 11	
		1974.1 14	100 11	10180.8 (11 ⁻)	D+Q	-0.27 10	
12356.8	(12 ⁻)	1351.1 9	100 10	11005.6 (11 ⁻)			
		1766.0 10	40 10	10590.9 (11 ⁻)			
12364.6	(12 ⁺)	1582.5 11	75 25	10781.7 (11 ⁺)			
		2171.4 15	50 25	10192.5 (11 ⁺)			
		2478.0 20	50 25	9886.8 (10 ⁺)			
		2697.0 20	50 25	9666.9 (10 ⁺)			
		3302.1 23	100 25	9062.7 (10 ⁺)	Q		
12570.1	(12 ⁺)	991.0 10	14 14	11579.3 (12 ⁺)			
		1157.0 8	29 7	11413.1 (11 ⁺)			
		1789.0 13	14 14	10781.7 (11 ⁺)			
		2174.9 15	14 14	10394.1 (10 ⁺)			
		2377.8 17	14 14	10192.5 (11 ⁺)			
		2390.1 17	100 14	10180.8 (11 ⁻)	D		
		2431.0 17	29 14	10137.2 (10 ⁺)			
		2682.9 21	43 14	9886.8 (10 ⁺)			
		3248.0 23	71 14	9322.1 (11 ⁺)	D+Q	-0.44 11	
		3507.0 25	14 14	9062.7 (10 ⁺)			
12719.2	(12 ⁺)	1306.0 10	33 33	11413.1 (11 ⁺)			
		2526.5 18	33 33	10192.5 (11 ⁺)			
		2928.0 20	33 33	9790.6 (10 ⁺)			
		3400.0 24	67 33	9322.1 (11 ⁺)			
		3655.0 26	100 33	9062.7 (10 ⁺)	Q		
12831.6	(13 ⁻)	1534.1 11	100 5	11297.7 (12 ⁻)			
		1713.6 12	77 5	11117.0 (11 ⁻)	Q		
		2652.2 19	41 5	10180.8 (11 ⁻)	Q		
12912.1	(13 ⁻)	915.7 6	100 5	11996.4 (12 ⁻)	D+Q		
		1657.0 12	30 5	11255.2 (11 ⁻)			
12928		3606.0 30	100	9322.1 (11 ⁺)			

Adopted Levels, Gammas (continued)

 $\gamma(^{58}\text{Ni})$ (continued)

E _i (level)	J ^π _i	E _γ [‡]	I _γ [‡]	E _f	J ^π _f	Mult. [†]
13016.6	(13 ⁻)	1718.3 12	100	11297.7	(12 ⁻)	D+Q
13048.2	(13 ⁻)	1749.8 12	100 7	11297.7	(12 ⁻)	D+Q
		2044.3 14	29 7	11005.6	(11 ⁻)	
13095.1	(12 ⁺)	3772.2 30	100	9322.1	(11 ⁺)	(D+Q)
13129.2	(12 ⁺)	3806.3 30	100 50	9322.1	(11 ⁺)	(D+Q)
13238.1	(13 ⁺)	518.9 4	40 2	12719.2	(12 ⁺)	D+Q
		668.0 5	100 4	12570.1	(12 ⁺)	D+Q
		873.3 6	44 4	12364.6	(12 ⁺)	D+Q
		1424.5 10	28 2	11814.3	(12 ⁻)	D
		1764.1 & 12	10 & 2	11474.5	(12 ⁺)	
		1941.7 14	30 2	11297.7	(12 ⁻)	D
		3045.0 21	2 2	10192.5	(11 ⁺)	
13356.6	(13 ⁺)	1881.5 13	100 17	11474.5	(12 ⁺)	
		3164.1 22	83 17	10192.5	(11 ⁺)	Q
13606.8	(12 ⁺)	3417.0 24	1×10 ² 1	10192.5	(11 ⁺)	
		4283.9 31	1×10 ² 1	9322.1	(11 ⁺)	D+Q
13632		4310 3	100	9322.1	(11 ⁺)	
13850.1	(14 ⁻)	938.0 7	100 5	12912.1	(13 ⁻)	D+Q
		1853.8 13	47 5	11996.4	(12 ⁻)	Q
13884.2	(13 ⁺)	755.0 10	33 33	13129.2	(12 ⁺)	
		789.0 10	33 33	13095.1	(12 ⁺)	(D+Q)
		2586.3 25	100 33	11297.7	(12 ⁻)	(D)
13943		3750.0 26	100	10192.5	(11 ⁺)	
14127.8	(14 ⁺)	889.6 6	100 5	13238.1	(13 ⁺)	D+Q
		1763.0 10	3.1 10	12364.6	(12 ⁺)	
14217.5	(14 ⁻)	1861.0 13	100 20	12356.8	(12 ⁻)	Q
		2062.0 15	80 20	12155.1	(12 ⁻)	Q
14455.8	(13 ⁺)	4261.7 30	100	10192.5	(11 ⁺)	
14853.1	(15 ⁻)	1835.6 13	8 4	13016.6	(13 ⁻)	
		2021.2 14	100 12	12831.6	(13 ⁻)	Q
14920.9	(14 ⁺)	1564.3 10	14 14	13356.6	(13 ⁺)	
		3445 2	100 14	11474.5	(12 ⁺)	Q
14934.7	(15 ⁻)	1084.8 8	100 11	13850.1	(14 ⁻)	D+Q
		2022.2 14	78 11	12912.1	(13 ⁻)	Q
15010.6	(14 ⁺)	1654.0 10	25 13	13356.6	(13 ⁺)	
		1773.0 10	25 13	13238.1	(13 ⁺)	
		3185.9 22	13 13	11824.7	(12 ⁺)	
		3431.0 24	25 13	11579.3	(12 ⁺)	
		3536.4 30	100 13	11474.5	(12 ⁺)	Q
15031.0	(14 ⁺)	1674.0 12	40 20	13356.6	(13 ⁺)	
		3206.0 30	20 20	11824.7	(12 ⁺)	

Adopted Levels, Gammas (continued)

 $\gamma(^{58}\text{Ni})$ (continued)

E _i (level)	J _i ^π	E _γ [‡]	I _γ [‡]	E _f	J _f ^π	Mult. [†]	δ [†]
15031.0	(14 ⁺)	3451.4 24 3556.0 25	20 20 100 20	11579.3 (12 ⁺) 11474.5 (12 ⁺)			
15105.2		1221.0 10 1976.0 14	1×10 ² 1 1×10 ² 1	13884.2 (13 ⁺) 13129.2 (12 ⁺)		Q	
15187.0	(13 ⁺)	2057.0 20 4997 4	1×10 ² 1 1×10 ² 1	13129.2 (12 ⁺) 10192.5 (11 ⁺)			
15242.0		2193.7 15	100	13048.2 (13 ⁻)			
15266.3	(14 ⁺)	2249.9 16 2435.6 17	29 14 100 14	13016.6 (13 ⁻) 12831.6 (13 ⁻)		D	
15294.3	(14 ⁺)	1688.0 12 2277.9 16 2462.2 17	100 11 22 11 44 11	13606.8 (12 ⁺) 13016.6 (13 ⁻) 12831.6 (13 ⁻)		Q	
15324.1	(14 ⁺)	3498.4 25 3849.0 27	13 13 100 13	11824.7 (12 ⁺) 11474.5 (12 ⁺)			
15434.1	(13 ⁻)	3436.5 ^a 24	1×10 ² 1	11996.4 (12 ⁻)			
		4136 4	1×10 ² 1	11297.7 (12 ⁻)			
15709.3	(15 ⁺)	1581.3 11 2470.0 17	100 7 29 7	14127.8 (14 ⁺) 13238.1 (13 ⁺)		D+Q	-0.22 4
15736.9	(15 ⁺)	706.0 10 726.5 5 1609.4 11 2501.1 18	5.6 13 19 6 100 6 63 6	15031.0 (14 ⁺) 15010.6 (14 ⁺) 14127.8 (14 ⁺) 13238.1 (13 ⁺)		Q	
15858.2	(15 ⁺)	847.6 6 1731.1 12	5 5 100 5	15010.6 (14 ⁺) 14127.8 (14 ⁺)		D+Q	
16167.2		1062.0 10 2283.0 16	1×10 ² 1 1×10 ² 1	15105.2 13884.2 (13 ⁺)			
16171.0	(15 ⁺)	847.0 10 1715.0 12	67 33 100 33	15324.1 (14 ⁺) 14455.8 (13 ⁺)		Q	
16246.6	(16 ⁻)	1312.0 9 2396.1 17	80 20 100 20	14934.7 (15 ⁻) 13850.1 (14 ⁻)		D+Q	
16496.6	(16 ⁻)	2279.0 19	100	14217.5 (14 ⁻)		Q	
16567.0	(16 ⁺)	708.6 10 857.6 6 1645.6 12	33 17 83 17 100 17	15858.2 (15 ⁺) 15709.3 (15 ⁺) 14920.9 (14 ⁺)		D+Q	
16676.4	(16 ⁺)	818.4 6 940.4 7 1644.6 12 1665.0 12 2546.0 18	87 13 75 13 50 13 100 13 50 13	15858.2 (15 ⁺) 15736.9 (15 ⁺) 15031.0 (14 ⁺) 15010.6 (14 ⁺) 14127.8 (14 ⁺)		D+Q	
16798.0	(15 ⁻)	1363.8 10 1385.4 10	40 20 20 20	15434.1 (13 ⁻) 15412.6 (13 ⁻)		Q	

Adopted Levels, Gammas (continued)

 $\gamma(^{58}\text{Ni})$ (continued)

29

E_i (level)	J_i^π	E_γ^\ddagger	I_γ^\ddagger	E_f	J_f^π	Mult. [†]	δ^\ddagger
16798.0	(15 ⁻)	1474 ^a		15324.1 (14 ⁺)			
		1503.9 11	100 20	15294.3 (14 ⁺)		D	
		1531.9 11	20 20	15266.3 (14 ⁺)			
		1556.0 10	40 20	15241.9 (13 ⁻)			
		3750		13048.2 (13 ⁻)			
		3965 3	20 20	12831.6 (13 ⁻)		Q	
17019.6		2166.5 15	100	14853.1 (15 ⁻)			
17163.1	(16 ⁺)	992.1 10	100 33	16171.0 (15 ⁺)		D+Q	
		1839.1 13	67 13	15324.1 (14 ⁺)		Q	
		1896.6 13	67 13	15266.3 (14 ⁺)			
17197		2092.0 20	100	15105.2			
17290.0	(16 ⁺)	1965.0 14	15 8	15324.1 (14 ⁺)		Q	
		1996.0 14	100 8	15294.3 (14 ⁺)		Q	
		2023.9 12	15 8	15266.3 (14 ⁺)			
		2436		14853.1 (15 ⁻)			
17530.0	(17 ⁺)	854.0 6	70 10	16676.4 (16 ⁺)			
		962.8 7	15 5	16567.0 (16 ⁺)			
		1793.3 13	100 10	15736.9 (15 ⁺)		Q	
		1819.5 13	60 10	15709.3 (15 ⁺)			
17681.4	(17 ⁺)	1004.8 7	50 10	16676.4 (16 ⁺)			
		1113.8 8	30 10	16567.0 (16 ⁺)			
		1823.7 13	100 10	15858.2 (15 ⁺)		Q	
		1944.8 14	70 10	15736.9 (15 ⁺)		Q	
		1972.7 14	40 10	15709.3 (15 ⁺)		Q	
18261.1	(17 ⁺)	1097.9 10	17 17	17163.1 (16 ⁺)		D+Q	
		2090.0 10	100 33	16171.0 (15 ⁺)		Q	
18341.5	(16 ⁻)	1583.0 11	17 17	16758 (14 ⁻)			
		1596.0 11	17 17	16745 (14 ⁻)			
		1634.0 11	83 17	16707 (14 ⁻)		Q	
		1668.0 12	100 17	16673 (14 ⁻)			
		3489.4 24	17 17	14853.1 (15 ⁻)	D(+Q)	-0.02 14	
18461.0	(17 ⁻)	1170.5 8	17 6	17290.0 (16 ⁺)	D+Q	-0.10 6	
		1664.0 12	100 6	16798.0 (15 ⁻)		Q	
18638.9	(18 ⁺)	957.5 7	75 8	17681.4 (17 ⁺)			
		1109.0 10	17 8	17530.0 (17 ⁺)			
		1962.2 14	100 8	16676.4 (16 ⁺)		Q	
		2073.0 15	33 8	16567.0 (16 ⁺)		Q	
19196		2949.0 30	100	16246.6 (16 ⁻)			
19205.4	(17 ⁻)	864.0 10	33 33	18341.5 (16 ⁻)	D+Q		
		1598.0 10	67 33	17607 (15 ⁻)			
		1623.6 11	100 33	17582 (15 ⁻)			
		1723.0 13	67 33	17482 (15 ⁻)			

Adopted Levels, Gammas (continued)

 $\gamma(^{58}\text{Ni})$ (continued)

E_i (level)	J_i^π	E_γ^\ddagger	I_γ^\ddagger	E_f	J_f^π	Mult. [†]	E_i (level)	J_i^π	E_γ^\ddagger	I_γ^\ddagger	E_f	J_f^π	Mult. [†]
19482.5	(18 ⁺)	1221.1 10	50 50	18261.1 (17 ⁺)			24611	(22 ⁻)	1280	2400.0 17	100 17	23331	(21 ⁻)
		2320.0 16	100 50	17163.1 (16 ⁺)	(Q)				2400.0 17	100 17	22211.3	(20 ⁻)	Q
19566.9	(18 ⁺)	2276.9 16	100	17290.0 (16 ⁺)	Q		25141	(22 ⁺)	3002.8 21	100	22138	(20 ⁺)	Q
19945.7	(19 ⁺)	1307.3 9	58 8	18638.9 (18 ⁺)			25552	(23 ⁻)	2750.5 19	100	22800.4	(21 ⁻)	Q
		2263.4 16	100 8	17681.4 (17 ⁺)	Q		25918	(23 ⁻)	2587.0 18	100	23331	(21 ⁻)	Q
		2415.0 17	42 8	17530.0 (17 ⁺)	Q		26059.7	(23 ⁺)	1848.0 13	50 50	24211.9	(22 ⁺)	
20135.4	(18 ⁻)	930.0 10	33 11	19205.4 (17 ⁻)					3291.0 23	100 50	22767.9	(21 ⁺)	Q
		1794.0 13	100 11	18341.5 (16 ⁻)	Q		27366	(24 ⁻)	2755.0 20	100	24611	(22 ⁻)	Q
20450.1	(19 ⁻)	1988.7 14	100	18461.0 (17 ⁻)	Q		28709	(25 ⁻)	3157.0 22	100	25552	(23 ⁻)	Q
20826.2	(19 ⁺)	2565.0 18	100	18261.1 (17 ⁺)	Q		28931	(25 ⁻)	3014.0 21	100	25918	(23 ⁻)	Q
21106.3	(19 ⁻)	971.0 10	14 14	20135.4 (18 ⁻)	D+Q				3379		25552	(23 ⁻)	
		1901.0 14	100 14	19205.4 (17 ⁻)	Q		30491	(26 ⁻)	3125		27366	(24 ⁻)	
21248.0	(20 ⁺)	1301.8 9	50 7	19945.7 (19 ⁺)	Q		32175	(27 ⁻)	3466		28709	(25 ⁻)	
		2609.0 18	100 14	18638.9 (18 ⁺)	Q		32495	(27 ⁻)	3564		28931	(25 ⁻)	
22138	(20 ⁺)	2570.9 18	100	19566.9 (18 ⁺)	Q				3786		28709	(25 ⁻)	
22211.3	(20 ⁻)	1105.0 10	22 11	21106.3 (19 ⁻)	D+Q		33972	(28 ⁻)	3480		30491	(26 ⁻)	
		2076.0 15	100 11	20135.4 (18 ⁻)	Q		36045	(29 ⁻)	3870		32175	(27 ⁻)	
22239.6	(20 ⁺)	2757.0 19	100	19482.5 (18 ⁺)	Q		36535	(29 ⁻)	4040		32495	(27 ⁻)	
22767.9	(21 ⁺)	1519.2 11	37 13	21248.0 (20 ⁺)			37810	(30 ⁻)	3838		33972	(28 ⁻)	
		2824.3 20	100 13	19945.7 (19 ⁺)	Q		40333	(31 ⁻)	4288		36045	(29 ⁻)	
22800.4	(21 ⁻)	2349.7 16	100	20450.1 (19 ⁻)	E2		40931	(31 ⁻)	4396		36535	(29 ⁻)	
23331	(21 ⁻)	1120		22211.3 (20 ⁻)			42007	(32 ⁻)	4197		37810	(30 ⁻)	
		2225.0 16	100	21106.3 (19 ⁻)	Q		2868.1+x		2868		x		
23741	(21 ⁺)	2914.5 25	100	20826.2 (19 ⁺)	Q		6083.2+x		3215		2868.1+x		
24211.9	(22 ⁺)	1444.0 10	29 14	22767.9 (21 ⁺)			9667.3+x		3584		6083.2+x		
		2964.0 19	100 14	21248.0 (20 ⁺)	Q								

[†] Mainly from $\gamma(\theta)$ in (p,p'γ). Some assignments are from DCO values in (²⁴Mg, α 2pγ). The multipolarity assignments for γ rays from (γ , γ') are from polarization asymmetry measurements.

[‡] Values represent averages of all available data. For γ rays taken from (γ , γ') work only, values are level-energy differences, without applying any correction for recoil, which is at most 1 keV.

[#] Total theoretical internal conversion coefficients, calculated using the BrIcc code (2008Ki07) with “Frozen Orbitals” approximation based on γ-ray energies, assigned multipolarities, and mixing ratios, unless otherwise specified.


[@] Multiply placed.

[&] Multiply placed with intensity suitably divided.

^a Placement of transition in the level scheme is uncertain.


Adopted Levels, GammasLevel Scheme

Intensities: Relative photon branching from each level



Adopted Levels, GammasLevel Scheme (continued)

Intensities: Relative photon branching from each level




Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level


- - - - - → γ Decay (Uncertain)

Adopted Levels, Gammas

Legend


Level Scheme (continued)

Intensities: Relative photon branching from each level

- - - - - ► γ Decay (Uncertain)

Adopted Levels, GammasLevel Scheme (continued)


Intensities: Relative photon branching from each level



Adopted Levels, Gammas

Level Scheme (continued)


Intensities: Relative photon branching from each level
 @ Multiply placed: intensity suitably divided



Adopted Levels, GammasLevel Scheme (continued)


Intensities: Relative photon branching from each level

@ Multiply placed: intensity suitably divided



Adopted Levels, Gammas**Level Scheme (continued)**


Intensities: Relative photon branching from each level
 @ Multiply placed: intensity suitably divided



Adopted Levels, Gammas**Level Scheme (continued)**

Intensities: Relative photon branching from each level

@ Multiply placed: intensity suitably divided



Adopted Levels, Gammas


Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level

@ Multiply placed: intensity suitably divided

-----► γ Decay (Uncertain)



Adopted Levels, Gammas


Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level

@ Multiply placed: intensity suitably divided

→ γ Decay (Uncertain)




Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level

@ Multiply placed: intensity suitably divided

- - - - - γ Decay (Uncertain)


Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level

@ Multiply placed: intensity suitably divided

- - - - - γ Decay (Uncertain)


Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level

@ Multiply placed: intensity suitably divided

- - - - - γ Decay (Uncertain)


Adopted Levels, Gammas


Legend


Level Scheme (continued)

Intensities: Relative photon branching from each level

@ Multiply placed: intensity suitably divided


- - - - - γ Decay (Uncertain)





Adopted Levels, Gammas

Adopted Levels, Gammas (continued)

Band(F): $\Delta J=1$ band based on 10694,
 10^-



Band(H): $\Delta J=1$ band based on 7724,
 8^+



Adopted Levels, Gammas (continued)