Adopted Levels, Gammas

	History									
	Туре	Author	Citation	Literature Cutoff Date						
Fu	Ill Evaluation	E. Browne, J. K. Tuli	NDS 114, 1849 (2013)	31-Dec-2012						
$Q(\beta^-)=237 \ 3; \ S(n)=8820 \ 4;$ Additional information 1. Others: Nuclear structure. 2012Lo04, 2012Mu09, 20111 2004Mi54, 2004Pa04, 20 Nuclear reactions: 2011Su04 Effect of strong magnetic fie Compilations. B(E2) values: 2012Go17, 20 Half-lives: 2011Ch65. Discovery of element iron: 2	S(p)=13176 4; Ba39, 2011Ut0 002Ca48, 1985I , 2006Sc16, 20 Id on ⁶⁰ Co ε do 12Pr08, 2011Pr 5007Li49.	$Q(\alpha) = -8556 \ 4$ 2012 1, 2009Su20, 2007Al45, 3111. 03Kn01. ecay: 2007Li49.	Wa38 2007Mo15, 2005Al47, 20	005Ch12, 2005Pu04, 2004Ag02,						

	Arg	guments for J^{2}	π assignments				
E(level)	L(t,p)#	$\gamma(\theta)$ in (t,p γ)	$\gamma(heta)$ in (⁴⁸ Ca,2n γ)	L(α,	² He)#	(⁶⁴ Ni,Xγ)	Adopt
0	0				0+	0+	
824	2	2			2+	2+	
1974	0					0 ⁺	
2115	4	2,4			4+	4^{+}	
2300	2	2			2+	a	
2356	0					0+	
2673	2	1,2,3				2+	
2756	2					2+	
2793					4^{+}	4+	
3039	2	1,2,3				2+	
3072	4	2,4		4		4+	
3293	3					3-	
3308		1,2,3,4				<	
3499	4				<	(4 ⁺)	
3516				4,5	(5 ⁺)	(5 ⁺)	
3520			(6')		(6')	(6')	
3520	(2)		(4')		(4')	(4')	
3562	(3)		((+))			(3)	
3582	2		(6')			(6^{+})	
3035	2					2 · 0 +	
3098	U C					0 ⁻	
2004	2		(6+)			5 (6 ⁺)	
2020	r		(0)			2+	
2022	2		6+			2 6 ⁺	
3058			6 ⁽⁻⁾			6 ⁽⁻⁾	
3950			(7^+)		(7^{+})	(7^+)	
4053	3		(r)		(7)	3-	
4176	2					2 ⁺	
4280	3					3-	
4296	5		7(-)			7(-)	
4358			7 ⁽⁻⁾	7		7 ⁽⁻⁾	
4359	5		-	-		5-	
4440	3					3-	
4451	-		6+			6+	
4503	4		-			4+	
4650	2					2+	

4755	(3))				(3-)		
4958	4					4^{+}		
5006			8(-)			8(-)		
5029	4					4^{+}		
5103	2					2+		
5218	3					3-		
5310	(5)			5.7	(5^{-})		
5333	(5)	, ,	8 (+)			8(+)		
5529			Q (-)			Q (-)		
5550			2 8+			2 8+		
5550	(7	N N	0		7 5	(7^{-})		
5020	0.)	0-		7,5	(7)		
5/55			9			9		
6475			10			10,		
6550			10()			10()		
6620	(8,	5)			8+6	(8+,6+)		
6740			(9,10)			(9,10)		
7250			$11^{(-)}$			11(-)		
7632			$11^{(-)}$			$11^{(-)}$		
7890			11			11		
8059			12^{+}			12^{+}		
8536			$12^{(-)}$			$12^{(-)}$		
9503			(13 ⁻)			(13 ⁻)		
9996			14^{+}			14^{+}		
10721			(14^{-})			(14^{-})		
11810			15			15		
12116			(15^{-})			(15^{-})		
12319			(16^+)			(16^+)		
12833			(16^{-})			(16^{-})		
14583			(10^{-})			(17^{-})		
14905			(17)			(18^+)		
17056			(10)			(10)		
# J^{π} o a $J^{\pi}=2^+$	f ⁵⁸ Fe not	(g.s.) is 0 ⁺ . consistent with 1	log <i>ft</i> in	ι <i>β</i> - α	lecay. ⁶⁰ Fe Levels			
			(Cross F	Reference (XREF)	Flags		
		60			50			64 . 2 7
		A $^{00}Mn \beta^{-} d$	ecay (1.77 s)	E	50 Fe(t,p γ)		I	⁰⁴ Ni(³ He, ⁷ Be)
		B 60 Mn β^- d	ecay (0.28 s)	F	64 Ni(d, ⁶ Li)		J	62 Ni(14 C, 16 O)
		$C = {}^{14}C({}^{48}Ca,2)$	nγ)	G	$^{48}Ca(^{15}N,2np\gamma),$	$(^{18}\text{O},2\text{n}\alpha\gamma)$	K	208 Pb(64 Ni,X γ)
		D = 58Fe(t,p)		Н	58 Fe(α , ² He)		L	⁶⁴ Ni(²³⁸ U,X)
E(level) [†]	Jπ‡	$T_{1/2}^{\#}$	XREF			Co	mmen	its
0.00	0			<i>a</i> . 0-	100			
0.04	0^{+}	$2.62 \times 10^{\circ} \text{ y } 4$ ABC	DEFGHIJKL	%β ⁻ =	=100	a		
				$T_{1/2}$:	From 2009Ru08.	Specific activ	vity m	easurement. Measured activity
				of	²⁰ Fe in the source	, its isotopic	compo	osition, and the number of iron
				ato	ms in the source.			
				T _{1/2} :	Other values: 1.4	9×10 ⁶ y 27, :	specifi	c activity measurement and
				rad	ioisotope concenti	ation (1984K	(u28).	3×10 ⁵ y (1957Ro54).
				$T_{1/2}$:	a larger sample n	naterial and a	more	accurate determination of the
				nur	nber of atoms sug	gests the resu	ult in 2	2009Ru08 is the most accurate.
				T1/2	The half-life of 6	Fe plays a p	romin	ent role in various
				- 1/2· asti	ophysical matters		- 0.11111	
873 83 <mark>0</mark> 0	2+	79 ns 8 ADC	DEEC TIME	Tue	From 20101 :01 :	n 64 Ni (238 I T	X) ∩+	her value: 8.0 ps 15
023.03 7	~	··> ps o ADC	PELO IJKE	+1/2· (10	77Wa10)		1. OL	ner value. 0.0 ps 15
1974.0 5	0^{+}	R	DE	(1)				
	-	_						

Continued on next page (footnotes at end of table)

⁶⁰Fe Levels (continued)

E(level) [†]	$J^{\pi \ddagger}$	$T_{1/2}^{\#}$	XREF	Comments
2114.60 ^a 12	4+	0.83 ps 21	A CDEFG I K	
2299.67 11	·	0.00 p5 21	A DE K	$J^{\pi}=2^+$ from 1476 γ M1 to 2 ⁺ , and 2300 γ E2 to 0 ⁺ . However, log <i>ft</i> =5.0 to 2300-keV level is not consistent with β^- decay from 60 Mn(4 ⁺) to 60 Ee(2 ⁺).
2356.2 10	0^{+}		BDF	
2672.9 <mark>&</mark> 9	2^{+}		DE	
$2755.9^{\&}10$	- 2+		DF	
2792 68 11	$\frac{2}{3^{+}}4^{+}$			I^{π} . $I^{\pi}-3^{+}$ from (M1) γ rays to $I^{\pi}-2^{+}$ A^{+} levels in ${}^{60}\text{Fe}$ $I^{\pi}-4^{+}$
2792.00 11	J , 1		n k	$from^{208} Pb(^{64}Ni, X\gamma).$
3038.9° 10	2+		DEF	
3072.01 23	4+		A DE HI	
3193.51 24	2-		A	
32934	3		D	
3307.9 9			EF	
3332.90			A	
3400.02 24	(4^{+})		A D	
251615b18	()	40 mg 21		VDEE: $H(2520)$
5510.15 16	(5)	49 ps 21	C ON K	$T_{1/2}$: assignment to 3516 level is uncertain.
3520 [@] 50	(4^{+})		Н	
3520.12 ^{<i>a</i>} 22	6+		C K	
3562 5	(3-)		D	
3582.21 18	(6^+)		СК	
3635 4	2*		D F	
3647.9 [∞] 9	o.+		E	
3698 5	0^{+}		D	
3713.9 × 10			E	
3867 5	3-		D	
3874.9 ^{&} 9			E	
3904.5 <i>3</i>	(6+)		C	
3929.9 10	2+		DE	
3931.8/18	6'	. 0.4	C	
3958.20° 18	(7^+)	>0.4 ps	CG	
5959.15 25 4053 8	$(7)_{3^{-}}$			
4055 8	2+		ם	
4280 8	3-		D	
4296.49 18	7(-)	>0.4 ps	C G	
4298.2 4		1	K	
4358.30 <mark>b</mark> 18	$7^{(-)}$		СН	
4359.5 3	5-		D G K	
4440 10	3-		D	
4451.4 <i>3</i>	6+		С	
4503 10	4+		D	
4650 10	2+		D	
4755 9	(3^{-})		D	
4958 9	4'	0.0 12 1	D	
5006.08° 19	8(-) 4+	0.8 ps $+13-4$	CG	
5029 10 5103 10	4 2+		ע	
5218 16	<u>2</u> 3-		DF	
~5310@	(5^{-})		. п	
~5510	(\mathbf{J})		п	

⁶⁰Fe Levels (continued)

E(level) [†]	$J^{\pi \ddagger}$	XRE	EF	Comments
5333.39 ^a 19	8+	С		
5434 17		D		
5528.75 ^b 20	9(-)	С		
5549.6 5	8+	С	K	
5596 18		D		
≈5620 [@]	(7-)		н	
5754.62 19	9-	С		
6314.7 5		С		
6475.27 ^a 21	10+	С		
6550.10 [°] 21	$10^{(-)}$	С		
6578.62 22		С		
≈6620 [@]	$(8^+, 6^+)$		H	J^{π} : configuration= $(\nu g_{9/2})^2 8^+$ or $((\nu g_{9/2})(\nu d_{5/2}))6^+$.
6740.1 ^d 4	(9,10)	С		
6944.4 5		С		
7003.4 6		С		
7243.13 24		С		
7250.07 ^b 23	$11^{(-)}$	С		
7631.9 4	$11^{(-)}$	С		
7664.9 <i>3</i>		С		
7890.41 ^d 22	11	С		
8059.49 ^a 23	12+	С		
8536.5 [°] 3	$12^{(-)}$	С		
8920.2 5		С		
8974.4 7		С		
9503.2 ^b 4	(13 ⁻)	С		
9559.43 ^d 25	13	С		
9995.83 ^a 25	14^{+}	С		
10670.4 13		С		
10721.0 ^C 8	(14 ⁻)	С		
11810.5 ^d 6	15	С		
12116.2 ^b 11	(15^{-})	С		
12319.0 ^{<i>a</i>} 16	(16^{+})	С		
12833.1 ^c 17	(16 ⁻)	С		
14583.4 ^b 15	(17 ⁻)	С		
14984.6 ^{<i>a</i>} 17	(18^{+})	С		
17956 ^a 4	(20^{+})	С		

[†] Levels connected by γ 's are from least-squares fit to E γ , others are from (t,p), unless stated otherwise. [‡] See separate table. See ⁵⁸Fe(t,p) and ⁵⁸Fe(α ,²He) references for level configurations used in DWBA calculations. [#] From ⁴⁸Ca(¹⁵N,2np γ),(¹⁸O,2n $\alpha\gamma$); recoil distance method for levels below 3.6 MeV and DSA method for levels above this energy, except the g.s. ^(a) From ${}^{58}\text{Fe}(\alpha,{}^{2}\text{He})$. [&] From ${}^{58}\text{Fe}(t,p\gamma)$ (1977Wa10). Kept fixed in least-squares fit.

^{*a*} Band(A): Yrast band.

^b Band(B): Band based on $J^{\pi}=5^{(-)}$.

^{*c*} Band(b): Band based on $J^{\pi} = 6^{(-)}$.

^d Band(C): Band based on $J^{\pi} = (9, 10)$.

$\gamma(^{60}\text{Fe})$

E _i (level)	\mathbf{J}_i^{π}	Eγ	I_{γ}^{b}	E_f	\mathbf{J}_f^{π}	Mult.	α^{\dagger}	Comments
823.83	2+	823.8 1	100	0.0	0+	[E2]	0.000319 5	B(E2)(W.u.)=13.6 14 α =0.000319 5; α (K)=0.000287 4; α (L)=2.75×10 ⁻⁵ 4; α (M)=3.78×10 ⁻⁶ 6; α (N+)=1.740×10 ⁻⁷ 25 α (N)=1.740×10 ⁻⁷ 25
1974.0 2114.60	0' 4 ⁺	1150.2 <i>4</i> 1290.8 <i>1</i>	100	823.83 823.83	2+ 2+	[E2]	0.0001356 <i>19</i>	B(E2)(W.u.)=14 4 α =0.0001356 19; α (K)=9.84×10 ⁻⁵ 14; α (L)=9.36×10 ⁻⁶ 14; α (M)=1.289×10 ⁻⁶ 18; α (N+)=2.65×10 ⁻⁵ α (N)=5.98×10 ⁻⁸ 9; α (IPF)=2.65×10 ⁻⁵ 4
2299.67		1475.8 ^{&} 1	80 ^{&} 4	823.83	2+	(M1)	0.0001331 19	$ \begin{array}{l} \alpha = 0.0001331 \ 19; \ \alpha(\mathrm{K}) = 6.70 \times 10^{-5} \ 10; \\ \alpha(\mathrm{L}) = 6.35 \times 10^{-6} \ 9; \ \alpha(\mathrm{M}) = 8.74 \times 10^{-7} \\ 13; \ \alpha(\mathrm{N}+) = 5.89 \times 10^{-5} \ 9 \\ \alpha(\mathrm{N}) = 4.08 \times 10^{-8} \ 6; \ \alpha(\mathrm{IPF}) = 5.88 \times 10^{-5} \\ 9 \end{array} $
		2299.7 ^{&} 2	100 ^{&} 7	0.0	0^{+}			Mult.: (E2) from ΔJ^{π} in 2010Ho13 not consistent with β^{-} decay.
2356.2	0^{+}	1532.4	100	823.83	2+			
2672.9	2+	1849 [@]	100	823.83	2^{+}			
2755.9	2+	1932 [@]	100 [@]	823.83	2+			
2792.68	3+,4+	493.0 ^{&} 1	36.2 ^{&} 19	2299.67		(M1)	0.000697 10	$ \begin{array}{l} \alpha = 0.000697 \ 10; \ \alpha(\mathrm{K}) = 0.000628 \ 9; \\ \alpha(\mathrm{L}) = 6.02 \times 10^{-5} \ 9; \ \alpha(\mathrm{M}) = 8.29 \times 10^{-6} \\ 12; \ \alpha(\mathrm{N}+) = 3.84 \times 10^{-7} \ 6 \\ \alpha(\mathrm{N}) = 3.84 \times 10^{-7} \ 6 \end{array} $
		678.1 ^{&} 1	5.1 ^{&} 6	2114.60	4+	(M1)	0.000348 5	$ \begin{array}{l} \alpha = 0.000348 \ 5; \ \alpha(\mathrm{K}) = 0.000314 \ 5; \\ \alpha(\mathrm{L}) = 3.00 \times 10^{-5} \ 5; \ \alpha(\mathrm{M}) = 4.13 \times 10^{-6} \\ 6; \ \alpha(\mathrm{N}+) = 1.92 \times 10^{-7} \ 3 \\ \alpha(\mathrm{N}) = 1.92 \times 10^{-7} \ 3 \end{array} $
		1968.8 ^{&} 1	100 ^{&} 5	823.83	2+	(M1)	0.000285 4	$\alpha = 0.000285 \ 4; \ \alpha(K) = 4.00 \times 10^{-5} \ 6; \\ \alpha(L) = 3.78 \times 10^{-6} \ 6; \ \alpha(M) = 5.21 \times 10^{-7} \\ 8; \ \alpha(N+) = 0.000241 \ 4 \\ \alpha(N) = 2.43 \times 10^{-8} \ 4; \ \alpha(IPF) = 0.000241 \ 4$
3038.9	2+	2215 [@]	$100^{@}$	823.83	2+			
3072.01	4+	279.6 ^{&} 7	11 ^{&} 7	2792.68	3+,4+	M1	0.00260 4	$\alpha = 0.00260 \ 4; \ \alpha(K) = 0.00234 \ 4; \\ \alpha(L) = 0.000226 \ 4; \ \alpha(M) = 3.11 \times 10^{-5} \\ 5; \ \alpha(N+) = 1.431 \times 10^{-6} \ 22 \\ \alpha(N) = 1.431 \times 10^{-6} \ 22$
		957.5 ^{&} 3	48 ^{&} 7	2114.60	4+	M1	0.0001708 24	$\alpha = 0.0001708 \ 24; \ \alpha(K) = 0.0001540 \ 22; \\ \alpha(L) = 1.464 \times 10^{-5} \ 21; \\ \alpha(M) = 2.02 \times 10^{-6} \ 3 \\ \alpha(N) = 9.39 \times 10^{-8} \ 14$
		2248.0 ^{&} 3	100 ^{&} 22	823.83	2+	E2	0.000471 7	α =0.000471 7; α (K)=3.35×10 ⁻⁵ 5; α (L)=3.17×10 ⁻⁶ 5; α (M)=4.36×10 ⁻⁷ 7; α (N+)=0.000434 6 α (N)=2.03×10 ⁻⁸ 3; α (IPF)=0.000434 6
3193.51		401.0 ^c 10 1078.9 2	77 100 <i>14</i>	2792.68 2114.60	3 ⁺ ,4 ⁺ 4 ⁺			

Continued on next page (footnotes at end of table)

$\gamma(^{60}\text{Fe})$ (continued)

E _i (level)	\mathbf{J}_i^{π}	Eγ	I_{γ}^{b}	E_f	\mathbf{J}_f^{π}	Mult.	$lpha^{\dagger}$	Comments
3307.9		635 [@]	100 [@] 20	2672.9	2+			
2252.0		2484 ^w	100 ^w 20	823.83	2+			
3352.9		1238.3 5	100	2114.60	4'			
3486.02	$(4\pm)$	$13/1.4^{\circ}$ 2	100%	2114.60	4 · 4+	MI	0.0001201.17	~ -0.0001201 17. $\sim (K) -7.55 \times 10^{-5}$ 11.
3498.0?	(4*)	1384.000 10	100-	2114.00	4	IVII	0.0001201 17	$\begin{array}{l} \alpha = 0.0001201 \ T7, \ \alpha(\text{K}) = 7.33 \times 10^{-5} \ T1, \\ \alpha(\text{L}) = 7.15 \times 10^{-6} \ I0; \ \alpha(\text{M}) = 9.86 \times 10^{-7} \\ I4; \ \alpha(\text{N}+) = 3.65 \times 10^{-5} \ 6 \\ \alpha(\text{N}) = 4.60 \times 10^{-8} \ 7; \ \alpha(\text{IPF}) = 3.64 \times 10^{-5} \ 6 \end{array}$
3516.15	(5 ⁻)	1401.56 [#] <i>19</i>	100	2114.60	4+	[E1]	0.000233 4	B(E1)(W.u.)= $0.32 \times 10^{-5} 14$ α =0.000233 4; α (K)= $4.17 \times 10^{-5} 6$; α (L)= $3.95 \times 10^{-6} 6$; α (M)= 5.44×10^{-7} 8; α (N+)= $0.000187 3$ α (N)= $2.53 \times 10^{-8} 4$; α (IPF)= $0.000187 3$
3520.12	6+	1405.4 [‡] 3	100	2114.60	4+			
3582.21	(6+)	1467.4 [‡] 3	100	2114.60	4+			
3647.9		2824 [@]	100	823.83	2^{+}			
3713.9		2890 [@]	100	823.83	2+			
3874.9		227 [@]		3647.9				
		3051 [@]		823.83	2+			
3904.5	(6+)	1789.1 ^{<i>a</i>} 7	100 ^{<i>a</i>}	2114.60	4+			
3929.9	2+	3106	100	823.83	2+			
3931.87	6+	349.5 ^{<i>a</i>} 1 1817.4 ^{<i>a</i>} 2	15.6 ^{<i>a</i>} 16 100 ^{<i>a</i>} 5	3582.21 2114.60	(6 ⁺) 4 ⁺			
3958.20	6(-)	375.9 ^a 1	27 <mark>a</mark> 1	3582.21	(6+)			
		437.9 ^{<i>a</i>} 3	100 ^{<i>a</i>} 3	3520.12	6+			
		441.9^{a} I	$47^{a} 2$	3516.15	$(5)_{4^+}$			
2050 12	(7^{+})	$377.0^{\ddagger}.2$	1.5 /	2592.21	(6^+)			
3939.13	(7)	420.0 [±] 2		2520.12	(0)			
		439.0^{\ddagger} 3		2516.15	(5-)			
1206 10	7(-)	$442.9^{\circ} 5$ 338.2 ^{<i>a</i>} 1	1000 1	3058 20	(3)			
7270.77	,	364.5^{a} 1	37^{a} 1	3931.87	6+			
		714.4 ^a 1	53 ^a 2	3582.21	(6 ⁺)			
		780.6 ^a 10	99 <mark>a</mark> 4	3516.15	(5 ⁻)			
4298.2		339.1 [‡] <i>3</i>	100	3959.13	(7^{+})			
4358.30	$7^{(-)}$	399.9 ^{<i>a</i>} 1	5.7 ^a 6	3958.20	6(-)			
		426.4^{a} 4	$1.1^{a} 6$	3931.87	6^+			
		455.7^{a} 2 842 3 ^a 1	$2.9^{a} 0$	3904.5	(0^{-})			
1350 5	5-	842.3 I 843.3 <mark>#</mark> .2	100 4	3516.15	(5^{-})			
4451.4	5 6 ⁺	2336.9^{a} 20	100^{a}	2114.60	(3^{+})			
5006.08	8(-)	647.7 ^{<i>a</i>} 2	4.7 ^{<i>a</i>} 6	4358.30	7 ⁽⁻⁾			
		1047.9 ^{<i>a</i>} 1	100 ^{<i>a</i>} 3	3958.20	6 ⁽⁻⁾	[E2]	0.0001749 25	B(E2)(W.u.)=40.1 α =0.0001749 25; α (K)=0.0001577 22; α (L)=1.503×10 ⁻⁵ 21; α (M)=2.07×10 ⁻⁶ 3
		~	a					$\alpha(N)=9.57\times10^{-8}$ 14
5333.39	8+	882.0 ^{<i>a</i>} 2 1036.8 ^{<i>a</i>} 1	22^{a} 1 100^{a} 3	4451.4 4296.49	6+ 7 ⁽⁻⁾			

Continued on next page (footnotes at end of table)

$\gamma(^{60}\text{Fe})$ (continued)

E _i (level)	\mathbf{J}_i^{π}	E_{γ}	I_{γ}^{b}	E_f	\mathbf{J}_f^π
5333.39	8+	1751.6 ^a 2	58 ^a 2	3582.21	(6 ⁺)
		1813.4 ^a 4	21 ^{<i>a</i>} 1	3520.12	6+
5528.75	9(-)	522.8 ^a 2	4.6 ^a 7	5006.08	8(-)
		1170.4 ^a 1	100 ^a 3	4358.30	$7^{(-)}$
		1232.3 ^a 6	4.6 ^a 7	4296.49	$7^{(-)}$
5549.6	8^{+}	1967.4 ^a 4	100 ^{<i>a</i>}	3582.21	(6^{+})
5754.62	9-	748.5 ^a 1	18 ^a 1	5006.08	8(-)
		1458.1 ^{<i>a</i>} 1	100 ^a 3	4296.49	$7^{(-)}$
6314.7		765.1 ^a 2	100 ^a	5549.6	8+
6475.27	10^{+}	720.3 ^a 3	12.8 ^a 6	5754.62	9-
		1141.9 ^a 1	100 ^{<i>a</i>} 3	5333.39	8+
6550.10	$10^{(-)}$	1021.2 ^a 15	1.9 ^a 9	5528.75	9(-)
		1544.0 ^a 1	100 ^a 3	5006.08	8(-)
6578.62		824.0 ^a 1	100 ^{<i>a</i>}	5754.62	9-
6740.1	(9, 10)	1734.0 ^a 5	100 ^a	5006.08	8(-)
6944.4		1415.6 ^a 4	100 ^a	5528.75	9(-)
7003.4		688.7 ^a 3	100 ^{<i>a</i>}	6314.7	
7243.13		664.5 ^a 1	100 ^{<i>a</i>}	6578.62	
7250.07	$11^{(-)}$	1721.3 ^a 1	100 ^{<i>a</i>}	5528.75	9(-)
7631.9	$11^{(-)}$	1877.2 ^a 3	100 ^{<i>a</i>}	5754.62	9-
7664.9		421.8 ^{<i>a</i>} 1	100 a	7243.13	
7890.41	11	1150.3 ^a 3	45 ^a 5	6740.1	(9,10)
		1340.3 ^a 1	100 ^{<i>a</i>} 5	6550.10	$10^{(-)}$
		1415.1 ^a 10	75 ^a 5	6475.27	10^{+}
8059.49	12^{+}	1584.2 ^a 1	100 ^a	6475.27	10^{+}
8536.5	$12^{(-)}$	1986.4 ^a 2	100 ^{<i>a</i>}	6550.10	$10^{(-)}$
8920.2		1255.3 ^a 4	100 ^a	7664.9	
8974.4		2030.0 ^a 5	100 ^a	6944.4	
9503.2	(13^{-})	2253.1 ^a 3	100 ^{<i>a</i>}	7250.07	$11^{(-)}$
9559.43	13	1499.9 ^a 2	51 ^a 3	8059.49	12^{+}
		1669.0 ^a 2	100 ^a 3	7890.41	11
9995.83	14^{+}	1936.3 ^a 1	100 ^{<i>a</i>}	8059.49	12^{+}
10670.4		1696.0 ^a 11	100 ^{<i>a</i>}	8974.4	
10721.0	(14 ⁻)	2184.4 ^a 7	100 ^a	8536.5	$12^{(-)}$
11810.5	15	2251.0 ^a 5	100 ^a	9559.43	13
12116.2	(15^{-})	2612.9 ^a 10	100 ^{<i>a</i>}	9503.2	(13 ⁻)
12319.0	(16^{+})	2323.1 ^a 15	100 ^{<i>a</i>}	9995.83	14^{+}
12833.1	(16 ⁻)	2112.1 ^{<i>a</i>} 15	100 ^a	10721.0	(14 ⁻)
14583.4	(17-)	2467.2 ^a 10	100 ^a	12116.2	(15^{-})
14984.6	(18^{+})	2665.6^{a} 7	100 ^a	12319.0	(16^{+})
17956	(20^{+})	2971 ^a 3	100 ^{<i>a</i>}	14984.6	(18^{+})

[†] Additional information 2.
[‡] From (⁶⁴Ni,Xγ).
[#] From ⁴⁸Ca(¹⁵N,2npγ).
[@] From ⁵⁸Fe(t,pγ). Energy from level separation, not included in energy fit.
[&] From ⁶⁰Mn β-decay (1.77 s).
^a From ¹⁴C(⁴⁸Ca,2nγ) (2007De56).
^b Relative photon branching from each level.
^c Placement of transition in the level scheme is uncertain.

Adopted Levels, Gammas Legend Level Scheme $\begin{array}{l} I_{\gamma} < \ 2\% \times I_{\gamma}^{max} \\ I_{\gamma} < 10\% \times I_{\gamma}^{max} \\ I_{\gamma} > 10\% \times I_{\gamma}^{max} \end{array}$ ٠ Intensities: Type not specified 1 2971 100 (20+) 17956 + 2003.6 100 8 (18^{+}) - 246) - 2 14984.6 (17^{-}) 14583.4 + 2113, 100 907 (16^{-}) 1 - 26/- 200 12833.1 + ²³²³ | (16^+) S. 12319.0 23/10 (15^{-}) 12116.2 15 11810.5 | 5/84.400 | 001 0:00¹ (14^{-}) 10721.0 + 193,553 + 10670.4 + 100 + 100 + 126.00 12 Ş 9995.83 14^{+} <u>ج</u> 9559.43 $\frac{13}{(13^{-})}$ *%* -0; - 50301 9503.2 . ري ري 8974.4 8 8920.2 8 150 4 4. 1564,2 ŝ 5 g $12^{(-)}$ 8536.5 1512, - 0°. ŝ 12^{+} 8059.49 8 11 S. 7890.41 7664.9 S $\frac{11^{(-)}}{11^{(-)}}$ 7631.9 E 8 7250.07 7243.13 8 7003.4 8 0 6944.4 8 \sim (9,10) Ś 6740.1 9 S 6578.62 2 10(-) 6550.10 ć. 10^{+} 6475.27 6314.7 $\frac{9^{-}}{8^{+}}$ 5754.62 5549.6 5528.75 8^+ 5333.39 8(-) 0.8 ps +13-4 5006.08 0^+ 0.0 2.62 $\times 10^6$ y 4

⁶⁰₂₆Fe₃₄

Adopted Levels, Gammas







 ${}^{60}_{26}\text{Fe}_{34}$

Adopted Levels, Gammas



⁶⁰₂₆Fe₃₄