

Adopted Levels, Gammas

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	E. Browne, J. K. Tuli		NDS 111,1093 (2010)	3-Mar-2009

Q(β^-)=-5175.3; S(n)=11058.6 10; S(p)=8924.6 10; Q(α)=-4578.1 8 [2012Wa38](#)
 Note: Current evaluation has used the following Q record -5175.3 11058.9 108924.5 10-4578.1 8 [2009AuZZ,2003Au03](#).
[Additional information 1.](#)

⁶⁶Zn Levels

Configuration: configurations used in DWBA analysis of (α ,²He), (¹²C,¹⁰C), and (¹²C,¹⁰Be) data.

Cross Reference (XREF) Flags

A ⁶⁶ Cu β^- decay	K ⁶⁹ Ga(p, α)	U Coulomb excitation
B ⁶⁶ Ga ϵ decay	L ⁶⁵ Cu(d,n)	V ⁶⁷ Zn(d,t)
C ⁶⁴ Ni(α ,2n γ), ⁵⁵ Mn(¹⁴ N,2pn γ)	M ⁶⁷ Zn(p,d)	W ⁶² Ni(⁶ Li,d)
D ⁶⁶ Zn(p,p')	N ⁶⁸ Zn(p,t)	X ⁶⁴ Ni(¹⁶ O, ¹⁴ C)
E ⁶⁶ Zn(γ , γ')	O ⁶⁴ Zn(t,p)	Y ⁶⁴ Ni(³ He,n)
F ⁶⁶ Zn(p,p' γ)	P ⁶⁵ Cu(³ He,d)	Z ⁶⁵ Cu(α ,t)
G ⁶⁵ Cu(p, γ)	Q ⁶⁶ Zn(α , α')	Others:
H ⁶⁶ Zn(n,n' γ)	R inelastic scattering	AA ⁶⁴ Zn(α , ² He)
I ⁶⁵ Cu(p,n), (p,n γ) IAR	S ⁶⁶ Zn(e,e')	AB ⁶⁴ Ni(¹² C, ¹⁰ Be), ⁶⁴ Zn(¹² C, ¹⁰ C)
J ⁶³ Cu(α ,p), (α ,p γ)	T ⁶⁶ Zn(α , α' γ), (³² S, ³² S' γ)	AC ⁶³ Cu(⁶ He,p2n γ)

E(level) [†]	J π [#]	T _{1/2} [‡]	XREF	Comments
0.0	0 ⁺	stable	ABCDEFGHIJKLMN OPQRSTUVWXYZ	XREF: Others: AA , AB , AC Configuration=(ν f _{5/2} 0 ⁺)
1039.2279 21	2 ⁺ [@]	1.68 ps 3	BCDEFGH JKLMN OPQRSTU VWX Z	XREF: Others: AA , AB , AC μ =+0.80 8; μ =+0.9 2 (2005St24); Q=+0.24 8 (2003Ko51) μ =+0.80 8, transient field integral perturbed angular correlation (tf) (2002Ke02). μ =+1.06 10, transient field integral perturbed angular correlation (tf) (2004An14). μ =+0.9 2, perturbed angular correlation after ion implantation (IMPAC) (1979Fa06). Q: from Coulomb excitation (2003Ko51). Other: +0.24 9 (2020Ro06 , from Coul. ex., preliminary value as stated by authors). Comment added March 30, 2021 by B. Singh. T _{1/2} : weighted average of 1.73 ps 7 DSAM (2006Le24), 1.68 ps 3 DSAM (2002Ke02), 1.61 ps 10 Coul ex. (2003Ko51), 1.74 ps 11 ⁶⁶ Zn(γ , γ') (1998Ba02 , 1981Ca10 , 1972Ka22 , 1972ArZD , 1967Be39), 1.56 ps 10 Coul ex. (1973Fi15), 1.66 ps 10 ⁶⁶ Zn(e,e') (1977Ne05). Other measurements: 1.3 ps 8 ⁶³ Cu(α ,p γ) (1974Iv01), 1.5 ps 7 ⁶⁶ Zn(α , α' γ) (1972Yo01), 1.7 ps +35-14 ⁶⁴ Ni(α ,2n γ), <2 ns ⁶⁶ Cu β^- decay (1953En06). Isotope shift: <r ² > ^{1/2} =3.9496 13 (2004An14). XREF: Others: AB , AC T _{1/2} : others: <1.4 ps from (α , α' γ), and 0.83 ps +35-21 from ⁶⁴ Ni(α ,2n γ), 1.7 ps 5 (2003Ko51).
1872.7653 24	2 ⁺	0.19 ^a ps 7	ABCDEFGHIGH JKLM OPQRSTU VWX	

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued)

⁶⁶Zn Levels (continued)

E(level) [†]	J ^π #	T _{1/2} [‡]	XREF	Comments
2372.353 4	0 ⁺	>0.21 ps	AB DEFGH K1 OP	T _{1/2} : from (n,n'γ); other: <60 ps from (p,p'γ).
2451.01 5	4 ⁺ @	0.34 ps 5	CD FGH JKLM OPQR TUVW	XREF: Others: AB, AC J ^π : 4 from γ(θ) in ⁶⁴ Ni(α,2nγ); π from E2 γ to 2 ⁺ . T _{1/2} : weighted average of 0.19 ps 6 (α,pγ), 0.76 ps 14 Coul ex. (2006Le24), and 0.35 ps 2 Coul. ex. (2003Ko51). Others: 0.83 ps +42-28 from (α,2nγ), 0.19 ps 6 from (α,pγ), 0.17 ps +5-3 from (n,n'γ), and <0.7 ps from (α,α'γ). μ=+2.6 8, transient field integral perturbed angular correlation (tf) (2004An14). μ=+2.6 8, from γ-factor=+0.65 20 in Coul. ex. (2006Le24).
2703.6 4	(3)		CD FGH K VW	E(level): possible doublet indicated by conflicting parity measurements. J ^π : (1 ⁻ ,3 ⁻) from proton-capture yield in ⁶⁵ Cu(p,γ); (2 ⁻ ,3 ⁻) from L=(0) for a level at 2704 in ⁶⁷ Zn(d,t); (3 ⁺) from Hauser-Feshbach analysis in ⁶⁶ Zn(p,p').
2762.8 6	(2)		E	J ^π : from γ(θ) in ⁶⁶ Zn(γ,γ').
2765.56 7	4 ⁺	>7 ps	CD FGH JK mn U	XREF: Others: AC J ^π : 4 from γ(θ) in ⁶⁴ Ni(α,2nγ); π from L(p,p')=4. T _{1/2} : by DSA from ⁶⁴ Ni(α,2nγ). Other: >2.1 ps by DSA from ⁶³ Cu(α,pγ).
2780.157 7	2 ⁺	0.26 ps 7	B DEFGH K mnOp V	T _{1/2} : weighted average of 0.25 ps 8 from (γ,γ') and 0.28 ps 14 from (n,n'γ).
2826.69 5	3 ⁻ @	0.180 ps 7	BCD FGH JK OpQRSTU WX Z	XREF: Others: AB, AC T _{1/2} : From Coul. ex. (2006LeZU). T _{1/2} : Others: 0.18 ps 4 from ⁶⁶ Zn(α,α'γ), 0.23 ps 14 from ⁶³ Cu(α,pγ), and 0.17 ps 4 from (n,n'γ). Other: 1.0 ps +6-3 from ⁶⁴ Ni(α,2nγ). μ=+2.1 9, transient field integral perturbed angular correlation (tf) (2004An14). μ=+2.1 9, from γ-factor=+0.7 3 in Coul. ex. (2006Le24).
2938.074 3	2 ⁺	0.044 ps 16	B DEFGH JKLM OP V	T _{1/2} : unweighted average of 0.06 ps 5 from (α,pγ) and 0.028 ps 3 from (n,n'γ).
3030	(0 ⁺)			J ^π : from L(⁶ Li,d)=(0).
3077.73 23	4 ⁺	1.04 ps 7	CD FGH JK M OPQR TUV	XREF: Others: AB, AC Configuration=(π f _{5/2} 4 ⁺) T _{1/2} : Coul. ex. (2006Le24). T _{1/2} : 1.7 ps +10-3 in ⁶⁴ Ni(α,2nγ), 0.5 ps +3-2 in (α,pγ), 0.13 ps +56-10 in (α,α'γ), and 0.09 ps +5-3 in (n,n'γ).
3105.040 4	0 ⁺		B DEFGH K O	
3212.582 8	2 ⁺	0.083 ps +21-14	B DEFGH K m O	T _{1/2} : from DSAM in (n,n'γ).
3226.2 11			DE K m R	
3228.885 3	1 ⁺	0.12 ps 3	B D FGH K P V	T _{1/2} : from DSAM in (n,n'γ). J ^π : from log ft=6.14 4 in ε decay from 0 ⁺ and γ to 0 ⁺ ; π=+ from M1,E2 to 2 ⁺ .
3241.2? 11			E	
3331.441 6	2 ⁺	0.083 ps +21-14	B DEFGH KLm O R V	T _{1/2} : from DSAM in (n,n'γ).
3380.944 4	1 ⁻	20 fs 5	B DEFGH K p V	T _{1/2} : other: 42 fs +21-14 from DSA in (n,n'γ). J ^π : 1 from γ(θ) in ⁶⁶ Zn(γ,γ'); π from L(p,p')=(1).

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued)

⁶⁶Zn Levels (continued)

E(level) [†]	J ^π #	T _{1/2} [‡]	XREF					Comments	
3427.406 18	1,2 ⁻		B	GH			p		J ^π : log ft=8.9 1 from ε decay from 0 ⁺ ; γ to 2 ⁺ .
3432.408 4	1 ⁻	30 fs +19-8	B	DEFGH	K			R	J ^π : 0,1,2 ⁻ from log ft=7.03 4 in ε decay from 0 ⁺ ; J=0 ruled out by γ to 0 ⁺ , J ^π =2 ⁻ ruled out by T _{1/2} ; π=(-) from L(p,p')=(1).
3507.249 23	2 ⁺		B	DEFGH	K	m		V	J ^π : L(p,p')=2.
3523.6 8				CD	FG	K	m	o	
3531.692 14	0 ⁺		B	DEFG	K			o	
3576.370 22	4 ⁺		B	DEFGH	K			R	J ^π : L(p,p')=4.
3670.72 5	2 ⁺		B	D	GH	K	m	OP	
3689.01 16	1 ⁺ ,2 ⁺ ,3 ⁺			D	G	KLm		V	J ^π : L(d,t)=L(d,n)=1.
3709.4 3	(5)	0.6 ^a ps +6-2	CD	G	JK				J ^π : from γ(θ) in ⁶⁴ Ni(α,2nγ).
3725.3 5				D	F	K			
3731.6 5					G				
3738.207 21	+	14 fs 2	B	DEFGH	K			O	Lower J ^π member of a J ^π =(2 ⁺)+(4 ⁺) level doublet at 3737 10 reported in ⁶⁴ Zn(t,p). J ^π : J ^π =(2 ⁺) from L(t,p)=(2) not adopted for this lower J ^π member of 3737 10 doublet since J=1 from γ(θ) and reduced dipole γ rays in ⁶⁶ Zn(γ,γ') reported for a 3739.1 level. T _{1/2} : from (γ,γ') with Γ _{γ0} /Γ=0.75 3 from adopted gammas.
3738.24 4	(4 ⁺)							O	E(level): higher J ^π member of a J ^π =(2 ⁺)+(4 ⁺) level doublet reported in ⁶⁴ Zn(t,p).
3747.03 19	5 ⁻	46 ps 3	CD	GH	JK			p	XREF: Others: AB, AC Configuration=((π p _{3/2})(π g _{9/2}))5 ⁻ J ^π : 5 from γ(θ) in ⁶⁴ Ni(α,2pnγ); π from E2 to 3 ⁻ . T _{1/2} : by recoil distance in ⁵⁵ Mn(¹⁴ N,npγ). DSA measurements of 0.8 ps +11-4 from ⁶³ Cu(α,pγ) and 6 ps +14-3 from ⁶⁴ Ni(α,2nγ) do not take fully into account strong feeding from the 4252 (T _{1/2} =133 ps 11) and 4076 (T _{1/2} =29.8 ps 14) levels. Other: 0.21 ps +14-7 from (n,n'γ).
3753.01 4	4 ⁺		B	D	FGH	K		p	J ^π : L(p,p')=4.
3770 30	(1 ⁻)							QR	J ^π : from L(⁶ Li,d)=(1).
3791.123 3	1 ⁺		B	D	FGH	K	m	V	J ^π : 1 ⁺ from log ft=5.00 4 from 0 ⁺ .
3806.4 10				D		K	m		
3825.0 3	0 ⁺		B	DEF		K			J ^π : from γ(θ) in (γ,γ').
3874 5				D		K			
3882.424 10	(2 ⁺)		B	DE	G	K			J ^π : from γ(θ) in ⁶⁶ Zn(γ,γ').
3898.3 6	5 ⁻		CD	G	K		o	q	J ^π : 6 ⁺ from L(p,p')=6; γ's to 0 ⁺ and 2 ⁺ levels rule it out; level could be a doublet.
3924.71 20				D	FGH	K		q	
3946 2	(1 ⁻)			D		K		pqR	J ^π : L(p,p')=(1).
3969 2	(4 ⁺)			D		K	m	Op	
4005 10	4 ⁺			D			m		J ^π : L(p,p')=4.
4011.7				E			m	V	E(level): from ⁶⁶ Zn(γ,γ').
4019.2 15	2 ⁺			D	FG	K	m	O	
4075.7 3	(6 ⁻)	29.8 ps 14	C	E		J	L		XREF: Others: AC μ=0.9 2; Q=-0.081 13 (2005St24) μ: By recoil into gas and or vacuum (RIGV) (1983Ba69). Q: From electron scattering (Es); recalculated

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued)

⁶⁶Zn Levels (continued)

E(level) [†]	J ^π #	T _{1/2} [‡]	XREF	Comments
				value (1981Ko06). T _{1/2} : by recoil distance in ⁵⁵ Mn(¹⁴ N,2pnγ). Other: >1.4 ps from (α,pγ). J ^π : (6) from γ(θ), γ yields and decay systematics in ⁶⁴ Ni(α,2nγ); π from (M1) to 5 ⁻ .
4081.0 15			D F K m	
4085.983 4	1 ⁺		B D FGH K m V	J ^π : log ft=5.99 4 from 0 ⁺ .
4108.5 10			d F m R	π=+ from L(p,d)=1 for a level at 4100 30; π=- from L(p,p')=3 for a level at 4110 10, indicating a possible doublet.
4119.0 5	(1 ⁻)		d G K 0 q	
4182.7 5	(6 ⁺)	0.15 ps 6	CD JK	XREF: Others: AC J ^π : (6) from γ(θ) and γ yields in ⁶⁴ Ni(α,2nγ); π from (E2) to 4 ⁺ . L(p,p')=4 at 4189 7. T _{1/2} : unweighted mean of DSA measurements 0.08 ps +6-4 in ⁶³ Cu(α,pγ) and 0.21 ps +10-3 in ⁶⁴ Ni(α,2nγ).
4186 7	(3 ⁻)		K 0	E(level): weighted average of 4184 10 in (t,p) and 4188 10 in (p,α).
4223 2	(1 ⁻)		D K	XREF: Others: AB J ^π : L(p,p')=(1).
4251.9 3	(7 ⁻)	133 ps 10	C J Op	XREF: Others: AA μ=1.0 2 (2005St24,1989Ra17) Configuration=((ν F _{5/2})(ν G _{9/2}) ₇₋ +(ν P _{1/2})(ν G _{9/2}) ₅₋) E(level): unresolved doublet in (α, ² He) at 4220 50; L=7 assigned to the main level at 4220 with a tentative L=5 assigned to the 4400 level visible at some backward angles. μ: From recoil into gas and/or vacuum (RIGV) (1981Ko06). J ^π : (7) from γ(θ) and γ yields in ⁶⁴ Ni(α,2nγ); π from (E2) to 5 ⁻ . T _{1/2} : by recoil distance in ⁵⁵ Mn(¹⁴ N,2pnγ). Other: >0.55 ps in (α,pγ).
4258 2			D K p	
4267 7	4 ⁺		D K p	J ^π : L(p,p')=4.
4295.339 4	1 ⁺	4.2 fs +18-9	B DE K m	J ^π : 1 ⁺ from log ft=5.23 4 from 0 ⁺ in ε decay.
4321.83 20			D H K m	
4332 7	2 ⁺		D K 0	J ^π : L(t,p)=2.
4393.7 16	3 ⁻	0.07 ps +4-2	D K OPQ T	T _{1/2} : from DSA in ⁶⁶ Zn(α,α'γ). J ^π : 1 from γ(θ) in ⁶⁶ Zn(γ,γ').
4424 6	1	7.0 fs 12	DE H m	
4433 6	1 ⁻		D H m 0	
4439 7	2 ⁺		D m	J ^π : L(p,p')=2.
4454 5			D m	
4461.409 5	1 ⁺	7 fs +12-3	B DE H m	J ^π : 1 ⁺ from log ft=5.54 4 from 0 ⁺ in ε decay.
4472 7	3 ⁻		D m q	J ^π : L(p,p')=3.
4497.6 5			D H p	
4511 & 5	0 ⁺		D Opq	
4511 & 5	(2 ⁺)		D Opq	
4527 5			D pq	
4538 7	4 ⁺		D pq	J ^π : L(p,p')=4.
4565 2	3 ⁻		D q	J ^π : L(p,p')=3.
4567 10	5 ⁻		0 q	
4609? 2	(1)	8.4 fs +33-18	E p	J ^π : from γ(θ) in ⁶⁶ Zn(γ,γ').

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued)

⁶⁶Zn Levels (continued)

E(level) [†]	J ^π #	T _{1/2} [‡]	XREF			Comments
4610 5	4 ⁺		D		p	J ^π : L(p,p')=4.
4622 6			D	H	Op	
4635.3 11	(2)		DE			XREF: Others: AA
4638.24 14	1		B			J ^π : (2) from γ(θ) in ⁶⁶ Zn(γ,γ'). XREF: Others: AA
4645 10			D			J ^π : log ft=7.96 11 from 0 ⁺ ; γ to 0 ⁺ .
4655 7	(3 ⁻ ,4 ⁺)		D			XREF: Others: AA
4675.6 5	1 ⁺		B		1	J ^π : L(p,p')=3+4; probable doublet. XREF: Others: AA
4680 50						J ^π : log ft=8.35 15 from 0 ⁺ rules out 0 ⁺ to 0 ⁺ transition; γ to 2 ⁺ .
4683 10	(1)	7.1 fs +24-14	DE		1	XREF: Others: AA
4694 7	4 ⁺		D		1	XREF: Others: AA
4730 7	2 ⁺		D		1	J ^π : L(p,p')=4. XREF: Others: AA
4745 10			D		1	J ^π : L(p,p')=2.
4758 10			D		1	
4780 7	5 ⁻		D		q	
4796 7	(1 ⁻)		D		q	J ^π : L(p,p')=5. J ^π : L(p,p')=(1).
4806.199 5	1 ⁺	3.8 fs +13-8	B DE	H		J ^π : 1 ⁺ from log ft=4.89 4 from 0 ⁺ in ε decay.
4814.1 4	(7 ⁻)	0.6 ^d ps 4	C		J	XREF: Others: AB
4832 10			D			J ^π : (7) from γ(θ) and γ yields in ⁶⁴ Ni(α,2nγ); π from (M1) to (6 ⁻).
4849.93 3	1 ⁺		B D			J ^π : log ft=6.62 6 from 0 ⁺ rules out a 0 ⁺ to 0 ⁺ transition; γ to 2 ⁺ .
4866.056 16	1 ⁺		B D			J ^π : log ft=6.42 6 from 0 ⁺ ; γ to 0 ⁺ .
4875 10			D		q	
4885 10			D		q	
4907 10			D		q	
4918 10			D		q	
4945 10			D			
4958.2 4	1 ⁺		B D		1	J ^π : log ft=7.48 11 from 0 ⁺ rules out a 0 ⁺ to 0 ⁺ transition; γ to 2 ⁺ .
4984 10			D		1m	
5005.8 3	1 ⁺		B D		1m	J ^π : log ft=7.47 7 from 0 ⁺ ; γ to 0 ⁺ .
5025 10			D		1m	
5038 10			D			
5059 10			D		q	
5073 10			D		q	
5086 10			D		q	
5097 10			D		q	
5106 10			D		q	
5111.9 4	(8 ⁻)		CD		q	J ^π : from γ(θ) and γ decay systematics in ⁶⁴ Ni(α,2nγ).
5124 10			D		q	
5143 10			D		q	
5159 10			D			XREF: Others: AA
5169 10			D			XREF: Others: AA
5180 10			D			XREF: Others: AA
5198 10			D			XREF: Others: AA
5207.3 5	(8 ⁺)	>6 ps	C			XREF: Others: AA, AB

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued)

⁶⁶Zn Levels (continued)

E(level) [†]	J ^π #	T _{1/2} [‡]	XREF			Comments
						Configuration=(ν g _{9/2} 8 ⁺) configuration from (¹² C, ¹⁰ Be), (¹² C, ¹⁰ C) reaction where this level is seen strongly excited (1990Bo27). This level was seen weakly excited in the (α, ² He) reaction (1990Fi07). J ^π : (8) from γ(θ), γ yields and decay systematics in ⁶⁴ Ni(α,2nγ); π from (E2) to (6) ⁺ . T _{1/2} : by DSA in ⁶⁴ Ni(α,2nγ). XREF: Others: AA XREF: Others: AA XREF: Others: AA
5222 10			D			
5234 10			D			
5245 10			D			
5263 10			D			
5274 10			D			
5285 10			D			
5305 10			D			
5322 10			D			
5331 10			D	q		
5352 10			D	q		
5364 10			D	q		
5375 10			D	q		
5389 10			D	q		
5403 10			D	q		
5420 10			D		XREF: Others: AB	
5431 10			D			
5446 10			D			
5464.4 5	(9 ⁻)	1.9 ps 8	C	q		J ^π : (9) from γ(θ) in ⁶⁴ Ni(α,2nγ); π from (E2) to (7 ⁻). T _{1/2} : by recoil distance in ⁵⁵ Mn(¹⁴ N,2npγ). T _{1/2} =2.8 ps +2I-14 by DSA in ⁶⁴ Ni(α,2nγ).
5500 45				q		
5650 30	3 ⁻		D	q		XREF: Others: AB E(level): from ⁶⁶ Zn(p,p'). J ^π : L(p,p')=3.
5740 50			A	J	q	
6000 50						XREF: Others: AB
6292.6 6	(10 ⁺)	1.6 ps +7-3	C	J		J ^π : (10) from γ(θ) in ⁶⁴ Ni(α,2nγ); π from (E2) to (8 ⁺). T _{1/2} : by DSA in ⁶⁴ Ni(α,2nγ).
6419.0 8			C			
6850 50	(8 ⁺)					XREF: Others: AB Configuration=(π g _{9/2} 8 ⁺) J ^π : from DWBA analysis of (¹² C, ¹⁰ Be), (¹² C, ¹⁰ C) data.
7.17×10 ³ 18				J		
7367.4 4	1	1.47 fs 16	E			J ^π : from γ(θ) in ⁶⁶ Zn(γ,γ'). T _{1/2} : by DSA in ⁶⁴ Ni(α,2nγ).
7517.3 10		1.5 ps +6-3	C			XREF: Others: AB
7550 50	(6 ⁺)					Configuration=((π g _{9/2})(π d _{5/2}))6 ⁺ J ^π : from DWBA analysis of (¹² C, ¹⁰ Be), (¹² C, ¹⁰ C) data.
7693.3 3	1	2.2 fs 4	E			J ^π : 1 from γ(θ) in ⁶⁶ Zn(γ,γ').
11059.9 10	2 ⁻ ,3 ⁻			I		J ^π : from n(θ) in ⁶⁵ Cu(p,n) IAR.
11395 10				I		
11411 10				I		
11457 10				I		

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{66}Zn Levels (continued)

<u>E(level)[†]</u>	<u>J^π#</u>	<u>XREF</u>	<u>Comments</u>
11514	10	I	
11593	10	I	
11654	10	I	
11698	10	I	
11757	10	I	
11841	10 (2 ⁺)	I	J ^π : from n(θ) and γ (θ) in $^{65}\text{Cu}(\text{p,n}),(\text{p,n}\gamma)$ IAR.
11916	10	I	
12194?	10	I	
12218	10	I	
12293	10	I	
12324	10	I	
12401	10	I	
12433?	10	I	
12552	10	I	
12602	10	I	
12651	10	I	
12688	10	I	
12714	10	I	

[†] Except as noted, levels with E<10000 are as follows: 1) from a least-squares fit to adopted E γ data; 2) energies with $\Delta E=2-10$ are from $^{66}\text{Zn}(\text{p,p}')$; 3) energies with $\Delta E>10$ are from $^{66}\text{Zn}(\alpha,\alpha')$. Levels with E>10000 are from $^{65}\text{Cu}(\text{p,n}),(\text{p,n}\gamma)$ IAR.

[‡] Except as noted, T_{1/2} is from measured widths in $^{66}\text{Zn}(\gamma,\gamma')$ and adopted γ branchings.

From L-transfer in $^{64}\text{Zn}(\text{t,p})$, except as noted.

@ Consistent with angular distribution and analyzing-power data in (pol p,p') ([1993Mo15](#)).

& Doublet reported in $^{64}\text{Zn}(\text{t,p})$. E=4511.5 given for one level in $^{66}\text{Zn}(\text{p,p}')$.

^a By DSA from $^{63}\text{Cu}(\alpha,\text{p}\gamma)$.

Adopted Levels, Gammas (continued)

E _i (level)	J _i ^π	E _γ [‡]	I _γ [#]	E _f	J _f ^π	Mult.	γ(⁶⁶ Zn)		I _(γ+ce)	Comments
							δ ^a	α [†]		
1039.2279	2 ⁺	1039.220 3	100.0 3	0.0	0 ⁺	E2 [@]		0.000269 4		α(K)=0.000241 4; α(L)=2.41×10 ⁻⁵ 4; α(M)=3.46×10 ⁻⁶ 5; α(N+..)=1.384×10 ⁻⁷ 20 α(N)=1.384×10 ⁻⁷ 20 B(E2)(W.u.)=17.5 4
1872.7653	2 ⁺	833.5324 21	100.0 4	1039.2279	2 ⁺	M1+E2	-1.6 2	0.000434 9		α(K)=0.000389 8; α(L)=3.91×10 ⁻⁵ 8; α(M)=5.60×10 ⁻⁶ 11; α(N+..)=2.24×10 ⁻⁷ 5 α(N)=2.24×10 ⁻⁷ 5 B(M1)(W.u.)=0.056 23; B(E2)(W.u.)=3.3×10 ² 13 Mult.: D+Q from γ(θ) in ⁶⁶ Cu β ⁻ decay; M1+E2 from RUL. δ: From γγ(θ) in ⁶⁶ Ga ε DECAFY (2002Ga20). Other value: -1.9 3, as given by 1974Kr16 from analysis of 1969Ha46 data.
		1872.740 6	0.39 3	0.0	0 ⁺	[E2]		0.000328 5		α(K)=7.04×10 ⁻⁵ 10; α(L)=6.99×10 ⁻⁶ 10; α(M)=1.001×10 ⁻⁶ 14; α(N+..)=0.000250 4 α(N)=4.05×10 ⁻⁸ 6; α(IPF)=0.000250 4 B(E2)(W.u.)=0.032 12
2372.353	0 ⁺	499.590 6	0.41 10	1872.7653	2 ⁺	E2		0.00199 3		α(K)=0.001782 25; α(L)=0.000182 3; α(M)=2.61×10 ⁻⁵ 4; α(N+..)=1.013×10 ⁻⁶ 15 α(N)=1.013×10 ⁻⁶ 15 B(E2)(W.u.)<22 B(E2)(W.u.)<40
		1333.112 5	100.0 4	1039.2279	2 ⁺	E2		0.000190 3		α(K)=0.0001383 20; α(L)=1.379×10 ⁻⁵ 20; α(M)=1.98×10 ⁻⁶ 3; α(N+..)=3.61×10 ⁻⁵ 5 α(N)=7.96×10 ⁻⁸ 12; α(IPF)=3.61×10 ⁻⁵ 5 Mult.: Q from γ(θ) in ⁶⁶ Zn(p,p'γ); E2 from RUL.
		2372.375 ^d		0.0	0 ⁺	E0			6.6×10 ⁻³ 13	E _γ : from level energy difference; 2372.2 from (p,p'γ). Mult.: from internal conversion data and absence of γ ray in (p,p'γ).
2451.01	4 ⁺	1411.75 ^b 5	100 ^f	1039.2279	2 ⁺	E2 [@]		0.000194 3		I _(γ+ce) : for 100 transitions of 1333γ from (p,p'γ). α(K)=0.0001227 18; α(L)=1.222×10 ⁻⁵ 18; α(M)=1.751×10 ⁻⁶ 25 α(N)=7.06×10 ⁻⁸ 10; α(IPF)=5.69×10 ⁻⁵ 8 B(E2)(W.u.)=18 3 δ(M3/E2)=+0.04 4.
2703.6	(3)	2450 ^d 2 830.7 ^c 8	2 ^f 100 ^g 11	0.0 1872.7653	0 ⁺ 2 ⁺					E _γ : not confirmed by any other ⁶⁶ Zn data.

8

Adopted Levels, Gammas (continued)

γ(⁶⁶Zn) (continued)

<u>E_i(level)</u>	<u>J_i^π</u>	<u>E_γ[‡]</u>	<u>I_γ[#]</u>	<u>E_f</u>	<u>J_f^π</u>	<u>Mult.</u>	<u>δ^a</u>	<u>α[†]</u>	<u>Comments</u>
2703.6	(3)	1664.4 ^b 4	7.1 ^g 10	1039.2279	2 ⁺	(D+Q)	8.5 15		Mult.,δ: from γ(θ) in ⁶⁶ Zn(n,n'γ).
2762.8	(2)	2762	100	0.0	0 ⁺				E _γ : reported only in ⁶⁶ Zn(γ,γ').
2765.56	4 ⁺	314.6 ^b 1 892.7 ^b 1	36 ^b 2 100 ^b 3	2451.01 1872.7653	4 ⁺ 2 ⁺	E2 [@]		0.000388 6	B(E2)(W.u.)<4.0 α(K)=0.000348 5; α(L)=3.50×10 ⁻⁵ 5; α(M)=5.01×10 ⁻⁶ 7; α(N+..)=2.00×10 ⁻⁷ 3 α(N)=2.00×10 ⁻⁷ 3 δ(M3/E2)=-0.04 6.
		1726.4 ^b 2	87 ^b 3	1039.2279	2 ⁺	E2 [@]		0.000274 4	B(E2)(W.u.)<0.13 α(K)=8.21×10 ⁻⁵ 12; α(L)=8.16×10 ⁻⁶ 12; α(M)=1.169×10 ⁻⁶ 17; α(N+..)=0.000183 3 α(N)=4.73×10 ⁻⁸ 7; α(IPF)=0.000183 3 δ(M3/E2)=+0.00 3.
2780.157	2 ⁺	328.8 ^b 5 907.394 19 1740.904 16	11 ^b 3 17.7 12 23.1 3	2451.01 1872.7653 1039.2279	4 ⁺ 2 ⁺ 2 ⁺	M1+E2	0.33 28	0.000241 8	α(K)=7.74×10 ⁻⁵ 13; α(L)=7.67×10 ⁻⁶ 13; α(M)=1.100×10 ⁻⁶ 19; α(N+..)=0.000155 7 α(N)=4.47×10 ⁻⁸ 8; α(IPF)=0.000155 7 B(M1)(W.u.)=0.0022 7; B(E2)(W.u.)=0.13 +20-13
		2780.095 16	100.0 24	0.0	0 ⁺	E2		0.000722 11	α(K)=3.51×10 ⁻⁵ 5; α(L)=3.47×10 ⁻⁶ 5; α(M)=4.98×10 ⁻⁷ 7; α(N+..)=0.000683 10 α(N)=2.02×10 ⁻⁸ 3; α(IPF)=0.000683 10 B(E2)(W.u.)=0.54 15 Mult.: Q from γ(θ) in ⁶⁶ Zn(n,n'γ); E2 from RUL.
2826.69	3 ⁻	953.93 9 1787.44 9	11.3 ^g 13 100 ^g 9	1872.7653 1039.2279	2 ⁺ 2 ⁺	(E1) ^{&}		0.000526 8	α(K)=4.21×10 ⁻⁵ 6; α(L)=4.16×10 ⁻⁶ 6; α(M)=5.95×10 ⁻⁷ 9; α(N+..)=0.000479 7 α(N)=2.41×10 ⁻⁸ 4; α(IPF)=0.000479 7 B(E1)(W.u.)=0.00036 5 δ(M2/E1)=-0.04 5.
2938.074	2 ⁺	1065.305 9 1898.823 8	0.60 12 100.0 8	1872.7653 1039.2279	2 ⁺ 2 ⁺	(M1+E2)	0.03 1	0.000288 4	α(K)=6.58×10 ⁻⁵ 10; α(L)=6.52×10 ⁻⁶ 10; α(M)=9.35×10 ⁻⁷ 13; α(N+..)=0.000215 3 α(N)=3.80×10 ⁻⁸ 6; α(IPF)=0.000215 3 B(M1)(W.u.)=(0.043 17); B(E2)(W.u.)=(0.017 14) Mult.: D+Q from γ(θ) in ⁶⁶ Zn(n,n'γ); M1+E2 from ΔJ ^π . δ: from γ(θ) in (n,n'γ); sign convention not specified.
3077.73	4 ⁺	2941 ^h 312.0 ^c 8	71 24 10 ^c 3	0.0 2765.56	0 ⁺ 4 ⁺				E _γ ,I _γ : reported only in ⁶⁶ Zn(γ,γ'). E _γ : doublet. E _γ is from level energy difference in

Adopted Levels, Gammas (continued)

$\gamma(^{66}\text{Zn})$ (continued)										
$E_i(\text{level})$	J_i^π	E_γ^\ddagger	$I_\gamma^\#$	E_f	J_f^π	Mult.	δ^a	α^\ddagger	$I_{(\gamma+ce)}$	Comments
3077.73	4 ⁺	627.6 ^c 4	100 ^c 10	2451.01	4 ⁺	(M1+E2)&	-0.25 12	0.000688 23		⁶⁴ Ni(α ,2n γ). $E_\gamma=312.8$ from level energy difference in ⁶⁵ Cu(p, γ). $\alpha(\text{K})=0.000617$ 21; $\alpha(\text{L})=6.19\times 10^{-5}$ 21; $\alpha(\text{M})=8.9\times 10^{-6}$ 3; $\alpha(\text{N}+..)=3.57\times 10^{-7}$ 12 $\alpha(\text{N})=3.57\times 10^{-7}$ 12 B(M1)(W.u.)=(0.056 9); B(E2)(W.u.)=(15 14) $\alpha(\text{K})=0.0001723$ 25; $\alpha(\text{L})=1.721\times 10^{-5}$ 25; $\alpha(\text{M})=2.47\times 10^{-6}$ 4; $\alpha(\text{N}+..)=9.43\times 10^{-6}$ 1 $\alpha(\text{N})=9.91\times 10^{-8}$ 14; $\alpha(\text{IPF})=9.33\times 10^{-6}$ 15 B(E2)(W.u.)=3.1 7 $\delta(\text{M3/E2})=-0.15$ 15.
		1204.2 ^c 5	33 ^c 6	1872.7653	2 ⁺	(E2)&		0.000201 3		
3105.040	0 ⁺	1232.264 8 2065.778 7 3105.064	100 4 6.2 3	1872.7653 2 ⁺ 1039.2279 2 ⁺ 0.0 0 ⁺		E0			2.4 $\times 10^{-2}$ 5	E_γ : from level energy difference; 3107.0 from (p,p' γ). Mult.: from internal conversion data and absence of γ ray in (p,p' γ). $I_{(\gamma+ce)}$: for 100 transitions of 1234 γ from (p,p' γ). E_γ : placed as depopulating the 3213 level in ⁶⁵ Cu(p, γ), ⁶⁶ Zn(p,p' γ), ⁶⁶ Zn(n,n' γ) and by 1971Ca14, 1994En02 in ⁶⁶ Ga ϵ decay. Placed as depopulating a proposed level at 4047 by 1970Ph01 in ϵ decay, which level, however, is not observed by 1994En02.
3212.582	2 ⁺	2173.319 15	100 6	1039.2279 2 ⁺						E_γ : placed as depopulating the 3213 level in ⁶⁵ Cu(p, γ), ⁶⁶ Zn(p,p' γ), ⁶⁶ Zn(n,n' γ) and by 1971Ca14, 1994En02 in ⁶⁶ Ga ϵ decay. Placed as depopulating a proposed level at 4047 by 1970Ph01 in ϵ decay, which level, however, is not observed by 1994En02.
3228.885	1 ⁺	3212.499 19 290.8105 11 448.73 2	2.2 5 0.92 3 2.01 7	0.0 0 ⁺ 2938.074 2 ⁺ 2780.157 2 ⁺		M1+E2	-0.02 3	0.001419 20		$\alpha(\text{K})=0.001272$ 18; $\alpha(\text{L})=0.0001283$ 19; $\alpha(\text{M})=1.84\times 10^{-5}$ 3; $\alpha(\text{N}+..)=7.39\times 10^{-7}$ $\alpha(\text{N})=7.39\times 10^{-7}$ 11
		856.527 10 1356.104 9 2189.616 6	2.09 12 6.7 7 100.0 5	2372.353 0 ⁺ 1872.7653 2 ⁺ 1039.2279 2 ⁺		M1+E2	0.12 2	0.000398 6		$\alpha(\text{K})=5.12\times 10^{-5}$ 8; $\alpha(\text{L})=5.07\times 10^{-6}$ 7; $\alpha(\text{M})=7.26\times 10^{-7}$ 11; $\alpha(\text{N}+..)=0.000341$ 5 $\alpha(\text{N})=2.95\times 10^{-8}$ 5; $\alpha(\text{IPF})=0.000341$ 5 B(M1)(W.u.)=0.012 3; B(E2)(W.u.)=0.060 25 Mult.: from ce measurement in ϵ decay.
		3228.800 6	28.3 2	0.0 0 ⁺		M1		0.000812 12		$\alpha(\text{K})=2.68\times 10^{-5}$ 4; $\alpha(\text{L})=2.64\times 10^{-6}$ 4; $\alpha(\text{M})=3.79\times 10^{-7}$ 6; $\alpha(\text{N}+..)=0.000782$ 11

Adopted Levels, Gammas (continued)

$\gamma(^{66}\text{Zn})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ [‡]	I_γ [#]	E_f	J_f^π	Mult.	δ^a	α^\dagger	Comments
									$\alpha(\text{N})=1.544\times 10^{-8}$ 22; $\alpha(\text{IPF})=0.000782$ 11 $\text{B}(\text{M1})(\text{W.u.})=0.0011$ 3
3331.441	2 ⁺	551.284 22 1458.662 12	7.2 7 100.0 23	2780.157 1872.7653	2 ⁺ 2 ⁺	(M1+E2)	-0.01 9	0.0001741 25	$\alpha(\text{K})=0.0001070$ 15; $\alpha(\text{L})=1.062\times 10^{-5}$ 15; $\alpha(\text{M})=1.523\times 10^{-6}$ 22 $\alpha(\text{N})=6.18\times 10^{-8}$ 9; $\alpha(\text{IPF})=5.49\times 10^{-5}$ 8 $\text{B}(\text{M1})(\text{W.u.})=(0.058 +10-15)$
3380.944	1 ⁻	2292.171 13 3331.351 14 442.873 14 600.788 21 1008.588 12 1508.158 7 2341.673 11	17.6 12 23 3 1.06 8 0.92 6 4.0 5 37.8 2 0.22 5	1039.2279 0.0 2938.074 2780.157 2372.353 1872.7653 1039.2279	2 ⁺ 0 ⁺ 2 ⁺ 2 ⁺ 0 ⁺ 2 ⁺ 2 ⁺				
3427.406	1,2 ⁻	3380.850 6 1554.62 3	100.0 6 100	1872.7653	0 ⁺ 2 ⁺				
3432.408	1 ⁻	494.336 1 1060.051 1 1559.627 1 2393.129 7	32.0 3 15.4 4 7.6 6 82 3	2938.074 2372.353 1872.7653 1039.2279	2 ⁺ 0 ⁺ 2 ⁺ 2 ⁺	E1		0.000924 13	$\alpha(\text{K})=2.73\times 10^{-5}$ 4; $\alpha(\text{L})=2.70\times 10^{-6}$ 4; $\alpha(\text{M})=3.86\times 10^{-7}$ 6; $\alpha(\text{N}+..)=0.000894$ 13 $\alpha(\text{N})=1.569\times 10^{-8}$ 22; $\alpha(\text{IPF})=0.000894$ 13 $\text{B}(\text{E1})(\text{W.u.})=0.00035 +10-23$
3507.249	2 ⁺	3432.309 7 680.56 10 1634.46 7 2467.97 7	100.0 13 18 5 42 7 100 9	0.0 2826.69 1872.7653 1039.2279	0 ⁺ 3 ⁻ 2 ⁺ 2 ⁺				
3523.6		758.0 ^c 8	100	2765.56	4 ⁺				
3531.692	0 ⁺	2492.42 3	100	1039.2279	2 ⁺				
3576.370	4 ⁺	749.68 10 796.21 5 1703.59 5 2537.09 5	25 8 53 12 100 4 93 20	2826.69 2780.157 1872.7653 1039.2279	3 ⁻ 2 ⁺ 2 ⁺ 2 ⁺				
3670.72	2 ⁺	441.822 1219.1 ^b 2 1797.94 9 2631.44 9	100 9 39 ^b 7 18 5 28 11	3228.885 2451.01 1872.7653 1039.2279	1 ⁺ 4 ⁺ 2 ⁺ 2 ⁺				E_γ : from level energy difference; 440.5 1 from (n,n' γ). I_γ : from (n,n' γ).
3689.01	1 ⁺ ,2 ⁺ ,3 ⁺	1815.4 ^h 5	100	1872.7653	2 ⁺				E_γ : reported only in ⁶⁵ Cu(p, γ).
3709.4	(5)	943.8 ^c 3	100 ^c	2765.56	4 ⁺	(D+Q) ^{&}	-1.5 +2-1		

Adopted Levels, Gammas (continued)

$\gamma(^{66}\text{Zn})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\ddagger	$I_\gamma^\#$	E_f	J_f^π	Mult.	δ^a	α^\dagger	Comments
3725.3		293.0 ^d 5	11 ^f	3432.408	1 ⁻				
		962 ^d 1	100 ^f	2765.56	4 ⁺				
3731.6		966.0 ^e 5	100 ^g	2765.56	4 ⁺				E_γ : reported only in ⁶⁵ Cu(p, γ).
3738.207	+	800.13 5	7 4	2938.074	2 ⁺				
		2698.92 5	27 5	1039.2279	2 ⁺				
		3738.10 5	100 6	0.0	0 ⁺				
3747.03	5 ⁻	669.5 ^c 3	21 ^c 2	3077.73	4 ⁺	(E1)&		0.000294 5	$\alpha(\text{K})=0.000264$ 4; $\alpha(\text{L})=2.63\times 10^{-5}$ 4; $\alpha(\text{M})=3.77\times 10^{-6}$ 6; $\alpha(\text{N}+..)=1.511\times 10^{-7}$ 22 $\alpha(\text{N})=1.511\times 10^{-7}$ 22 $\text{B}(\text{E}1)(\text{W.u.})=4.9\times 10^{-6}$ 7 $\delta(\text{M}2/\text{E}1)=-0.04$ 6.
		919.9 ^c 3	4.9 ^c 5	2826.69	3 ⁻	E2@		0.000360 5	$\alpha(\text{K})=0.000323$ 5; $\alpha(\text{L})=3.24\times 10^{-5}$ 5; $\alpha(\text{M})=4.65\times 10^{-6}$ 7; $\alpha(\text{N}+..)=1.85\times 10^{-7}$ 3 $\alpha(\text{N})=1.85\times 10^{-7}$ 3 $\text{B}(\text{E}2)(\text{W.u.})=0.045$ 7 $\delta(\text{M}3/\text{E}2)=-0.00$ 6.
		981.5 ^c 5	<6 ^c	2765.56	4 ⁺				
		1295.6 ^c 4	100 ^c 10	2451.01	4 ⁺	(E1+M2)&	-0.04 2	0.000193 3	$\alpha(\text{K})=7.15\times 10^{-5}$ 11; $\alpha(\text{L})=7.09\times 10^{-6}$ 11; $\alpha(\text{M})=1.015\times 10^{-6}$ 16; $\alpha(\text{N}+..)=0.0001134$ $\alpha(\text{N})=4.10\times 10^{-8}$ 7; $\alpha(\text{IPF})=0.0001134$ 17 $\text{B}(\text{E}1)(\text{W.u.})=(3.2\times 10^{-6})$ 5; $\text{B}(\text{M}2)(\text{W.u.})=(0.014 +15-14)$
3753.01	4 ⁺	2713.73 5	100	1039.2279	2 ⁺				
3791.123	1 ⁺	283.87 3	0.016 4	3507.249	2 ⁺				
		410.178 12	0.289 12	3380.944	1 ⁻				
		459.683 14	0.387 17	3331.441	2 ⁺				
		562.241 10	0.029 3	3228.885	1 ⁺				
		578.540 19	0.26 4	3212.582	2 ⁺				
		686.080 6	1.11 4	3105.040	0 ⁺				
		853.038 8	0.334 9	2938.074	2 ⁺	M1+E2	0.37 18	0.000357 11	$\alpha(\text{K})=0.000321$ 10; $\alpha(\text{L})=3.20\times 10^{-5}$ 11; $\alpha(\text{M})=4.59\times 10^{-6}$ 15; $\alpha(\text{N}+..)=1.85\times 10^{-7}$ 6 $\alpha(\text{N})=1.85\times 10^{-7}$ 6
		1010.957 19	0.119 7	2780.157	2 ⁺				
		1418.754 5	2.703 13	2372.353	0 ⁺				
		1918.329 5	8.76 4	1872.7653	2 ⁺	M1+E2	-0.07 3	0.000295 5	$\alpha(\text{K})=6.47\times 10^{-5}$ 9; $\alpha(\text{L})=6.40\times 10^{-6}$ 9; $\alpha(\text{M})=9.18\times 10^{-7}$ 13; $\alpha(\text{N}+..)=0.000223$ 4 $\alpha(\text{N})=3.73\times 10^{-8}$ 6; $\alpha(\text{IPF})=0.000223$ 4
		2751.835 5	100.0 5	1039.2279	2 ⁺	(M1+E2)	-0.12 2	0.000626 9	$\alpha(\text{K})=3.48\times 10^{-5}$ 5; $\alpha(\text{L})=3.43\times 10^{-6}$ 5; $\alpha(\text{M})=4.92\times 10^{-7}$ 7; $\alpha(\text{N}+..)=0.000588$ 9 $\alpha(\text{N})=2.00\times 10^{-8}$ 3; $\alpha(\text{IPF})=0.000588$ 9

Adopted Levels, Gammas (continued)

γ(⁶⁶Zn) (continued)

<u>E_i(level)</u>	<u>J_i^π</u>	<u>E_γ[‡]</u>	<u>I_γ[#]</u>	<u>E_f</u>	<u>J_f^π</u>	<u>Mult.</u>	<u>δ^a</u>	<u>α[†]</u>	<u>Comments</u>
3791.123	1 ⁺	3791.004 8	2.941 24	0.0	0 ⁺	M1		0.001017 15	α(K)=2.08×10 ⁻⁵ 3; α(L)=2.05×10 ⁻⁶ 3; α(M)=2.95×10 ⁻⁷ 5; α(N+..)=0.000994 14 α(N)=1.200×10 ⁻⁸ 17; α(IPF)=0.000994 14
3825.0	0 ⁺	2785.7 3	100	1039.2279	2 ⁺				
3882.424	(2) ⁺	1507 ^h		2372.353	0 ⁺				E _γ ,I _γ : reported only in ⁶⁶ Zn(γ,γ').
		2009.628 16	100 21	1872.7653	2 ⁺				
		2843.130 16	54 11	1039.2279	2 ⁺				
3898.3	5 ⁻	1071.3 ^c 7	100 ^c	2826.69	3 ⁻	(E2(+M3))&	-0.04 20	0.00025 4	α(K)=0.00023 4; α(L)=2.3×10 ⁻⁵ 4; α(M)=3.2×10 ⁻⁶ 6; α(N+..)=1.30×10 ⁻⁷ 21 α(N)=1.30×10 ⁻⁷ 21
3924.71		1555 ^d 1	100 ^f	2372.353	0 ⁺				E _γ : possible doublet. E _γ =1553.0 in ⁶⁵ Cu(p,γ). I _γ : I _γ (1555)/I _γ (2888)<0.6 from ⁶⁵ Cu(p,γ). I _γ : relative branching from ⁶⁶ Zn(p,p'γ).
		2885.3 ^b 2	33	1039.2279	2 ⁺				
4019.2	2 ⁺	1239.0 ^d 15	100 ^f	2780.157	2 ⁺				
4075.7	(6 ⁻)	328.6 ^c 2	100 ^c	3747.03	5 ⁻	(M1+E2)@	+0.10 2	0.00299 5	α(K)=0.00268 5; α(L)=0.000272 5; α(M)=3.91×10 ⁻⁵ 7; α(N+..)=1.559×10 ⁻⁶ 25 α(N)=1.559×10 ⁻⁶ 25 B(M1)(W.u.)=(0.0206 10); B(E2)(W.u.)=(3.1 13)
4085.983	1 ⁺	347.77 5	0.14 5	3738.207	+				
		554.28 3	0.35 4	3531.692	0 ⁺				
		653.568 14	0.10 4	3432.408	1 ⁻				
		658.57 3	0.59 6	3427.406	1,2 ⁻				
		705.031 15	0.30 4	3380.944	1 ⁻				
		857.093 9	1.2 4	3228.885	1 ⁺				
		873.392 21	1.34 9	3212.582	2 ⁺				
		980.934 13	3.80 15	3105.040	0 ⁺				
		1147.896 10	6.15 21	2938.074	2 ⁺	M1+E2	-0.18 5	0.000192 3	α(K)=0.0001708 25; α(L)=1.700×10 ⁻⁵ 25; α(M)=2.44×10 ⁻⁶ 4; α(N+..)=2.28×10 ⁻⁶ 4 α(N)=9.87×10 ⁻⁸ 14; α(IPF)=2.18×10 ⁻⁶ 4
		1305.807 21	0.31 4	2780.157	2 ⁺				
		1713.602 12	1.92 9	2372.353	0 ⁺				
		2213.181 9	10.3 4	1872.7653	2 ⁺	M1+E2	-0.23 5	0.000410 6	α(K)=5.03×10 ⁻⁵ 7; α(L)=4.98×10 ⁻⁶ 7; α(M)=7.14×10 ⁻⁷ 10; α(N+..)=0.000354 6 α(N)=2.90×10 ⁻⁸ 4; α(IPF)=0.000354 6
		3046.684 9	4.47 18	1039.2279	2 ⁺	M1+E2	-0.8 2	0.000778 16	α(K)=2.97×10 ⁻⁵ 5; α(L)=2.94×10 ⁻⁶ 5; α(M)=4.21×10 ⁻⁷ 6; α(N+..)=0.000745 16 α(N)=1.712×10 ⁻⁸ 25; α(IPF)=0.000745 16
		4085.853 9	100.0 6	0.0	0 ⁺	M1		0.001117 16	α(K)=1.86×10 ⁻⁵ 3; α(L)=1.83×10 ⁻⁶ 3;

Adopted Levels, Gammas (continued)

$\gamma(^{66}\text{Zn})$ (continued)

<u>E_i(level)</u>	<u>J^{π}_i</u>	<u>E_{γ}[‡]</u>	<u>I_{γ}[#]</u>	<u>E_f</u>	<u>J^{π}_f</u>	<u>Mult.</u>	<u>δ^a</u>	<u>α^\dagger</u>	<u>Comments</u>
4119.0	(1 ⁻)	3079.7 ^e 5	100 ^g	1039.2279	2 ⁺				$\alpha(\text{M})=2.63\times 10^{-7}$ 4; $\alpha(\text{N}+..)=0.001096$ 16 $\alpha(\text{N})=1.070\times 10^{-8}$ 15; $\alpha(\text{IPF})=0.001096$ 16 E _{γ} : reported only in ⁶⁵ Cu(p, γ).
4182.7	(6 ⁺)	1732.9 ^c 5	100 ^c	2451.01	4 ⁺	(E2) [@]		0.000276 4	$\alpha(\text{K})=8.15\times 10^{-5}$ 12; $\alpha(\text{L})=8.10\times 10^{-6}$ 12; $\alpha(\text{M})=1.161\times 10^{-6}$ 17; $\alpha(\text{N}+..)=0.000186$ 3 $\alpha(\text{N})=4.69\times 10^{-8}$ 7; $\alpha(\text{IPF})=0.000186$ 3 B(E2)(W.u.)=15 6 $\delta(\text{M3/E2})<5.3\times 10^{-4}$ from RUL. $\delta=-0.10$ 5 from $\gamma(\theta)$ in (d,2n γ).
4251.9	(7 ⁻)	175.9 ^c 3	87 ^c 9	4075.7	(6 ⁻)	(M1+E2) [@]	+0.09 2	0.0144 4	$\alpha(\text{K})=0.0128$ 3; $\alpha(\text{L})=0.00133$ 4; $\alpha(\text{M})=0.000190$ 5; $\alpha(\text{N}+..)=7.48\times 10^{-6}$ 17 $\alpha(\text{N})=7.48\times 10^{-6}$ 17 B(M1)(W.u.)=(0.0139 21); B(E2)(W.u.)=(6 3)
		504.7 ^c 3	100 ^c 10	3747.03	5 ⁻	(E2) [@]		0.00193 3	$\alpha(\text{K})=0.001726$ 25; $\alpha(\text{L})=0.0001766$ 25; $\alpha(\text{M})=2.52\times 10^{-5}$ 4; $\alpha(\text{N}+..)=9.82\times 10^{-7}$ 14 $\alpha(\text{N})=9.82\times 10^{-7}$ 14 B(E2)(W.u.)=4.4 7 $\delta(\text{M3/E2})=-0.00$ 2.
4295.339	1 ⁺	412.916 16 557.13 5	0.088 13 0.161 17	3882.424 3738.207	(2) ⁺ +	M1+E2		0.0011 3	$\alpha(\text{K})=0.00103$ 25; $\alpha(\text{L})=0.00010$ 3; $\alpha(\text{M})=1.5\times 10^{-5}$ 4; $\alpha(\text{N}+..)=5.9\times 10^{-7}$ 14 $\alpha(\text{N})=5.9\times 10^{-7}$ 14
		718.97 5	0.260 20	3576.370	4 ⁺				
		763.64 3	0.233 20	3531.692	0 ⁺				
		862.926 13	0.398 20	3432.408	1 ⁻				
		867.93 3	0.114 14	3427.406	1,2 ⁻				
		914.388 14	0.71 4	3380.944	1 ⁻				
		963.892 15	0.38 3	3331.441	2 ⁺				
		1066.450 12	0.062 12	3228.885	1 ⁺				
		1082.75 2	0.348 20	3212.582	2 ⁺				
		1190.287 7	3.35 19	3105.040	0 ⁺				
		1357.250 12	4.3 13	2938.074	2 ⁺	M1+E2	-0.18 5	0.0001689 24	$\alpha(\text{K})=0.0001231$ 18; $\alpha(\text{L})=1.223\times 10^{-5}$ 18; $\alpha(\text{M})=1.753\times 10^{-6}$ 25 $\alpha(\text{N})=7.11\times 10^{-8}$ 10; $\alpha(\text{IPF})=3.18\times 10^{-5}$ 5 B(M1)(W.u.)=0.054 +20-29; B(E2)(W.u.)=1.5 +10-12
		1515.162 20	0.162 15	2780.157	2 ⁺				
		2422.525 7	49.37 24	1872.7653	2 ⁺	M1+E2	0.01 3	0.000491 7	$\alpha(\text{K})=4.30\times 10^{-5}$ 6; $\alpha(\text{L})=4.25\times 10^{-6}$ 6; $\alpha(\text{M})=6.10\times 10^{-7}$ 9; $\alpha(\text{N}+..)=0.000443$ 7 $\alpha(\text{N})=2.48\times 10^{-8}$ 4; $\alpha(\text{IPF})=0.000443$ 7 B(M1)(W.u.)=0.112 +24-48; B(E2)(W.u.)=0.003 +19-3

Adopted Levels, Gammas (continued)

γ(⁶⁶Zn) (continued)

<u>E_i(level)</u>	<u>J_i^π</u>	<u>E_γ[‡]</u>	<u>I_γ[#]</u>	<u>E_f</u>	<u>J_f^π</u>	<u>Mult.</u>	<u>δ^α</u>	<u>α[†]</u>	<u>Comments</u>
4295.339	1 ⁺	3256.021 9	2.47 10	1039.2279	2 ⁺	M1+E2	1.5 2	0.000889 14	α(K)=2.70×10 ⁻⁵ 4; α(L)=2.66×10 ⁻⁶ 4; α(M)=3.81×10 ⁻⁷ 6; α(N+..)=0.000859 14 α(N)=1.552×10 ⁻⁸ 22; α(IPF)=0.000859 14 B(M1)(W.u.)=0.00071 +21-34; B(E2)(W.u.)=0.24 +6-11
		4295.187 10	100.0 8	0.0	0 ⁺				
4321.83		3282.5 ^b 2	100	1039.2279	2 ⁺				
4393.7	3 ⁻	1565.2 20	≈100	2826.69	3 ⁻				E _γ : reported only in ⁶⁶ Zn(α,α'γ). I _γ : from ⁶⁶ Zn(α,α'γ).
		3357.0 25	≈100	1039.2279	2 ⁺				E _γ : reported only in ⁶⁶ Zn(α,α'γ). I _γ : from ⁶⁶ Zn(α,α'γ).
4424	1	4424 6	100	0.0	0 ⁺				E _γ : from ⁶⁶ Zn(n,n'γ). E _γ =4426 in ⁶⁶ Zn(γ,γ'). I _γ : from ⁶⁶ Zn(n,n'γ).
4433	1 ⁻	4433 ^h 6	100	0.0	0 ⁺				E _γ ,I _γ : reported only in ⁶⁶ Zn(n,n'γ).
4461.409	1 ⁺	375.398 17	0.25 7	4085.983	1 ⁺				
		670.251 14	0.48 8	3791.123	1 ⁺				
		708.36 5	1.01 9	3753.01	4 ⁺				
		723.17 5	0.40 6	3738.207	+				
		885.00 5	0.22 6	3576.370	4 ⁺				
		929.68 3	0.53 7	3531.692	0 ⁺				
		954.12 7	0.52 8	3507.249	2 ⁺				
		1129.923 18	1.59 9	3331.441	2 ⁺				
		1232.480 15	6.5 22	3228.885	1 ⁺				
		1248.779 22	0.12 4	3212.582	2 ⁺				
		1356.320 15	14.3 22	3105.040	0 ⁺				
		1523.279 15	0.64 6	2938.074	2 ⁺				
		2088.985 13	1.3 3	2372.353	0 ⁺				
		2588.553 13	3.07 18	1872.7653	2 ⁺	M1+E2	0.35 27	0.000568 16	α(K)=3.86×10 ⁻⁵ 6; α(L)=3.81×10 ⁻⁶ 6; α(M)=5.47×10 ⁻⁷ 9; α(N+..)=0.000525 16 α(N)=2.22×10 ⁻⁸ 4; α(IPF)=0.000525 16 B(M1)(W.u.)=0.0022 +10-22; B(E2)(W.u.)=0.06 +10-6
		3422.040 8	100.0 7	1039.2279	2 ⁺	M1+E2	-0.06 2	0.000885 13	α(K)=2.44×10 ⁻⁵ 4; α(L)=2.41×10 ⁻⁶ 4; α(M)=3.46×10 ⁻⁷ 5; α(N+..)=0.000858 12 α(N)=1.408×10 ⁻⁸ 20; α(IPF)=0.000858 12 B(M1)(W.u.)=0.034 +15-34; B(E2)(W.u.)=0.017 +14-17
		4461.202 9	97.7 13	0.0	0 ⁺				
4497.6		3458.3 ^b 5	100	1039.2279	2 ⁺				
4609?	(1)	4609 ^h	100	0.0	0 ⁺				E _γ ,I _γ : reported only in ⁶⁶ Zn(γ,γ').
4622		4622 6	100	0.0	0 ⁺				E _γ ,I _γ : reported only in ⁶⁶ Zn(n,n'γ).
4635.3	(2)	2762 ^h	100	1872.7653	2 ⁺				E _γ ,I _γ : reported only in ⁶⁶ Zn(γ,γ').

Adopted Levels, Gammas (continued)

γ(⁶⁶Zn) (continued)

<u>E_i(level)</u>	<u>J_i^π</u>	<u>E_γ[‡]</u>	<u>I_γ[#]</u>	<u>E_f</u>	<u>J_f^π</u>	<u>Mult.</u>	<u>δ^α</u>	<u>α[†]</u>	<u>Comments</u>
4638.24	1	1106.53 24	77 24	3531.692	0 ⁺				
		1409.35 24	1.0×10 ² 5	3228.885	1 ⁺				
		2265.84 24	9×10 ¹ 4	2372.353	0 ⁺				
4675.6	1 ⁺	2802.8 5	100	1872.7653	2 ⁺				
4683	(1)	4685 ^h	100	0.0	0 ⁺				E _γ , I _γ : reported only in ⁶⁶ Zn(γ, γ').
4806.199	1 ⁺	1015.081 18	0.66 16	3791.123	1 ⁺				
		1135.47 9	0.25 5	3670.72	2 ⁺				
		1274.50 3	0.38 3	3531.692	0 ⁺				
		1298.95 7	0.205 24	3507.249	2 ⁺				
		1425.25 2	0.32 3	3380.944	1 ⁻				
		1577.308 20	0.21 4	3228.885	1 ⁺				
		1868.105 20	1.5 3	2938.074	2 ⁺				
		2026.016 25	0.14 4	2780.157	2 ⁺				
		2433.807 18	0.40 4	2372.353	0 ⁺				
		2933.358 9	11.45 16	1872.7653	2 ⁺	M1+E2	1.6 2	0.000762 12	α(K)=3.19×10 ⁻⁵ 5; α(L)=3.15×10 ⁻⁶ 5; α(M)=4.52×10 ⁻⁷ 7; α(N+..)=0.000727 12 α(N)=1.84×10 ⁻⁸ 3; α(IPF)=0.000727 12 B(M1)(W.u.)=0.0060 +17-24; B(E2)(W.u.)=2.9 +7-10
		3766.850 9	8.0 3	1039.2279	2 ⁺	M1+E2	0.11 4	0.001009 15	α(K)=2.11×10 ⁻⁵ 3; α(L)=2.08×10 ⁻⁶ 3; α(M)=2.98×10 ⁻⁷ 5; α(N+..)=0.000986 14 α(N)=1.212×10 ⁻⁸ 17; α(IPF)=0.000986 14 B(M1)(W.u.)=0.0069 +15-24; B(E2)(W.u.)=0.010 8
		4806.007 9	100.0 6	0.0	0 ⁺				E _γ : γ-ray, in ⁶⁶ Ga decay, often used for the efficiency calibration of germanium detectors.
4814.1	(7 ⁻)	738.4 ^c 3	100 ^c 9	4075.7	(6 ⁻)	(M1+E2) [@]	+0.11 2	0.000472 7	α(K)=0.000423 6; α(L)=4.24×10 ⁻⁵ 6; α(M)=6.08×10 ⁻⁶ 9; α(N+..)=2.45×10 ⁻⁷ 4 α(N)=2.45×10 ⁻⁷ 4 B(M1)(W.u.)=(0.06 5); B(E2)(W.u.)=(2.2 18)
		915.5 ^c 8	45 ^c 9	3898.3	5 ⁻				
4849.93	1 ⁺	1468.97 5	6.0 17	3380.944	1 ⁻				
		2977.08 4	100 10	1872.7653	2 ⁺				
		3810.59 5	40 4	1039.2279	2 ⁺				
4866.056	1 ⁺	1195.32 9	2.9 11	3670.72	2 ⁺				
		1433.63 4	5.9 12	3432.408	1 ⁻				
		1534.60 4	19 5	3331.441	2 ⁺				
		1927.96 4	7.2 24	2938.074	2 ⁺				
		2085.86 4	7 5	2780.157	2 ⁺				
		2993.21 3	100 10	1872.7653	2 ⁺				
		4865.87 4	8.8 7	0.0	0 ⁺				
4958.2	1 ⁺	3085.4 4	100	1872.7653	2 ⁺				

Adopted Levels, Gammas (continued)

γ(⁶⁶Zn) (continued)

<u>E_i(level)</u>	<u>J_i^π</u>	<u>E_γ[‡]</u>	<u>I_γ[#]</u>	<u>E_f</u>	<u>J_f^π</u>	<u>Mult.</u>	<u>δ^α</u>	<u>α[†]</u>	<u>Comments</u>
5005.8	1 ⁺	5005.6 3	100	0.0	0 ⁺				
5111.9	(8 ⁻)	860.8 ^c 8	11 ^c 3	4251.9	(7 ⁻)	(M1+E2)&	+0.21 15	0.000344 8	α(K)=0.000308 7; α(L)=3.08×10 ⁻⁵ 8; α(M)=4.42×10 ⁻⁶ 11; α(N+..)=1.78×10 ⁻⁷ 4 α(N)=1.78×10 ⁻⁷ 4
5207.3	(8 ⁺)	1036.0 ^c 3 954.2 ^c 5	100 ^c 3 82 ^c 16	4075.7 4251.9	(6 ⁻) (7 ⁻)	(E1)&		0.0001395 20	B(E1)(W.u.)<3.6×10 ⁻⁵ α(K)=0.0001252 18; α(L)=1.244×10 ⁻⁵ 18; α(M)=1.78×10 ⁻⁶ 3 α(N)=7.18×10 ⁻⁸ 10 δ(M2/E1)=-0.00 8. E _γ : doublet reported in ⁶⁴ Ni(α,2nγ).
		1025.8 ^c 5	100 ^c 11	4182.7	(6 ⁺)	(E2)@		0.000277 4	α(K)=0.000248 4; α(L)=2.49×10 ⁻⁵ 4; α(M)=3.56×10 ⁻⁶ 5; α(N+..)=1.427×10 ⁻⁷ 20 α(N)=1.427×10 ⁻⁷ 20 B(E2)(W.u.)<2.9 δ(M3/E2)=-0.04 6.
5464.4	(9 ⁻)	1212.5 ^c 4	100 ^c	4251.9	(7 ⁻)	(E2)@		0.000200 3	α(K)=0.0001697 24; α(L)=1.695×10 ⁻⁵ 24; α(M)=2.43×10 ⁻⁶ 4; α(N+..)=1.073×10 ⁻⁵ α(N)=9.76×10 ⁻⁸ 14; α(IPF)=1.063×10 ⁻⁵ 17 B(E2)(W.u.)=7 3 δ(M3/E2)=-0.04 4.
6292.6	(10 ⁺)	828 ^{ch} 1 1085.3 ^c 4	<22 ^c 100 ^c 9	5464.4 5207.3	(9 ⁻) (8 ⁺)	(E2)@		0.000243 4	α(K)=0.000218 3; α(L)=2.18×10 ⁻⁵ 3; α(M)=3.12×10 ⁻⁶ 5; α(N+..)=1.252×10 ⁻⁷ 18 α(N)=1.252×10 ⁻⁷ 18 B(E2)(W.u.)=13 +4-7 δ(M3/E2)=+0.04 6.
6419.0		1307.1 ^c 7	100 ^c	5111.9	(8 ⁻)				
7367.4	1	2732 3071 ^h 3293 ^h 3356 ^h 3485 3544 3793 ^h 3835	1.9 12 2.1 12 ≈0.5 0.4 2 2.1 7 1.6 5 ≈0.5 1.6 7	4635.3 4295.339 4075.7 4011.7 3882.424 3825.0 3576.370 3531.692	(2) 1 ⁺ (6 ⁻) (2) ⁺ 0 ⁺ 4 ⁺ 0 ⁺				
		4141	1.0 6	3226.2					E _γ : rounded-off value from level energy difference; 3840 from (γ,γ').

Adopted Levels, Gammas (continued)

$\gamma(^{66}\text{Zn})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\ddagger	$I_\gamma^\#$	E_f	J_f^π	Comments
7367.4	1	4262	8.1 20	3105.040	0 ⁺	
		4428	2.1 6	2938.074	2 ⁺	
		4587	3.8 12	2780.157	2 ⁺	E_γ : rounded off value from level energy difference; 4591 from (γ, γ') .
		4995 ^h	≤ 0.4	2372.353	0 ⁺	
		5495	14.3 7	1872.7653	2 ⁺	
		6326	≈ 0.3	1039.2279	2 ⁺	
		7368	100	0.0	0 ⁺	
7517.3		1224.7 ^c	100 ^c	6292.6	(10 ⁺)	
7693.3	1	4187	8 2	3507.249	2 ⁺	
		4263	25 3	3432.408	1 ⁻	
		4361	13 2	3331.441	2 ⁺	
		4452	7 2	3241.2?		
		4480	21 2	3212.582	2 ⁺	
		4587	8 1	3105.040	0 ⁺	
		4755	24 2	2938.074	2 ⁺	
		4930	<2	2762.8	(2)	I_γ : relative branching also reported as 8 in $^{66}\text{Zn}(\gamma, \gamma')$.
		5321	<3	2372.353	0 ⁺	
		5819	<2	1872.7653	2 ⁺	
		6654	42 1	1039.2279	2 ⁺	
		7693	100 1	0.0	0 ⁺	

[†] Additional information 2.

[‡] From ^{66}Ga ϵ decay, except as noted. Energies of γ 's depopulating levels above 7 MeV are from $^{66}\text{Zn}(\gamma, \gamma')$.

[#] Relative-photon branching from each level from ^{66}Ga ϵ decay, except as noted. Branchings for γ 's depopulating levels above 7 MeV are from $^{66}\text{Zn}(\gamma, \gamma')$.

[@] From $\gamma(\theta)$ in $^{64}\text{Ni}(\alpha, 2n\gamma)$ and RUL.

[&] From $\gamma(\theta)$ in $^{64}\text{Ni}(\alpha, 2n\gamma)$ and ΔJ^π .

^a From $\gamma(\theta)$ in $^{64}\text{Ni}(\alpha, 2n\gamma)$, except as noted.

^b From $^{66}\text{Zn}(n, n'\gamma)$.

^c From $^{64}\text{Ni}(\alpha, 2n\gamma)$.

^d From $^{66}\text{Zn}(p, p'\gamma)$.

^e From level energy differences in $^{65}\text{Cu}(p, \gamma)$. Uncertainty not given, but has been chosen by the evaluators to be 0.5 keV (an approximate upper limit).

^f From $^{66}\text{Zn}(p, p'\gamma)$. Uncertainties not given in the source data but are estimated as $\approx 2\text{-}20\%$.

^g From $^{65}\text{Cu}(p, \gamma)$.

^h Placement of transition in the level scheme is uncertain.


Adopted Levels, Gammas

Legend

Level Scheme

Intensities: Relative photon branching from each level

-----▶ γ Decay (Uncertain)



$^{66}_{30}\text{Zn}_{36}$


Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level

-----▶ γ Decay (Uncertain)



⁶⁶Zn₃₆


Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level

-----▶ γ Decay (Uncertain)



$^{66}_{30}\text{Zn}_{36}$


Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level

-----▶ γ Decay (Uncertain)




Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level

-----> γ Decay (Uncertain)



$^{66}_{30}\text{Zn}_{36}$


Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level

-----> γ Decay (Uncertain)



$^{66}_{30}\text{Zn}_{36}$