

Coulomb excitation

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	Huo Junde, Huang Xiaolong, J. K. Tuli		NDS 106,159 (2005)	1-Apr-2005

1998Si25: Ep=2.0-4.5MeV,measured I_γ,B(E2),Ge(Li).
 1975Th01: E(α)=3.20-4.86 MeV, measured E_γ, I_γ, γ(θ), γ-ray yields.
 1977An27: E(α)=7.5 MeV, E(¹²C)=30 MeV, E(¹⁴N)=35 MeV, E(¹⁶O)=39 MeV, γ-ray yields.
 1961Ho05: E(α)=3.3 MeV, measured T_{1/2} from pulsed beam technique.
 Others: 1956Te26, 1962Ri09, 1964Al27, 1964Al28, 1976AnZK.
 B(E2) values are taken from 1998Si25 or 1975Th01.

⁶⁷Zn Levels

E(level) [†]	J ^π [‡]	T _{1/2}	Comments
0.0	5/2 ⁻		
93.3 6	1/2 ⁻		B(E2)↑=0.0018 7(1998Si25)
184.5 3	3/2 ⁻	1.01 ns 10	B(E2)↑=0.0192 10(1998Si25) BE2=0.0190 14(1975Th01). T _{1/2} : from 1961Ho05.
393.6 3	3/2 ⁻		B(E2)↑=0.014 4(1998Si25) BE2=0.00049 3(1975Th01).
815.21 19	7/2 ⁻		B(E2)↑=0.0285 20(1998Si25) BE2=0.029 2(1975Th01).
870.9? 10	3/2,5/2,7/2		J=5/2, 7/2 from γ(θ) (1975Th01). B(E2)↑=0.04 2 (1975Th01)
888.0 6	5/2 ⁻		E(level): only observed by 1975Th01. J=3/2, 5/2, 7/2 from γ(θ) (1975Th01). B(E2)↑=0.0088 6(1998Si25) BE2=0.0086 6(1975Th01). J=5/2 strongly favored from γ(θ) (1975Th01) when combined with the γγ(θ) data of 1973Ba54.

[†] From Adopted Levels.

[‡] From Adopted Levels. Supporting arguments from this data set are indicated.

γ(⁶⁷Zn)

E _i (level)	J _i ^π	E _γ [†]	I _γ [‡]	E _f	J _f ^π	Mult.#	δ [#]	α [@]	Comments
93.3	1/2 ⁻	93.3		0.0	5/2 ⁻	E2		0.873	
184.5	3/2 ⁻	91.3	13.1 4	93.3	1/2 ⁻	M1+E2	+0.06 5	0.083 8	
		184.6	86.9 25	0.0	5/2 ⁻	M1+E2	0.34 4	0.0180 13	
393.6	3/2 ⁻	209.0	9.6 3	184.5	3/2 ⁻	M1+E2	-0.034 21	0.00913 6	
		300.3	70.3 21	93.3	1/2 ⁻	M1+E2	+0.20 8		
		393.6	20.1 6	0.0	5/2 ⁻				δ: -0.17 8 or -2.4 3 for M1+E2.
815.21	7/2 ⁻	421.6 2	0.70 8	393.6	3/2 ⁻				
		630.7 2	5.1 2	184.5	3/2 ⁻	E2			
		815.2 2	94.2 28	0.0	5/2 ⁻	M1+E2	+5.5 5		
870.9?	3/2,5/2,7/2	870.9	100	0.0	5/2 ⁻				
888.0	5/2 ⁻	494.4	23.8 11	393.6	3/2 ⁻	M1+E2	-0.14 3		
		703.4	4.8 4	184.5	3/2 ⁻				δ: -0.09 28 or +8.0 18 FOR (M1+E2).
		794.7	18.4 7	93.3	1/2 ⁻	E2(+M3)	-0.04 4		
		888.0	53.2 16	0.0	5/2 ⁻	M1+E2	+0.96 9		

Continued on next page (footnotes at end of table)

Coulomb excitation (continued) **$\gamma({}^{67}\text{Zn})$ (continued)**

† From [1975Th01](#) where uncertainties are given; rest are from level energy differences.

‡ Percent photon branching from each level ([1975Th01](#)).

From adopted gammas.

@ Total theoretical internal conversion coefficients, calculated using the BrIcc code ([2008Ki07](#)) with “Frozen Orbitals” approximation based on γ -ray energies, assigned multipolarities, and mixing ratios, unless otherwise specified.

Coulomb excitation**Level Scheme**

Intensities: % photon branching from each level

