

Adopted Levels, Gammas

Type	Author	Citation	History	Literature Cutoff Date
Full Evaluation	G. Gürdal, E. A. Mccutchan	NDS 136, 1 (2016)		1-Jul-2016

Q(β^-)=-6.22x10³ 5; S(n)=11532.5 16; S(p)=8523.0 15; Q(α)=-4087.7 11 2012Wa38
 S(2n)=19725.7 21; S(2p)=15132.9 11 (2012Wa38).

α : [Additional information 1](#).

⁷⁰Ge Levels

Cross Reference (XREF) Flags

A	⁷⁰ Ga β^- decay	J	⁶⁹ Ga(p, γ)	S	⁷⁰ Ge(p,p'),(pol p,p')
B	⁷⁰ As ϵ decay	K	⁶⁹ Ga(d,n)	T	⁷⁰ Ge(d,d')
C	⁷⁰ Zn 2 β^- decay	L	⁶⁹ Ga(α ,t)	U	⁷⁰ Ge(⁶ Li, ⁶ Li')
D	¹² C(⁶⁶ Zn, ⁸ Be γ)	M	⁶⁹ Ga(³ He,d)	V	⁷⁰ Ge(α , α')
E	⁴⁶ Ti(²⁸ Si,4p γ)	N	Coulomb excitation	W	⁷⁰ Ge(e,e')
F	⁶⁴ Ni(¹² C, α 2n γ)	O	⁷⁰ Ge(γ , γ')	X	⁷² Ge(p,t)
G	⁶⁵ Cu(⁷ Li,2n γ), ⁶⁰ Ni(¹² C,2p γ)	P	⁷⁰ Ge(pol γ , γ')	Y	⁷⁴ Se(d, ⁶ Li)
H	⁶⁶ Zn(⁶ Li,d)	Q	⁷⁰ Ge(n,n' γ)		
I	⁶⁸ Zn(α ,2n γ), ⁶⁷ Zn(α ,n γ)	R	⁷⁰ Ge(p,p' γ)		

E(level) [†]	J ^{π}	T _{1/2}	XREF	Comments
0.0 ^b	0 ⁺	stable	ABCDEFGHIJKLMN OPQRSTUVWXYZ	R=4.055 fm 8; where R is the rms value of charge distribution from (e,e').
1039.506 ^b 9	2 ⁺	1.31 ps 2	AB DEFG IJ LMN QRSTUVWXY	Q=+0.04 3 (2003Su01); μ =+0.91 5 T _{1/2} : from B(E2)=0.179 3 in Coulomb excitation. Others: 1.38 ps 8 from B(E2)=0.169 10 from (e,e'), 1.32 ps 14 from DSAM in ¹² C(⁶⁶ Zn, ⁸ Be), 1.3 ps 3 from DSAM in ⁶⁸ Zn(α ,2n γ), ⁶⁷ Zn(α ,n γ) and >5 ps in (⁷ Li,2n γ). J ^{π} : from 1039.49 γ E2 to 0 ⁺ . Q: from multiple Coulomb Excitation (2003Su01). μ : from weighted average of 0.88 8 (2013Gu23, from g-factor using TF), 0.90 16 (2007Bo41, from g-factor using TF), 0.86 24 (2006Le24, from g-factor using TF), 0.740 178 (1987La20, from g-factor using TF), 0.94 52 (1984Pa20, from g-factor using TF), 0.94 20 (1977Fa07, from g-factor using IMPAC). Others: 0.76 16 (1977Fa07), 1.18 58 (1969He11); the same data reanalyzed by 1974Hu01 gave μ =1.76 42; the same data reanalyzed by 1977Fa07 according to the latest understanding of the experiment and corrections μ = 0.94 20 (from g=0.47 10) (this value is included in the weighted average).
1215.621 ^e 15	0 ⁺	3.7 ns 2	AB DE GHIJKLMN QRSTUV XY	T _{1/2} : from electron spectrometer measurement with pulsed-cyclotron beam in (p,p' γ). Others: 2.9 ns 4 from ⁷⁰ Ga β^- decay; 4.8 ns 7 from B(E2) in Coulomb excitation. J ^{π} : L=0 from ⁷² Ge(p,t) and ⁶⁶ Zn(⁶ Li,d); E0 transition to g.s.. The level interpreted as deformed-intruder state (2003Su01).
1707.689 ^c 14	2 ⁺	1.94 ps 28	B DEFG IJ LMN QRSTUV X	Q=-0.07 4 (2003Su01); μ =+1.3 7 (2013Gu23) J ^{π} : from 1708 γ E2 transition to g.s. T _{1/2} : from DSAM in ¹² C(⁶⁶ Zn, ⁸ Be). Other: 1.1 ps +10-14 from (α ,2n γ), (α ,n γ), 4.2 ps +26-14 from Coulomb

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{70}Ge Levels (continued)

E(level) [†]	J^π	$T_{1/2}$	XREF	Comments
				excitation and > 7 ps in $^{65}\text{Cu}(^7\text{Li}, 2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C}, 2p\gamma)$.
2153.084 ^b 20	4 ⁺	0.76 ps 14	B DEFG IJ L N QR UV X	Q: from multiple Coulomb Excitation. μ : from g-factor measurements using TF in Coulomb excitation. Other: 0.8 12 from $^{12}\text{C}(^{66}\text{Zn}, ^8\text{Be})$ using TF. Q=+0.22 5 (2003Su01); μ =+1.7 8 $T_{1/2}$: from DSAM in $^{12}\text{C}(^{66}\text{Zn}, ^8\text{Be})$. Others: 4 ps 1 from DSAM in $^{65}\text{Cu}(^7\text{Li}, 2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C}, 2p\gamma)$ and 1.7 ps 4 from Coulomb excitation and 0.8 ps 2 from DSAM in $^{68}\text{Zn}(\alpha, 2n\gamma)$, $^{67}\text{Zn}(\alpha, n\gamma)$. J^π : from 1113.60 γ E2 to 2 ⁺ ; assumed E2 cascade member. Q: from multiple Coulomb Excitation (2003Su01). μ : from g-factor measurements using TF in Coulomb excitation (2013Gu23, 2007Bo41).
2156.744 ^e 21	2 ⁺		B D G J MN QRS	Q=+0.26 10 (2003Su01) J^π : 941.10 γ E2 to 0 ⁺ . Q: from multiple Coulomb Excitation.
2160 2307.0 5	0 ⁺	≤ 40 ps	J LM QRS X	$T_{1/2}$: centroid-shift time measurement in (p,p' γ). J^π : E0 transition to g.s.
2451.313 ^c 21	3 ⁺	1.7 [#] ps +10-3	B D FG IJ LM QRS	J^π : from 743.62 γ M1(+E2) to 2 ⁺ ; J = 3 from angular distribution and yield function in ($\alpha, 2n\gamma$).
2534.95 4	2 ⁺	0.6 [#] ps 2	B D IJ LM QRS X	J^π : L(p,t)=2; J^π not consistent with observed $\log ft=7.8$ from 4 ⁺ . $T_{1/2}$: Other: >0.4 ps from DSAM in (n,n' γ). $\mu=0.3$ 9 (2007Bo41)
2562.049 ^d 20	3 ⁻	0.50 ps 7	B DE G I LMN QRSTUVWX	$T_{1/2}$: weighted average of 0.55 ps 7 in $^{12}\text{C}(^{66}\text{Zn}, ^8\text{Be}\gamma)$ (from DSAM) and 0.4 ps 1 in $^{65}\text{Cu}(^7\text{Li}, 2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C}, 2p\gamma)$ (from DSAM). Others: 2.3 ps 5 from $^{65}\text{Cu}(^7\text{Li}, 2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C}, 2p\gamma)$. J^π : L(p,t)=3. B(E3)=0.073 10 in $^{70}\text{Ge}(e, e')$; 0.068 from Coulomb excitation. μ : from transient field method in Coulomb excitation (2007Bo04).
2806.25 ^c 3	4 ⁺	0.6 [#] ps 2	B DEFG IJ LM QRS V X	J^π : L(p,t)=4; 1098.54 γ E2 to 2 ⁺ . Discrepant with $L(\alpha, \alpha')=3$ for E=2800 keV 10 which may be a different level.
2887.4 7	0 ⁺		LM QRS X	J^π : from L(p,t)=0.
2945.0 10	2 ⁺		J LM QRS V X	J^π : from L(p,t)=2.
3046.439 20	3 ⁺		B IM QRS U X	J^π : from 889.72 γ D+Q to 2 ⁺ ; 893.59 γ D+Q to 4 ⁺ , $\log ft=5.75$ from 4 ⁺ parent.
3058.695 ^e 16	4 ⁺	1.4 [#] ps 3	B DE G I m QRST V	$T_{1/2}$: other: 1.0 ps 5 from DSAM in $^{65}\text{Cu}(^7\text{Li}, 2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C}, 2p\gamma)$. J^π : L(p,t)=4.
3105.7 7	(0 ⁺)		M QRS X	J^π : from excitation function in (p,p' γ).
3130 10			V	
3180.6 10	2 ⁺	0.015 ps 6	M QRS UV X	J^π : L(p,t)=2. $T_{1/2}$: from DSAM in (n,n' γ).
3194.2 6	4 ⁺		L S V X	J^π : L(p,t)=4; discrepant with $L(\alpha, \alpha')=(5)$ for E=3200 keV 10. Also, L(p,p')=4.

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{70}Ge Levels (continued)

E(level) [†]	J ^π	T _{1/2}	XREF	Comments
3240.5 10	1 ⁺		LM O RS	J ^π : 1 from $\gamma(\theta)$ in (γ,γ') and $\pi=+$ from L=1 in ($^3\text{He,d}$).
3294.79 8	3 ⁺ ,4 ⁺		B e QR X	J ^π : from 1587 γ to 2 ⁺ and L(p,p') = 4.
3296.98 ^b 3	6 ⁺	0.5 [#] ps 1	eFG I Q S	T _{1/2} : others: 2.6 ps 6 from DSAM in $^{65}\text{Cu}(^7\text{Li},2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C},2p\gamma)$.
3308				J ^π : from 1143.89 γ E2 to 4 ⁺ ; band assignment.
3314.5 7	1 ⁻		M O QRS UV X	J ^π : 1 from $\gamma(\theta)$ in (γ,γ') , $\pi=-$ from L(α,α')=1. B(E1) [†] : from $^{70}\text{Ge}(\gamma,\gamma')$.
3334.8 10	0 ⁺ to 3 ⁺		LM QRS X	J ^π : from L=1(+3) in ($^3\text{He,d}$) and from observed 2295.3 γ to 2 ⁺ .
3345 2			S	
3351 2			S	
3371.57 10	(3,4)	0.3 ps 2	B Q	J ^π : log ft=7.7 1 from 4 ⁺ in ε decay; J=5 unlikely because of 2333 γ to 2 ⁺ level. T _{1/2} : from DSAM in (n,n' γ).
3416.32 ^d 4	5 ⁻	13.7 [@] ps 10	DE G I M Q S X	J ^π : L(p,t)=5. T _{1/2} : Other: > 14 ps in $^{65}\text{Cu}(^7\text{Li}, 2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C},2p\gamma)$.
3423 2	(2 ⁺)		V X	J ^π : from L(p,t)=(2).
3428 2	5 ⁻		STU	J ^π : from L(α,α')=5.
3432 2	3 ⁻		M S V X	J ^π : from L(α,α')=3.
3456 2	4 ⁺		S	J ^π : from L=4 in (p,p') and ($^6\text{Li},^6\text{Li}'$).
3466? 6			M	
3482.3 5	1 ⁺ ,2 ⁺ ,3 ⁺		M RS X	J ^π : from L=1+3 in ($^3\text{He,d}$).
3488.276 21	(3,4 ⁺)		B QRS	J ^π : log ft=6.0 1 from 4 ⁺ in ε decay; J=4 ⁻ and 5 unlikely because of 2449 γ to 2 ⁺ level.
3517? 6			M	
3540?			T	
3562.7 6			M RS X	
3570.44 7	(3) ⁻		B M S UV	J ^π : from 2531.7 γ to 2 ⁺ ; 1471.24 γ to 4 ⁺ ; L=4(+2) in ($^3\text{He,d}$).
3580.7 10	4 ⁺	0.6 ps 2	Q S	J ^π : L(α,α')=4. T _{1/2} : from DSAM in (n,n' γ).
3590.3 5			M RS	
3631.5 10	(2) ⁺	0.5 ps 1	LM QRS X	J ^π : from L(p,t)=(2); $\pi=+$ from L($^3\text{He,d}$)=1. T _{1/2} : from DSAM in (n,n' γ).
3637 10	0 ⁺		V	J ^π : L(α,α')=0.
3666.78 ^d 6	6 ⁻	35 [@] ps 3	E G I M Q S	J ^π : J=6 from $\gamma(\theta)$ in $^{46}\text{Ti}(^{28}\text{Si},4p\gamma)$, π from 250 γ M1(+E2) to 5 ⁻ . T _{1/2} : others: 40 ps 8 in $^{68}\text{Zn}(\alpha,2n\gamma)$, $^{67}\text{Zn}(\alpha,n\gamma)$ and 74 ps 6 in (HI,xn γ).
3669.4 ^c 10	(5 ⁺)	1 [#] ps 1	F I M	J ^π : 1218 γ to 3 ⁺ ; band assignment.
3675.76 7	4 ⁺		B L RS V	J ^π : from L(α,α')=4.
3683 3	0 ⁺			J ^π : from L(p,t)=0.
3687 3	1 ⁺ ,2 ⁺ ,3 ⁺		M S	J ^π : L=1+3 in ($^3\text{He,d}$).
3708.5 9			M RS	
3733 3	1 ⁺ ,2 ⁺ ,3 ⁺		M	J ^π : L=1+3 in ($^3\text{He,d}$).
3740 3	0 ⁺		S X	J ^π : L(p,t)=0.
3753.2 ^c 4	6 ⁺	1.6 [#] ps 5	EFG I S	J ^π : from stretched 946.7 γ E2 to 4 ⁺ , L(p,p')=(6); band assignment.
3776 2	3 ⁻		S V X	J ^π : from L(α,α')=3.
3782 2	2 ⁺		M S	XREF: M(3775). J ^π : from L(p,t)=2.

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{70}Ge Levels (continued)

E(level) [†]	J ^π	T _{1/2}	XREF		Comments
3850 3			LM	S	
3856 2	(2) ⁻			S	J ^π : from L=4+2(+0) in (³ He,d).
3870 2	3 ⁻			S V X	J ^π : from L(α,α')=3.
3890 3	1 ⁺ ,2 ⁺ ,3 ⁺		M	S	J ^π : from L=1+3 in (³ He,d).
3895.2 10	1		L O	X	J ^π : from 3895.1γ D to 0 ⁺ .
3900.6 7	(4 ⁻ ,5,6,7 ⁻)		E		J ^π : from 234γ to 6 ⁻ , 484γ to 5 ⁻ .
3903.9 7	+		M	RS V	J ^π : L=1+3 in (³ He,d), L(p,p')=(0). E(level): from (p,p'γ).
3911 3				S	
3928 3	4 ⁺			S X	J ^π : from L(p,t)=4.
3941? 10				S	
3955.11 ^d 8	7 ⁻	17.0 [@] ps 10	E G I	Q S V	J ^π : J=7 from γ(θ) in (α,2nγ), π from 288γ M1+(E2) to 6 ⁻ .
3964 3	(2) ⁻		LM	V	J ^π : from L=4+2(+0) in (³ He,d).
3976 3	1 ⁺ ,2 ⁺ ,3 ⁺			S	J ^π : from L(α,α')=2.
3990 3			M	S	
4003 2				RS V	
4024 3	4 ⁺		M	X	J ^π : from L(p,t)=4.
4037 3	(4 ⁺)			S V	J ^π : from L(α,α')=(4).
4053.3 10			E		
4054 3			M	S	
4061 2	1 ⁺ ,2 ⁺ ,3 ⁺			S	J ^π : from L=1+3 in (³ He,d).
4080 3	1 ⁺ ,2 ⁺ ,3 ⁺		M		J ^π : from L=1+3 in (³ He,d).
4086 3	4 ⁺			V X	J ^π : L(p,t)=4.
4096.1 20	3 ⁻			RS	J ^π : from L(α,α')=3.
4101.45 5	3 ⁻ ,4 ⁻		B	S	J ^π : from 688γ to 5 ⁻ and L(p,p')=3.
4103.5 ^{±e} 5	6 ⁺		G		J ^π : 1295γ Q to 4 ⁺ ; band assignment in ⁶⁵ Cu(⁷ Li, 2nγ), ⁶⁰ Ni(¹² C, 2pγ) (2010Su05).
4119 3			M	S	
4131 2	2 ⁻		M	S	J ^π : from L=4+2+0 in (³ He,d).
4144.7 20	1 ⁻			RS V	J ^π : from L(α,α')=1.
4155 2	1 ⁺ ,2 ⁺ ,3 ⁺		M	S	J ^π : L=1+3 in (³ He,d).
4166 3				S	
4180 3	2 ⁺			S X	J ^π : L(p,t)=2.
4203.5 ^b 4	8 ⁺	8 [@] ps 2	EFG I	M S	J ^π : from 906.6γ E2 to 6 ⁺ ; band assignment.
4212 3	3 ⁺ ,4 ⁺ ,5 ⁺		M	S	J ^π : from L(p,p')=4.
4226 3	2 ⁺			S X	J ^π : from L(p,t)=2.
4238 3	1 ⁺ ,2 ⁺ ,3 ⁺		M	S X	J ^π : from L=1+3 in (³ He,d).
4243.11 15			B		
4261 10	2 ⁺			S X	J ^π : from L(p,t)=2.
4268 10	5 ⁻			V	J ^π : from L(α,α')=5.
4282 10	3 ⁺ ,4 ⁺ ,5 ⁺			S	J ^π : from L(p,p')=4.
4287 3	1 ⁺ ,2 ⁺ ,3 ⁺		M		J ^π : from L(³ He,d)=1+3.
4299.3 3	7 ⁻	3 [@] ps 1	E G I	S V	J ^π : from L(α,α')=7.
4330 3			LM	S V X	L=4+2(+0) in (³ He,d); L(α,α')=(3+5), L(p,t)=0.
4352 3	(2) ⁻		M P	X	J ^π : from L=4+2(+0) in (³ He,d).
4356.7 7	1 ^{(-)&}			P	B(E1)↑=0.0023 4 B(E1) from ⁷⁰ Ge(pol γ, γ').
4357 10	+			S X	L(p,t)=2 at 4357 20; L(p,p')=4 at 4357 10.
4365 10	(3 ⁻)			S V	J ^π : from L(α,α')=(3) for 4373 10.
4378 10				S	
4391 3	1 ⁺ ,2 ⁺ ,3 ⁺		M	S	J ^π : from L(³ He,d)=1+3.
4409 10	4 ⁺			S V	J ^π : from L(α,α')=4.
4419 3	2 ⁻ ,3 ⁻ ,4 ⁻		M	S	J ^π : from L(³ He,d)=4+2.

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{70}Ge Levels (continued)

E(level) [†]	J ^π	T _{1/2}	XREF	Comments
4431.4 4	8 ⁺	0.4 [#] ps 2	EFG I	J ^π : from 1134.6γ E2 to 6 ⁺ and yield function in $^{68}\text{Zn}(\alpha,2n\gamma)$.
4447.5 8	1 ⁻ &		M P	B(E1)↑=0.0036 7 B(E1) from $^{70}\text{Ge}(\text{pol } \gamma, \gamma')$.
4448 2	2 ⁺		S V X	J ^π : from L(p,t)=2.
4473 2	4 ⁺		M S V X	J ^π : from L(p,t)=4. Other: L(α,α')=(3+5) is discrepant.
4520 3	2 ⁻ ,3 ⁻ ,4 ⁻		M S	J ^π : from L(^3He ,d)=4+2.
4520.9 8	1 ⁻		P	B(E1)↑<0.0005 B(E1) from $^{70}\text{Ge}(\text{pol } \gamma, \gamma')$.
4534 10	(4 ⁺)		S V	J ^π : from L(α,α')=(4).
4539 3	0 ⁺		X	J ^π : from L(p,t)=0.
4546 10			S	
4552.1 10	(8)	104 [@] ps +70-35	I	J ^π : from 253γ to 7 ⁺ and 1253.2γ to 6 ⁺ .
4555 3			M S	
4574 3			M	
4577.18 15	(3,4 ⁺)		B S	J ^π : log ft=6.4 from 4 ⁺ in ε decay; J=5 unlikely because of 2421γ to 2 ⁺ level.
4606 10			S	
4613 3	1 ⁺ ,2 ⁺ ,3 ⁺		M S	J ^π : from L(^3He ,d)=1+3.
4629 3	(4 ⁺)		S V X	J ^π : L(α,α')=(4).
4642 3	(2) ⁻		M S	J ^π : L(^3He ,d)=4+2(+0).
4657 10			V	E(level): multiplet.
4675.39 21	(3,4 ⁺)		B S	J ^π : log ft=6.2 2 from 4 ⁺ in ε decay; J=5 or 4 ⁻ are unlikely because of 2968.1γ to 2 ⁺ level.
4687 2	(2) ⁻		LM S	J ^π : L=4+2(+0) in (^3He ,d).
4707 10			S X	
4716 10	(2 ⁺)		S V	J ^π : from L(p,t)=(2).
4727 10			S	
4736 3			M	
4768 3	(2) ⁻		M	J ^π : L(^3He ,d)=4+2(+0).
4775 10	(4 ⁺)		V	J ^π : from L(α,α')=(4).
4790.6 19	1 ⁽⁻⁾ &		P	
4810 10	3 ⁻		V	J ^π : from L(α,α')=3.
4820.2 ^c 11	(8 ⁺)		F	J ^π : 1067γ to 6 ⁺ ; band assignment.
4851.9 4	(8 ⁻) ^a	>3 [#] ps	E G I M	J ^π : from (M1+E2) γ to 7 ⁻ .
4877 3	2 ⁻		M	J ^π : L=4+2+0 in (^3He ,d).
4886.6 13	1&		P	
4905 3	3 ⁻		M	J ^π : from L(α,α')=3.
4908.1 ^d 10	(9 ⁻) ^a		E	
4915 10			V	
4935 3	1 ⁻		X	J ^π : from L(p,t)=1.
4940 10	3 ⁻		V	J ^π : from L(α,α')=3.
4943 3	(2) ⁻		M	J ^π : L(^3He ,d)=4+2(+0).
4979 3	(2) ⁻		M	J ^π : L=4+2(+0) in (^3He ,d).
4985.0 10			I	
5008 3	2 ⁻		M V	J ^π : from L=4+2+0 in (^3He ,d); L(α,α')=(3).
5024 3	2 ⁺		X	J ^π : from L(p,t)=2.
5040 10	(3 ⁻)		I M V	J ^π : L(α,α')=(3).
5048.4 10	(4 ⁻)		I M	J ^π : from L(^3He ,d)=4+2+0; 1381.gy to 2 ⁻ .
5050 3	0 ⁺		L X	J ^π : from L(p,t)=0.
5078 3	1 ⁺ ,2 ⁺ ,3 ⁺		M V	J ^π : L=1+3 in (^3He ,d).
5102 3	1 ⁺ ,2 ⁺ ,3 ⁺		M	J ^π : L=1+3 in (^3He ,d).
5113 10	(3 ⁻)		V	J ^π : L(α,α')=(3).

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{70}Ge Levels (continued)

E(level) [†]	J ^π	T _{1/2}	XREF	Comments
5129.6 7	1- $\&$		P	B(E1) \uparrow =0.0029 8 B(E1) from $^{70}\text{Ge}(\text{pol } \gamma, \gamma')$.
5145 3	(3 ⁻)		M V	J ^π : L(α, α')=(3); discrepant with L($^3\text{He}, d$)=(4+2+0).
5184 3	0 ⁺			J ^π : L(p,t)=0.
5195 10	(4 ⁺)			J ^π : L(α, α')=(4).
5222.3 14			E	
5227 10	(3 ⁻)			J ^π : L(α, α')=(3).
5242.7 ^b 11	10 ⁺		EFG I	J ^π : from 1039.2 γ Q to 8 ⁺ ; band assignment.
5263.4 8	1(-) $\&$		P	B(E1) \uparrow =0.0022 4 B(E1) from $^{70}\text{Ge}(\text{pol } \gamma, \gamma')$.
5265.82 14			B	
5290 3	0 ⁺			J ^π : L(p,t)=0.
5299.2 4	9(-)		E G I	J ^π : from 1344.1 γ (E2) to 7 ⁻ , 1273 γ from 11(-); inconsistent with (6,7,8) from $\gamma(\theta)$ in ($^7\text{Li}, 2n\gamma$).
5338 3	0 ⁺			J ^π : L(p,t)=0.
5370.11 5			B	
5403 3	0 ⁺			J ^π : L(p,t)=0.
5410 3				
5435.5 ^{‡e} 11	8 ⁺		G	J ^π : from 1332 γ Q to 6 ⁺ ; band assignment in $^{65}\text{Cu}(^7\text{Li}, 2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C}, 2p\gamma)$ (2010Su05).
5441 3	(2 ⁺)			J ^π : L(p,t)=(2).
5465.3 8	1- $\&$		P	B(E1) \uparrow =0.0023 4 B(E1) from $^{70}\text{Ge}(\text{pol } \gamma, \gamma')$.
5467 3	0 ⁺			J ^π : L(p,t)=0.
5512.5 10	1(-) $\&$		P	B(E1) \uparrow =0.0019 3 B(E1) from $^{70}\text{Ge}(\text{pol } \gamma, \gamma')$.
5539.7 5	(10)	5 ns 2	FG I	T _{1/2} : from electronic timing in $^{68}\text{Zn}(\alpha, 2n\gamma)$, $^{67}\text{Zn}(\alpha, n\gamma)$. J ^π : T _{1/2} of this level suggests that the 1108 γ to 8 ⁺ may be M2, suggesting an assignment of J ^π =10 ⁻ . However, level is assigned as side band member based on 8 ⁺ in $^{64}\text{Ni}(^{12}\text{C}, \alpha 2n\gamma)$, suggesting an assignment of J ^π =10 ⁺ . Other: 9 ⁺ in $^{65}\text{Cu}(^7\text{Li}, 2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C}, 2p\gamma)$.
5552.5 [‡] 5	9(-)		G	J ^π : 1253.2 γ Q to 7 ⁻ .
5876.9 7	1(-) $\&$		P	B(E1) \uparrow =0.0014 4 B(E1) from $^{70}\text{Ge}(\text{pol } \gamma, \gamma')$.
5989.7 7	1(+) $\&$		P	
6006.9 11			E	
6160.1 ^d 14	(11 ⁻) ^a		E	
6297.0 14	1 $\&$		P	
6362.8 8	1 $\&$		P	
6549.1 14			E	
6572.2 11	11(-)		G	J ^π : from 1273 γ Q to 9(-).
6587.7 8	1(+) $\&$		P	
6604.2 11			E	
6636.6 15	1 $\&$		P	
6702.5 13	1(-) $\&$		P	B(E1) \uparrow =0.0027 5 B(E1) from $^{70}\text{Ge}(\text{pol } \gamma, \gamma')$.
6716.8 ^b 15	12 ⁺		FG	J ^π : Q 1474 γ to 10 ⁺ ; band assignment.
6779.7 11	(12)		F	
6786.1 ^d 17	(13 ⁻) ^a		E	
7306.3 8	1(+) $\&$		P	

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{70}Ge Levels (continued)

E(level) [†]	J ^π	XREF	Comments
7426.0 8	1 ^{(-)&}	P	B(E1)↑=0.0022 4 B(E1) from $^{70}\text{Ge}(\text{pol } \gamma, \gamma')$.
7619.7 15	(14)	F	J ^π : Q 840γ to (12).
7753.5 10	1 ^{(-)&}	P	B(E1)↑=0.0026 6 B(E1) from $^{70}\text{Ge}(\text{pol } \gamma, \gamma')$.
7767.8 ^b 18	14 ⁺	F	J ^π : Q 1051γ to 12 ⁺ , band assignment.
8058.1 ^d 20	(15 ⁻) ^a	E	
8245.7 18	(16)	F	J ^π : Q 626γ to (14).
8283.7 15	1 ^{(+)&}	P	
8878.5 14	1 ^{&}	P	
9423.7 21	(18)	F	J ^π : Q 1178γ to (16).
9619.2 ^d 22	(17 ⁻) ^a	E	
10269.7 23	(20)	F	J ^π : 846γ to (18).
11336.2 ^d 25	(19 ⁻) ^a	E	
13173 ^d 3	(21 ⁻) ^a	E	

[†] From a least-squares fit for levels connected by γ 's and from reaction data sets for others.

[‡] This level was only reported in [2010Su05](#) in $^{65}\text{Cu}(^7\text{Li},2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C},2p\gamma)$.

From DSAM in $^{68}\text{Zn}(\alpha,2n\gamma)$, $^{67}\text{Zn}(\alpha,n\gamma)$.

@ From RDM in $^{68}\text{Zn}(\alpha,2n\gamma)$, $^{67}\text{Zn}(\alpha,n\gamma)$ ([1982Cl02](#)).

& From mult in (pol γ, γ').

^a From ($^{28}\text{Si},4p\gamma$) based on DCO, and level cascades.

^b Band(A): sequence based on g.s..

^c Band(B): sequence based on 2⁺, 1707.7 keV level.

^d Band(C): sequence based on 3⁻, 2562.05 keV level.

^e Band(D): sequence based on 0⁺, 1215.62 keV level.

Adopted Levels, Gammas (continued)

$\gamma(^{70}\text{Ge})$										
$E_i(\text{level})$	J_i^π	E_γ †	I_γ †	E_f	J_f^π	Mult.	δ	α	$I_{(\gamma+ce)}$	Comments
1039.506	2 ⁺	1039.513 10	100	0.0	0 ⁺	E2 ^d		3.23×10 ⁻⁴		$\alpha(\text{K})=0.000289$ 4; $\alpha(\text{L})=2.96\times 10^{-5}$ 5; $\alpha(\text{M})=4.41\times 10^{-6}$ 7; $\alpha(\text{N})=2.88\times 10^{-7}$ 4 B(E2)(W.u.)=20.8 4 E_γ : from ⁷⁰ Ga β^- decay.
1215.621	0 ⁺	176.115 13	100 9	1039.506	2 ⁺	E2 ^f		0.0894		$\alpha(\text{K})=0.0790$ 11; $\alpha(\text{L})=0.00902$ 13; $\alpha(\text{M})=0.001337$ 19; $\alpha(\text{N})=7.73\times 10^{-5}$ 11 B(E2)(W.u.)=48 7 E_γ : from ⁷⁰ Ga β^- decay.
		1215.8 ^b		0.0	0 ⁺	E0 ^f			1.00 4	$I_{(\gamma+ce)}$: for 100 transitions of 176 γ from (p,p' γ) (1985Pa15).
1707.689	2 ⁺	492.09 5	4.9 4	1215.621	0 ⁺	E2		0.00247		$\alpha(\text{K})=0.00220$ 3; $\alpha(\text{L})=0.000232$ 4; $\alpha(\text{M})=3.45\times 10^{-5}$ 5; $\alpha(\text{N})=2.20\times 10^{-6}$ 3 B(E2)(W.u.)=16 3 Mult.: from RUL and decay pattern.
		668.21 4	100 6	1039.506	2 ⁺	M1+E2	-3.6 +11-6	9.80×10 ⁻⁴ 2		$\alpha(\text{K})=0.000875$ 21; $\alpha(\text{L})=9.08\times 10^{-5}$ 22; $\alpha(\text{M})=1.35\times 10^{-5}$ 4; $\alpha(\text{N})=8.74\times 10^{-7}$ 20 B(E2)(W.u.)=64 11; B(M1)(W.u.)=0.0015 9 Mult., δ : from RUL, δ and decay pattern. δ as evaluated by 1977Kr17;
		1707.61 2	79.2 23	0.0	0 ⁺	E2		2.87×10 ⁻⁴		$\alpha(\text{K})=0.0001011$ 15; $\alpha(\text{L})=1.025\times 10^{-5}$ 15; $\alpha(\text{M})=1.529\times 10^{-6}$ 22; $\alpha(\text{N})=1.007\times 10^{-7}$ 15 B(E2)(W.u.)=0.50 8 Mult.: from angular distribution in ⁶⁸ Zn(α ,2n γ), ⁶⁷ Zn(α ,n γ) and RUL.
2153.084	4 ⁺	445.6 10	0.7 4	1707.689	2 ⁺	[E2]		0.00338 6		$\alpha(\text{K})=0.00301$ 5; $\alpha(\text{L})=0.000318$ 5; $\alpha(\text{M})=4.74\times 10^{-5}$ 8; $\alpha(\text{N})=3.00\times 10^{-6}$ 5 B(E2)(W.u.)=17 10
		1113.60 4	100 6	1039.506	2 ⁺	E2		2.77×10 ⁻⁴		$\alpha(\text{K})=0.000247$ 4; $\alpha(\text{L})=2.52\times 10^{-5}$ 4; $\alpha(\text{M})=3.76\times 10^{-6}$ 6; $\alpha(\text{N})=2.46\times 10^{-7}$ 4 B(E2)(W.u.)=25 5 Mult.: from ⁶⁸ Zn(α ,2n γ), ⁶⁷ Zn(α ,n γ) and RUL. $\delta = -0.1$ 2 in ⁶⁸ Zn(α ,2n γ), ⁶⁷ Zn(α ,n γ) gives a large B(M3) which is excluded by RUL.
2156.744	2 ⁺	450.4 5	4.7 24	1707.689	2 ⁺	E2		0.00327		$\alpha(\text{K})=0.00291$ 5; $\alpha(\text{L})=0.000308$ 5; $\alpha(\text{M})=4.58\times 10^{-5}$ 7; $\alpha(\text{N})=2.90\times 10^{-6}$ 5 Mult.: from (p,p' γ). Other: D from ⁶⁵ Cu(⁷ Li,2n γ), ⁶⁰ Ni(¹² C,2p γ).
		941.10 4	62 3	1215.621	0 ⁺	E2		4.09×10 ⁻⁴		$\alpha(\text{K})=0.000366$ 6; $\alpha(\text{L})=3.76\times 10^{-5}$ 6; $\alpha(\text{M})=5.60\times 10^{-6}$ 8; $\alpha(\text{N})=3.65\times 10^{-7}$ 6 Mult.: from (p,p' γ).
		1117.28 4	100 6	1039.506	2 ⁺	E2		2.75×10 ⁻⁴		$\alpha(\text{K})=0.000245$ 4; $\alpha(\text{L})=2.50\times 10^{-5}$ 4;

Adopted Levels, Gammas (continued)

$\gamma(^{70}\text{Ge})$ (continued)										
$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult.	δ	α	$I_{(\gamma+ce)}$	Comments
2156.744	2 ⁺	2156.65 6	17.1 9	0.0	0 ⁺	[E2]		4.60×10 ⁻⁴		$\alpha(\text{M})=3.73\times 10^{-6}$ 6; $\alpha(\text{N})=2.44\times 10^{-7}$ 4 Mult.: from (p,p' γ).
2307.0	0 ⁺	599.1 ^b	82 [@] 7	1707.689	2 ⁺	E2 ^b		1.36×10 ⁻³		$\alpha(\text{K})=6.55\times 10^{-5}$ 10; $\alpha(\text{L})=6.62\times 10^{-6}$ 10; $\alpha(\text{M})=9.88\times 10^{-7}$ 14; $\alpha(\text{N})=6.52\times 10^{-8}$ 10
		1091.3 ^b		1215.621	0 ⁺	E0 ^f			0.013 2	$\alpha(\text{K})=0.001218$ 17; $\alpha(\text{L})=0.0001270$ 18; $\alpha(\text{M})=1.89\times 10^{-5}$ 3; $\alpha(\text{N})=1.216\times 10^{-6}$ 17 B(E2)(W.u.)>4.8 $I_{(\gamma+ce)}$: for 100 transitions of 1268 γ from (p,p' γ) (1985Pa15).
		1267.5 ^b	100 [@] 7	1039.506	2 ⁺	E2 ^b		2.28×10 ⁻⁴		$\alpha(\text{K})=0.000185$ 3; $\alpha(\text{L})=1.89\times 10^{-5}$ 3; $\alpha(\text{M})=2.82\times 10^{-6}$ 4; $\alpha(\text{N})=1.85\times 10^{-7}$ 3 B(E2)(W.u.)>0.14
		2307.1 ^b		0.0	0 ⁺	E0 ^f			0.040 9	$I_{(\gamma+ce)}$: for 100 transitions of 1268 γ from (p,p' γ) and includes pair production (1985Pa15).
2451.313	3 ⁺	294.60 16 297.88 8 743.62 4	0.37 15 2.4 4 100 7	2156.744 2 ⁺ 2153.084 4 ⁺ 1707.689 2 ⁺		M1(+E2)	+0.04 8	5.78×10 ⁻⁴ 9		$\alpha(\text{K})=0.000517$ 8; $\alpha(\text{L})=5.28\times 10^{-5}$ 8; $\alpha(\text{M})=7.89\times 10^{-6}$ 12; $\alpha(\text{N})=5.20\times 10^{-7}$ 8 B(E2)(W.u.)<0.5; B(M1)(W.u.)=0.022 +14-5 Mult., δ : D+Q from ⁶⁵ Cu(⁷ Li,2n γ), ⁶⁰ Ni(¹² C,2p γ), M1+(E2) from RUL. Other: $\delta=+3.5$ 9 from $\gamma(\theta)$ in (n,n' γ).
		1411.86 4	39.3 23	1039.506	2 ⁺	M1+E2	-2.2 +5-3	2.18×10 ⁻⁴ 4		$\alpha(\text{K})=0.0001463$ 22; $\alpha(\text{L})=1.487\times 10^{-5}$ 22; $\alpha(\text{M})=2.22\times 10^{-6}$ 4; $\alpha(\text{N})=1.460\times 10^{-7}$ 22 B(E2)(W.u.)=0.79 +48-17; B(M1)(W.u.)=0.00022 +15-10 Mult.: D+Q from $\gamma(\theta)$ in (n,n' γ); M1+E2 from RUL. δ : from $\gamma(\theta)$ in (n,n' γ).
2534.95	2 ⁺	827.24 10 1319.6	32 4 9.5 10	1707.689 2 ⁺ 1215.621 0 ⁺						I_γ : from (n,n' γ); $I_\gamma(827)/I_\gamma(1495)=76$ 10/100 10, in (n,n' γ) which ratio is different from the adopted I_γ from ⁷⁰ As ε decay.
		1495.43 5	100 8	1039.506	2 ⁺	M1+E2	-0.75	2.15×10 ⁻⁴		$\alpha(\text{K})=0.0001274$ 18; $\alpha(\text{L})=1.291\times 10^{-5}$ 18; $\alpha(\text{M})=1.93\times 10^{-6}$ 3; $\alpha(\text{N})=1.273\times 10^{-7}$ 18 B(E2)(W.u.)=1.9 7; B(M1)(W.u.)=0.0050 18 Mult., δ : D+Q from $\gamma(\theta)$ in (n,n' γ); M1+E2 from RUL.
2562.049	3 ⁻	1522.55 2	100	1039.506	2 ⁺	E1+M2 ^d	-0.11 ^d 10	3.42×10 ⁻⁴		$\alpha(\text{K})=6.7\times 10^{-5}$ 6; $\alpha(\text{L})=6.8\times 10^{-6}$ 6; $\alpha(\text{M})=1.01\times 10^{-6}$ 8; $\alpha(\text{N})=6.6\times 10^{-8}$ 6

Adopted Levels, Gammas (continued)

$\gamma(^{70}\text{Ge})$ (continued)								
$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult.	α	Comments
2806.25	4 ⁺	653.15 6	11.8 14	2153.084	4 ⁺	(M1)	7.66×10 ⁻⁴	B(E1)(W.u.)=0.00022 4 δ : other: -0.11 4 from $\gamma(\theta)$ in (n,n' γ); 0.02 5 from $^{65}\text{Cu}(^7\text{Li},2n\gamma)$. $\alpha(\text{K})=0.000685$ 10; $\alpha(\text{L})=7.01\times 10^{-5}$ 10; $\alpha(\text{M})=1.048\times 10^{-5}$ 15; $\alpha(\text{N})=6.90\times 10^{-7}$ 10 B(M1)(W.u.)=0.014 5 Mult.: D from $^{65}\text{Cu}(^7\text{Li},2p\gamma)$, $^{60}\text{Ni}(^{12}\text{C},2p\gamma)$ based on $\Delta J = 0$ dipole transition; $\Delta\pi$ =no from level scheme.
		1098.54 4	100 6	1707.689	2 ⁺	E2	2.84×10 ⁻⁴	$\alpha(\text{K})=0.000254$ 4; $\alpha(\text{L})=2.60\times 10^{-5}$ 4; $\alpha(\text{M})=3.88\times 10^{-6}$ 6; $\alpha(\text{N})=2.54\times 10^{-7}$ 4 B(E2)(W.u.)=31 11 Mult.: from $^{68}\text{Zn}(\alpha,2n\gamma)$, $^{67}\text{Zn}(\alpha,n\gamma)$ and RUL; $\delta=-0.2$ 2 in $^{68}\text{Zn}(\alpha,2n\gamma)$, $^{67}\text{Zn}(\alpha,n\gamma)$ gives a large B(M3) which is excluded by RUL.
2887.4	0 ⁺	730.8 [‡]	100 [‡] 10	2156.744	2 ⁺			
		1179.5 [‡]	100 [‡] 15	1707.689	2 ⁺			
2945.0	2 ⁺	1237.3 ^b	100 ^b	1707.689	2 ⁺			
3046.439	3 ⁺	239.90 10	1.1 3	2806.25	4 ⁺			
		595.11 4	100 6	2451.313	3 ⁺	[M1+E2]		
		889.72 4	14.0 9	2156.744	2 ⁺	M1+E2 ^h		
		893.50 4	10.0 5	2153.084	4 ⁺	M1+E2 ^h		
		1338.76 4	48 3	1707.689	2 ⁺	M1+E2 ^h		
3058.695	4 ⁺	2006.87 3	14.8 5	1039.506	2 ⁺	M1+E2 ^h		
		252.46 4	16.3 10	2806.25	4 ⁺	[M1+E2]		
		496.74 4	15.3 10	2562.049	3 ⁻	[E1]	7.15×10 ⁻⁴	$\alpha(\text{K})=0.000639$ 9; $\alpha(\text{L})=6.52\times 10^{-5}$ 10; $\alpha(\text{M})=9.72\times 10^{-6}$ 14; $\alpha(\text{N})=6.31\times 10^{-7}$ 9 B(E1)(W.u.)=0.00015 4 Mult.: D from $\gamma(\theta)$ in $^{65}\text{Cu}(^7\text{Li},2p\gamma)$, $^{60}\text{Ni}(^{12}\text{C},2p\gamma)$.
		607.34 4	26.2 15	2451.313	3 ⁺	M1+(E2) ^h		
		901.95 5	5.9 4	2156.744	2 ⁺	[E2]	4.54×10 ⁻⁴	$\alpha(\text{K})=0.000406$ 6; $\alpha(\text{L})=4.17\times 10^{-5}$ 6; $\alpha(\text{M})=6.22\times 10^{-6}$ 9; $\alpha(\text{N})=4.05\times 10^{-7}$ 6 B(E2)(W.u.)=1.00 23
		905.61 2	67 4	2153.084	4 ⁺	[M1+E2]		
		1350.90 6	2.8 3	1707.689	2 ⁺	[E2]	2.21×10 ⁻⁴	$\alpha(\text{K})=0.0001618$ 23; $\alpha(\text{L})=1.647\times 10^{-5}$ 23; $\alpha(\text{M})=2.46\times 10^{-6}$ 4; $\alpha(\text{N})=1.614\times 10^{-7}$ 23 B(E2)(W.u.)=0.063 16
		2019.16 2	100.0 25	1039.506	2 ⁺	E2	4.02×10 ⁻⁴	$\alpha(\text{K})=7.38\times 10^{-5}$ 11; $\alpha(\text{L})=7.46\times 10^{-6}$ 11; $\alpha(\text{M})=1.114\times 10^{-6}$ 16; $\alpha(\text{N})=7.35\times 10^{-8}$ 11 B(E2)(W.u.)=0.30 7 Mult.: from $^{68}\text{Zn}(\alpha,2n\gamma)$, $^{67}\text{Zn}(\alpha,n\gamma)$ and RUL. M3 is ruled out because $\delta=+0.2$ 2 from $^{68}\text{Zn}(\alpha,2n\gamma)$, $^{67}\text{Zn}(\alpha,n\gamma)$ gives a large B(M3).
3105.7	(0 ⁺)	1397.9 [‡]	33 [‡] 10	1707.689	2 ⁺			
		2066.3 [‡]	100 [‡] 7	1039.506	2 ⁺			
3180.6	2 ⁺	2141.1 [‡]	100	1039.506	2 ⁺			

Adopted Levels, Gammas (continued)

$\gamma(^{70}\text{Ge})$ (continued)									
$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult.	δ	α	Comments
3240.5	1 ⁺	3240.4	100	0.0	0 ⁺	M1		8.29×10 ⁻⁴	$\alpha(\text{K})=3.21\times 10^{-5}$ 5; $\alpha(\text{L})=3.23\times 10^{-6}$ 5; $\alpha(\text{M})=4.82\times 10^{-7}$ 7; $\alpha(\text{N})=3.19\times 10^{-8}$ 5 E_γ : from (γ, γ'). Mult.: D from $\gamma(\theta)$ in (γ, γ'), $\Delta\pi$ =yes from level scheme.
3294.79	3 ⁺ , 4 ⁺	760.2 5 1587.17 12 2255.16 11	8.×10 ¹ 4 100 13 44 6	2534.95 2 ⁺ 1707.689 2 ⁺ 1039.506 2 ⁺					
3296.98	6 ⁺	490 ^g 1143.89& 2	0.80 ^g 20 100 ^g	2806.25 4 ⁺ 2153.084 4 ⁺		[E2] E2		2.62×10 ⁻⁴	$\alpha(\text{K})=0.000232$ 4; $\alpha(\text{L})=2.37\times 10^{-5}$ 4; $\alpha(\text{M})=3.53\times 10^{-6}$ 5; $\alpha(\text{N})=2.31\times 10^{-7}$ 4 B(E2)(W.u.)=34 7 Mult.: From angular distribution in ⁶⁸ Zn($\alpha, 2n\gamma$), ⁶⁷ Zn($\alpha, n\gamma$) and RUL. $\delta = 0.0$ 2 from $\gamma(\theta)$.
3314.5	1 ⁻	2274.6‡ 3314.8		1039.506 2 ⁺ 0.0 0 ⁺		E1		1.41×10 ⁻³	$\alpha(\text{K})=2.09\times 10^{-5}$ 3; $\alpha(\text{L})=2.10\times 10^{-6}$ 3; $\alpha(\text{M})=3.14\times 10^{-7}$ 5; $\alpha(\text{N})=2.07\times 10^{-8}$ 3 E_γ : from (γ, γ'). Mult.: D from $\gamma(\theta)$ in (γ, γ'), $\Delta\pi$ =yes from level scheme.
3334.8	0 ⁺ to 3 ⁺	2295.3‡	100	1039.506 2 ⁺					
3371.57	(3,4)	1218.57 11 2331.59 24	100 21 26 6	2153.084 4 ⁺ 1039.506 2 ⁺					
3416.32	5 ⁻	357.72 5	59 4	3058.695 4 ⁺		E1 ^d		1.68×10 ⁻³	$\alpha(\text{K})=0.001499$ 21; $\alpha(\text{L})=0.0001533$ 22; $\alpha(\text{M})=2.28\times 10^{-5}$ 4; $\alpha(\text{N})=1.475\times 10^{-6}$ 21 B(E1)(W.u.)=0.000146 16 Mult.: From ⁶⁸ Zn($\alpha, 2n\gamma$), ⁶⁷ Zn($\alpha, n\gamma$) and RUL. $\delta = -0.06$ 3 in ⁶⁸ Zn($\alpha, 2n\gamma$), ⁶⁷ Zn($\alpha, n\gamma$) gives a large B(M2). δ : Other: 0.00 4 in ⁶⁵ Cu(⁷ Li, 2n γ), ⁶⁰ Ni(¹² C, 2p γ).
		854.6 4	97 6	2562.049 3 ⁻		E2		5.20×10 ⁻⁴	$\alpha(\text{K})=0.000464$ 7; $\alpha(\text{L})=4.78\times 10^{-5}$ 7; $\alpha(\text{M})=7.13\times 10^{-6}$ 10; $\alpha(\text{N})=4.64\times 10^{-7}$ 7 B(E2)(W.u.)=2.00 21 Mult.: From ⁶⁸ Zn($\alpha, 2n\gamma$), ⁶⁷ Zn($\alpha, n\gamma$) and RUL. $\delta = 0.02$ 5 in ⁶⁸ Zn($\alpha, 2n\gamma$), ⁶⁷ Zn($\alpha, n\gamma$) gives a large B(M3).
		1263.09& 6	100 6	2153.084 4 ⁺		E1(+M2) ^d	-0.05 ^d 5	1.90×10 ⁻⁴ 4	$\alpha(\text{K})=8.98\times 10^{-5}$ 24; $\alpha(\text{L})=9.07\times 10^{-6}$ 25; $\alpha(\text{M})=1.35\times 10^{-6}$ 4; $\alpha(\text{N})=8.90\times 10^{-8}$ 25 B(E1)(W.u.)=5.5×10 ⁻⁶ 6; B(M2)(W.u.)=0.040 4
3488.276	(3,4 ⁺)	953.30 7 1036.99 4	11.3 11 64 5	2534.95 2 ⁺ 2451.313 3 ⁺					

Adopted Levels, Gammas (continued)

$\gamma(^{70}\text{Ge})$ (continued)									
$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult.	δ	α	Comments
3488.276	(3,4 ⁺)	1331.58 7 1335.28 10 1780.52 2 2448.82 9	10.0 9 8.1 9 100.0 22 7.7 5	2156.744 2153.084 1707.689 1039.506	2 ⁺ 4 ⁺ 2 ⁺ 2 ⁺				
3570.44	(3) ⁻	1417.24 7 2531.7 2	100 10 6.0 20	2153.084 1039.506	4 ⁺ 2 ⁺				
3580.7	4 ⁺	1427.6 [‡]	100	2153.084	4 ⁺				
3631.5	(2) ⁺	2591.9 [‡]	100	1039.506	2 ⁺				
3666.78	6 ⁻	250.46 ^{&} 5	100	3416.32	5 ⁻	M1(+E2) ^d	0.03 ^d +2-5	0.00727	$\alpha(\text{K})=0.00648$ 10; $\alpha(\text{L})=0.000678$ 10; $\alpha(\text{M})=0.0001013$ 15; $\alpha(\text{N})=6.62\times 10^{-6}$ 10 B(E2)(W.u.)=0.85 7; B(M1)(W.u.)=0.040 3 δ : Other: +0.05 2 in ⁶⁵ Cu(⁷ Li,2n γ), ⁶⁰ Ni(¹² C,2p γ).
3669.4	(5 ⁺)	1218.1 ^{&}	100	2451.313	3 ⁺				
3675.76	4 ⁺	1523.2 7 2636.20 7	100 19 43.6 19	2153.084 1039.506	4 ⁺ 2 ⁺				
3753.2	6 ⁺	946.7 [#] 4	100 [#]	2806.25	4 ⁺	E2		4.03 $\times 10^{-4}$	$\alpha(\text{K})=0.000360$ 5; $\alpha(\text{L})=3.70\times 10^{-5}$ 6; $\alpha(\text{M})=5.52\times 10^{-6}$ 8; $\alpha(\text{N})=3.60\times 10^{-7}$ 5 B(E2)(W.u.)=27 9 Mult.: Q from $\gamma(\theta)$ in (α ,2n γ), M2 excluded by comparison to RUL.
3895.2	1	3895.1	100	0.0	0 ⁺	D		1.07 $\times 10^{-3}$	$\alpha(\text{K})=2.40\times 10^{-5}$ 4; $\alpha(\text{L})=2.41\times 10^{-6}$ 4; $\alpha(\text{M})=3.60\times 10^{-7}$ 5; $\alpha(\text{N})=2.39\times 10^{-8}$ 4 E_γ : from (γ,γ'). Mult.: from $\gamma(\theta)$ in (γ,γ'). E_γ : From ⁴⁶ Ti(²⁸ Si, 4p γ). E_γ : From ⁴⁶ Ti(²⁸ Si, 4p γ).
3900.6	(4 ⁻ ,5,6,7 ⁻)	234 484		3666.78 3416.32	6 ⁻ 5 ⁻				
3955.11	7 ⁻	288.33 ^{&} 5 658.1 ^{&} 4	100 ^{&} 10 19 ^{&} 4	3666.78 3296.98	6 ⁻ 6 ⁺	M1(+E2) ^d E1(+M2)	0.01 ^d 3 +0.02 5	0.00512 3.67 $\times 10^{-4}$ 10	$\alpha(\text{K})=0.00457$ 7; $\alpha(\text{L})=0.000476$ 7; $\alpha(\text{M})=7.12\times 10^{-5}$ 11; $\alpha(\text{N})=4.66\times 10^{-6}$ 7 B(E2)(W.u.)=0.081 12; B(M1)(W.u.)=0.045 7 $\alpha(\text{K})=0.000329$ 9; $\alpha(\text{L})=3.34\times 10^{-5}$ 9; $\alpha(\text{M})=4.98\times 10^{-6}$ 13; $\alpha(\text{N})=3.25\times 10^{-7}$ 9 B(E1)(W.u.)=1.3 $\times 10^{-5}$ 3; B(M2)(W.u.)=0.056 13 Mult., δ : D+Q from ⁶⁵ Cu(⁷ Li,2n γ), $\Delta\pi$ = yes from level scheme.
4053.3		1247 [#]	100 [#]	2806.25	4 ⁺				
4101.45	3 ⁻ ,4 ⁻	688 [@] 1045 [@] 1295.24 6		3416.32 3058.695 2806.25	5 ⁻ 4 ⁺ 4 ⁺				
			100 11			Q			Mult.: Q from $\gamma(\theta)$ in ⁶⁵ Cu(⁷ Li, 2n γ), ⁶⁰ Ni(¹² C, 2p γ).

Adopted Levels, Gammas (continued)

$\gamma(^{70}\text{Ge})$ (continued)										
$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult.	δ	α	Comments	
4101.45	3 ⁻ ,4 ⁻	1539.29 20 1944.21 16 1948.35 11	31 9 25 4 59 5	2562.049 3 ⁻ 2156.744 2 ⁺ 2153.084 4 ⁺	3 ⁻ 2 ⁺ 4 ⁺					
4103.5	6 ⁺	688 ^c 1045 ^c 1295 ^c 1948 ^c	7.3 ^c 6 100 ^c 9 14.8 ^c 13 5.5 ^c 6	3416.32 5 ⁻ 3058.695 4 ⁺ 2806.25 4 ⁺ 2156.744 2 ⁺	5 ⁻ 4 ⁺ 4 ⁺ 2 ⁺	Q			Mult.: from angular distribution of oriented nuclei (ADO) radios in ⁶⁵ Cu(⁷ Li, 2n γ), ⁶⁰ Ni(¹² C, 2p γ) (2010Su05). E γ : 1948 γ populates 2 ⁺ state according to level scheme given in 2010Su05. Placement of 1948 γ is questionable due to ΔJ .	
4203.5	8 ⁺	450 ^g 906.6 [@] 4	2.9 ^g 6 100 ^g 10	3753.2 6 ⁺ 3296.98 6 ⁺	6 ⁺ 6 ⁺	[E2] E2		4.48 $\times 10^{-4}$	$\alpha(K)=0.000401$ 6; $\alpha(L)=4.12\times 10^{-5}$ 6; $\alpha(M)=6.14\times 10^{-6}$ 9; $\alpha(N)=4.00\times 10^{-7}$ 6 B(E2)(W.u.)=6.7 17 Mult.: from angular distribution and linear-polarization data in ⁶⁸ Zn(α ,2n γ), ⁶⁷ Zn(α ,n γ) and RUL. $\delta=-0.2$ 2 is ⁶⁸ Zn(α ,2n γ), ⁶⁷ Zn(α ,n γ) gives a large B(M3) value, M3 is not possible because of RUL.	
4243.11		1196.66 15	100	3046.439 3 ⁺	3 ⁺					
4299.3	7 ⁻	344.1 ^{&} 4 1002.4 ^{&} 4	73 ^{&} 15 100 ^{&} 20	3955.11 7 ⁻ 3296.98 6 ⁺	7 ⁻ 6 ⁺	M1(+E2) E1+M2	0.1 3 0.11 2	0.0034 6 1.59 $\times 10^{-4}$ 4	$\alpha(K)=0.0030$ 6; $\alpha(L)=0.00031$ 6; $\alpha(M)=4.7\times 10^{-5}$ 9; $\alpha(N)=3.1\times 10^{-6}$ 6 B(E2)(W.u.)=10 4; B(M1)(W.u.)=0.08 3 Mult., δ : D+Q from angular distribution data in ⁶⁵ Cu(⁷ Li,2n γ), $\Delta\pi$ = no from level scheme. $\alpha(K)=0.000142$ 3; $\alpha(L)=1.44\times 10^{-5}$ 4; $\alpha(M)=2.15\times 10^{-6}$ 5; $\alpha(N)=1.41\times 10^{-7}$ 3 B(E1)(W.u.)=8.E-5 4 Mult., δ : D+Q from angular distribution data in ⁶⁵ Cu(⁷ Li,2n γ), $\Delta\pi$ = yes from level scheme.	
4356.7	1 ⁽⁻⁾	4356.6 ^a 7	100	0.0 0 ⁺	0 ⁺	E1 ^a		0.00182	$\alpha(K)=1.480\times 10^{-5}$ 21; $\alpha(L)=1.483\times 10^{-6}$ 21; $\alpha(M)=2.21\times 10^{-7}$ 3; $\alpha(N)=1.463\times 10^{-8}$ 21	
4431.4	8 ⁺	677 1134.6 [@] 4	3.4 10 100 9	3753.2 6 ⁺ 3296.98 6 ⁺	6 ⁺ 6 ⁺	[E2] E2		2.66 $\times 10^{-4}$	$\alpha(K)=0.000236$ 4; $\alpha(L)=2.42\times 10^{-5}$ 4; $\alpha(M)=3.60\times 10^{-6}$ 5; $\alpha(N)=2.36\times 10^{-7}$ 4 B(E2)(W.u.)=44 22 Mult.: from angular distribution in ⁶⁸ Zn(α ,2n γ), ⁶⁷ Zn(α ,n γ) and RUL. $\delta=-0.1$ 2 in ⁶⁸ Zn(α ,2n γ), ⁶⁷ Zn(α ,n γ) gives large B(M3). RUL rules out M3.	
4447.5	1 ⁻	4447.3 ^a 8	100	0.0 0 ⁺	0 ⁺	E1 ^a		0.00185	$\alpha(K)=1.443\times 10^{-5}$ 21; $\alpha(L)=1.446\times 10^{-6}$ 21; $\alpha(M)=2.16\times 10^{-7}$ 3; $\alpha(N)=1.426\times 10^{-8}$ 20	
4520.9	1 ⁻	4520.7 ^a 8	100	0.0 0 ⁺	0 ⁺	E1 ^a		0.00188	$\alpha(K)=1.414\times 10^{-5}$ 20; $\alpha(L)=1.417\times 10^{-6}$ 20; $\alpha(M)=2.11\times 10^{-7}$ 3; $\alpha(N)=1.398\times 10^{-8}$ 20	

Adopted Levels, Gammas (continued)

γ(⁷⁰Ge) (continued)

<u>E_i(level)</u>	<u>J_i^π</u>	<u>E_γ[†]</u>	<u>I_γ[†]</u>	<u>E_f</u>	<u>J_f^π</u>	<u>Mult.</u>	<u>δ</u>	<u>α</u>	<u>Comments</u>
4552.1	(8)	252.8&	100	4299.3	7 ⁻				
4577.18	(3,4 ⁺)	2419.88 24 2424.41 20	86 15 100 15	2156.744 2153.084	2 ⁺ 4 ⁺				
4675.39	(3,4 ⁺)	2521.8 3 2968.1 3	49 13 100 15	2153.084 1707.689	4 ⁺ 2 ⁺				
4790.6	1 ⁽⁻⁾	4790.4 ^a 19	100	0.0	0 ⁺	E1 ^a		0.00196	α(K)=1.317×10 ⁻⁵ 19; α(L)=1.319×10 ⁻⁶ 19; α(M)=1.97×10 ⁻⁷ 3; α(N)=1.302×10 ⁻⁸ 19
4820.2	(8 ⁺)	1067 ^g	100	3753.2	6 ⁺				
4851.9	(8 ⁻)	896.8& 4	100	3955.11	7 ⁻	(M1+E2) ^e	0.4 ^e 2	3.98×10 ⁻⁴ 11	α(K)=0.000356 10; α(L)=3.64×10 ⁻⁵ 11; α(M)=5.43×10 ⁻⁶ 15; α(N)=3.58×10 ⁻⁷ 10 B(E2)(W.u.)<4.9; B(M1)(W.u.)<0.01 δ: Other: +1.1 3 in ⁶⁵ Cu(⁷ Li,2nγ), ⁶⁰ Ni(¹² C,2nγ).
4886.6	1	4886.4 ^a 13	100	0.0	0 ⁺	D ^a			
4908.1	(9 ⁻)	953 [#]	100 [#]	3955.11	7 ⁻				
4985.0		1029.9&	100	3955.11	7 ⁻				
5048.4	(4 ⁻)	1381.6&	100	3666.78	6 ⁻				
5129.6	1 ⁻	5129.4 ^a 7	100	0.0	0 ⁺	E1 ^a		0.00207	α(K)=1.212×10 ⁻⁵ 17; α(L)=1.214×10 ⁻⁶ 17; α(M)=1.81×10 ⁻⁷ 3; α(N)=1.198×10 ⁻⁸ 17
5222.3		1169 [#]	100 [#]	4053.3					
5242.7	10 ⁺	1039.2&	100	4203.5	8 ⁺	E2		3.23×10 ⁻⁴	α(K)=0.000289 4; α(L)=2.96×10 ⁻⁵ 5; α(M)=4.41×10 ⁻⁶ 7; α(N)=2.88×10 ⁻⁷ 4 Mult.: Q from γ(θ) in ⁶⁵ Cu(⁷ Li, 2nγ), ⁶⁰ Ni(¹² C, 2pγ), assumed E2 band member.
5263.4	1 ⁽⁻⁾	5263.2 ^a 8	100	0.0	0 ⁺	E1 ^a		0.00211	α(K)=1.175×10 ⁻⁵ 17; α(L)=1.177×10 ⁻⁶ 17; α(M)=1.756×10 ⁻⁷ 25; α(N)=1.162×10 ⁻⁸ 17
5265.82		2219.34 14	100	3046.439	3 ⁺				
5299.2	9 ⁽⁻⁾	1344.1@ 4	100	3955.11	7 ⁻	(E2)			Mult.: from γ(θ) in ⁶⁵ Cu(⁷ Li,2nγ), ⁶⁰ Ni(¹² C,2nγ).
5370.11		1881.67 5 2325.42 18	100 6 21 2	3488.276 3046.439	(3,4 ⁺) 3 ⁺				
5435.5	8 ⁺	1332 ^c	100 ^c	4103.5	6 ⁺	Q			Mult.: from angular distribution of oriented nuclei (ADO) radios in ⁶⁵ Cu(⁷ Li, 2nγ), ⁶⁰ Ni(¹² C, 2pγ) (2010Su05).
5465.3	1 ⁻	5465.1 ^a 8	100	0.0	0 ⁺	E1 ^a		0.00216	α(K)=1.124×10 ⁻⁵ 16; α(L)=1.125×10 ⁻⁶ 16; α(M)=1.678×10 ⁻⁷ 24; α(N)=1.111×10 ⁻⁸ 16
5512.5	1 ⁽⁻⁾	5512.3 ^a 10	100	0.0	0 ⁺	E1 ^a		0.00217	α(K)=1.112×10 ⁻⁵ 16; α(L)=1.114×10 ⁻⁶ 16; α(M)=1.661×10 ⁻⁷ 24; α(N)=1.099×10 ⁻⁸ 16
5539.7	(10)	1108.3@ 4	100@	4431.4	8 ⁺	Q ^b		5.54×10 ⁻⁴	α(K)=0.000495 7; α(L)=5.10×10 ⁻⁵ 8; α(M)=7.62×10 ⁻⁶ 11; α(N)=5.03×10 ⁻⁷ 7 Mult.: from R(DCO) in ⁶⁴ Ni(¹² C,α2nγ). From T _{1/2}

Adopted Levels, Gammas (continued)

$\gamma(^{70}\text{Ge})$ (continued)									
$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult.	α	Comments	
								considerations, M2 is more likely. However, assignment to a band structure in $^{64}\text{Ni}(^{12}\text{C},\alpha 2n\gamma)$ suggests E2 character. Other: D in $^{65}\text{Cu}(^7\text{Li},2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C},2p\gamma)$.	
5552.5	9 ⁽⁻⁾	1253.2 [@]	4	100	4299.3	7 ⁻	Q	2.30×10 ⁻⁴	$\alpha(\text{K})=0.000190$ 3; $\alpha(\text{L})=1.94\times 10^{-5}$ 3; $\alpha(\text{M})=2.89\times 10^{-6}$ 4; $\alpha(\text{N})=1.89\times 10^{-7}$ 3 Mult.: Q from angular distribution of oriented nuclei (ADO) ratio in $^{65}\text{Cu}(^7\text{Li},2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C},2p\gamma)$.
5876.9	1 ⁽⁻⁾	5876.6 ^a	7	100	0.0	0 ⁺	E1 ^a	0.00227	$\alpha(\text{K})=1.031\times 10^{-5}$ 15; $\alpha(\text{L})=1.032\times 10^{-6}$ 15; $\alpha(\text{M})=1.540\times 10^{-7}$ 22; $\alpha(\text{N})=1.019\times 10^{-8}$ 15
5989.7	1 ⁽⁺⁾	5989.4 ^a	7	100	0.0	0 ⁺	M1 ^a	1.66×10 ⁻³	$\alpha(\text{K})=1.272\times 10^{-5}$ 18; $\alpha(\text{L})=1.276\times 10^{-6}$ 18; $\alpha(\text{M})=1.90\times 10^{-7}$ 3; $\alpha(\text{N})=1.262\times 10^{-8}$ 18
6006.9		1155 [#]		100 [#]	4851.9	(8 ⁻)			
6160.1	(11 ⁻)	1252 [#]		100 [#]	4908.1	(9 ⁻)			
6297.0	1	6296.7 ^a	14	100	0.0	0 ⁺	D ^a		
6362.8	1	6362.5 ^a	8	100	0.0	0 ⁺	D ^a		
6549.1		1641 [#]		100 [#]	4908.1	(9 ⁻)			
6572.2	11 ⁽⁻⁾	1273 [@]		100	5299.2	9 ⁽⁻⁾	Q		Mult.: from Angular Distribution of Oriented nuclei (ADO) ratio in $^{65}\text{Cu}(^7\text{Li}, 2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C}, 2p\gamma)$ and level scheme.
6587.7	1 ⁽⁺⁾	6587.4 ^a	8	100	0.0	0 ⁺	M1 ^a		
6604.2		1305 [#]		100 [#]	5299.2	9 ⁽⁻⁾			
6636.6	1	6636.3 ^a	15	100	0.0	0 ⁺	D ^a		
6702.5	1 ⁽⁻⁾	6702.2 ^a	13	100	0.0	0 ⁺	E1 ^a		
6716.8	12 ⁺	1474 ^g		100	5242.7	10 ⁺	E2		Mult.: Q in $^{65}\text{Cu}(^7\text{Li}, 2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C}, 2p\gamma)$ and $^{64}\text{Ni}(^{12}\text{C},\alpha 2n\gamma)$, assumed E2 from placement in band structure.
6779.7	(12)	1240 ^g		100	5539.7	(10)	Q		Mult.: from R(DCO) in $^{64}\text{Ni}(^{12}\text{C},\alpha 2n\gamma)$.
6786.1	(13 ⁻)	626 [#]		100 [#]	6160.1	(11 ⁻)			
7306.3	1 ⁽⁺⁾	7305.9 ^a	8	100	0.0	0 ⁺	M1 ^a		
7426.0	1 ⁽⁻⁾	7425.6 ^a	8	100	0.0	0 ⁺	E1 ^a		
7619.7	(14)	840 ^g		100	6779.7	(12)	Q		Mult.: from R(DCO) in $^{64}\text{Ni}(^{12}\text{C},\alpha 2n\gamma)$.
7753.5	1 ⁽⁻⁾	7753.0 ^a	10	100	0.0	0 ⁺	E1 ^a		
7767.8	14 ⁺	1051 ^g		100	6716.8	12 ⁺	E2		Mult.: Q from R(DCO) in $^{64}\text{Ni}(^{12}\text{C},\alpha 2n\gamma)$, E2 from assumed band structure.
8058.1	(15 ⁻)	1272 [#]		100 [#]	6786.1	(13 ⁻)			
8245.7	(16)	626 ^g		100	7619.7	(14)	Q		Mult.: from R(DCO) in $^{64}\text{Ni}(^{12}\text{C},\alpha 2n\gamma)$.
8283.7	1 ⁽⁺⁾	8283.2 ^a	15	100	0.0	0 ⁺	M1 ^a		
8878.5	1	8877.9 ^a	14	100	0.0	0 ⁺	D ^a		
9423.7	(18)	1178 ^g		100	8245.7	(16)	Q		Mult.: from R(DCO) in $^{64}\text{Ni}(^{12}\text{C},\alpha 2n\gamma)$.
9619.2	(17 ⁻)	1561 [#]		100 [#]	8058.1	(15 ⁻)			
10269.7	(20)	846 ^g		100	9423.7	(18)			
11336.2	(19 ⁻)	1717 [#]		100 [#]	9619.2	(17 ⁻)			
13173	(21 ⁻)	1837 [#]		100 [#]	11336.2	(19 ⁻)			

Adopted Levels, Gammas (continued)

$\gamma(^{70}\text{Ge})$ (continued)

† From ^{72}As ε decay, unless otherwise stated.

‡ From $(n,n'\gamma)$.

From $^{46}\text{Ti}(^{28}\text{Si}, 4p\gamma)$.

@ From $^{65}\text{Cu}(^7\text{Li}, 2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C}, 2p\gamma)$.

& From $^{68}\text{Zn}(\alpha, 2n\gamma)$, $^{67}\text{Zn}(\alpha, n\gamma)$.

^a From $^{70}\text{Ge}(\text{pol } \gamma, \gamma')$.

^b From $(p, p'\gamma)$.

^c from $^{65}\text{Cu}(^7\text{Li}, 2n\gamma)$, $^{60}\text{Ni}(^{12}\text{C}, 2p\gamma)$ (2010Su05).

^d From angular distribution and linear-polarization data in $^{68}\text{Zn}(\alpha, 2n\gamma)$, $^{67}\text{Zn}(\alpha, n\gamma)$.

^e From angular distribution in $^{68}\text{Zn}(\alpha, 2n\gamma)$, $^{67}\text{Zn}(\alpha, n\gamma)$ and RUL.

^f From internal conversion data in $(p, p'\gamma)$.


^g From $^{64}\text{Ni}(^{12}\text{C}, \alpha 2n\gamma)$.

^h From $\gamma(\theta)$ in ^{70}As ε decay, cases of D+Q with large, non-zero values for δ have been assumed to be M1+E2 in character.

Adopted Levels, Gammas

Level Scheme


Intensities: Relative photon branching from each level



Adopted Levels, Gammas

Level Scheme (continued)


Intensities: Relative photon branching from each level



Adopted Levels, Gammas

Level Scheme (continued)

Intensities: Relative photon branching from each level




⁷⁰Ge₃₈


Adopted Levels, Gammas

Level Scheme (continued)

Intensities: Relative photon branching from each level




⁷⁰Ge₃₈


Adopted Levels, GammasBand(A): Sequence based
on g.s. $^{70}_{32}\text{Ge}_{38}$

Adopted Levels, Gammas (continued)

Band(C): Sequence based
on 3^- , 2562.05 keV
level



Band(D): Sequence based
on 0^+ , 1215.62 keV
level

 $^{70}_{32}\text{Ge}_{38}$