

Adopted Levels, Gammas

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	Ameenah R. Farhan, Balraj Singh		NDS 110,1917 (2009)	30-Jun-2009

Q(β^-)=-3574.4; S(n)=10497.74 17; S(p)=10398.6 18; Q(α)=-6028.38 18 2012Wa38
 Note: Current evaluation has used the following Q record -3574.4 10497.7317 10398.418-6028.4 5 2009AuZZ,2003Au03.
 S(2n)=17916.59 18, s(2p)=18390.90 19 (2009AuZZ,2003Au03). Values in 2003Au03 are within \approx 0.1 keV of those in 2009AuZZ.
 Additional information 1.

Mass measurements: 1985E101, 1982Zu04, 1977De20.

Nuclear structure calculations: 2008Yo07 (high-spin levels, B(E2), shell-model); 2008Ah03 (levels, B(E2), g factor, projected shell model).

⁷⁸Se(e,e): 1988Kh02, 1987Ku21, 1986Kh07.

See ⁷⁷Se(n,n),(n, γ):resonances dataset for 38 resonances between 41.2 eV to 3.91 keV.

⁷⁸Se Levels

Cross Reference (XREF) Flags

A	⁷⁸ As β^- decay (90.7 min)	H	⁷⁷ Se(n, γ) E=112.0 eV	O	⁷⁸ Se(p,p' γ),($\alpha,\alpha'\gamma$)
B	Muonic atom	I	⁷⁷ Se(n, γ) E=211.6 eV	P	⁷⁸ Se(α,α')
C	⁷⁸ Br ϵ decay (6.45 min)	J	⁷⁷ Se(n, γ) E=340.8 eV	Q	⁷⁸ Se(d,d')
D	⁷⁶ Ge($\alpha,2n\gamma$)	K	⁷⁷ Se(n, γ) E=864.0 eV	R	Coulomb excitation
E	⁷⁶ Ge(¹⁶ O, ¹⁴ C)	L	⁷⁷ Se(d,p)	S	⁸⁰ Se(p,t)
F	⁷⁶ Se(t,p)	M	⁷⁸ Se(n,n' γ)		
G	⁷⁷ Se(n, γ) E=thermal	N	⁷⁸ Se(p,p'),(pol p,p')		

E(level) [†]	J ^π [‡]	T _{1/2}	XREF	Comments
0.0 [#]	0 ⁺	stable	ABCDEFGH LMNOPQRS	
613.727 [#] 3	2 ⁺	9.79 ps 21	ABCDEFGH J LMNOPQR	$\mu=+0.77$ 5 (1998Sp03) Q=-0.20 7 (2003Ha15) B(E2) $\uparrow=0.332$ 7 $\langle r^2 \rangle^{1/2}=4.1407$ fm 18 (2004An14 evaluation). B(E2) \uparrow : weighted average of 0.325 45 (2003Ha15), 0.392 66 (deduced from T _{1/2} =8.3 ps 14 (1987Sc07),RDM measurement in ($\alpha,2n\gamma$)), 0.327 7 (1977Le11), 0.385 35 (1962St02), 0.35 3 (1962Gal13), 0.36 7 (1960Le07) and 0.36 5 (1956Te26). All values, except for 1987Sc07, are from cross sections and yields in Coulomb excitation. Other: 0.335 9 (2001Ra27 evaluation). T _{1/2} : from B(E2)=0.332 7. Other: 9.69 ps 26 (2001Ra27 evaluation). J ^π : from L(t,p)=2. Also, L=2 and vector analyzing power in (p,p'). μ : from transient-field technique in Coul. ex. (1998Sp03), sign from 1969He11. Other: +0.78 22 (1969He11,IMPAC technique). See also 1989Ra17 evaluation and 2005St24 compilation. Q: from Coulomb excitation (2003Ha15). Others:-0.26 9 (1977Le11), -0.30 11 (1976VoZY). See also 1989Ra17 evaluation and 2005St24 compilation.
1308.644 [@] 5	2 ⁺	4.2 ps 3	A CD FG KLMNOPQR	$\mu=0.66$ 22 (1998Sp03) Q=+0.17 9 (2003Ha15) μ : from transient-field technique in Coul. ex. (1998Sp03). See also 2005St24 compilation. Q: from Coulomb excitation (2003Ha15). T _{1/2} : from B(E2) in Coulomb excitation. Other: 3.8 ps 10 from recoil-distance method in ($\alpha,2n\gamma$) (1987Sc07). Weighted

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued)

⁷⁸Se Levels (continued)

E(level) [†]	J ^{π‡}	T _{1/2}	XREF	Comments
				average of the two values is also 4.2 ps 3.
1498.599 9	0 ⁺	45 ps 8	A C FG LMNO qR	J ^π : from L(t,p)=2. Also, L=2 and vector analyzing power in (p,p') and J=2 from circular polarization in (n,γ). XREF: L(1510)q(1510). T _{1/2} : from B(E2)(↑) in Coul. ex. J ^π : 0 from γγ(θ) in (n,γ); L(d,p)=1.
1502.825# 13	4 ⁺	1.04 ps 5	A D G MNOPqR	μ=1.6 5 (1998Sp03) Q=-0.68 15 (2003Ha15) XREF: q(1510). μ: from transient-field technique in Coul. ex. (1998Sp03). See also 2005St24 compilation. J ^π : γ(θ) and linear polarization in (α,2nγ). T _{1/2} : weighted average of 1.05 ps 5 from B(E2) in Coul. ex. and 0.9 ps 2 from DSA in (α,2nγ) (1987Sc07).
1758.689 17	0 ⁺		A C G MNO Q	J ^π : J=0 from γγ(θ) in (n,γ); γ's to 2 ⁺ .
1853.927@ 12	3 ⁺	1.2 ps 4	A D G LMNO	XREF: L(1880). J ^π : γ(θ) and polarization measurements in (α,2nγ). T _{1/2} : DSA in (α,2nγ) (1987Sc07).
1995.897 8	2 ⁺	4.6 ps +32-14	A C FGH MNO QR	XREF: Q(2030). J ^π : L(t,p)=2; L(p,p')=2; J=2 from circular polarization in (n,γ). T _{1/2} : from B(E2)(↑) in Coulomb excitation.
2190.65@ 18	4 ⁺	0.7 ps 3	D MN Q	XREF: Q(2220). J ^π : γ(θ) and polarization measurements in (α,2nγ). T _{1/2} : DSA in (α,2nγ) (1987Sc07). J ^π : γ to 2 ⁺ suggests 0 ⁺ to 4 ⁺ .
2267.07 12			G	J ^π : γ to 0 ⁺ .
2299.8 5	1,2(+)		M	J ^π : M1+E2 γ to 2 ⁺ ; J=2 from γγ(θ) in (n,γ).
2327.329 19	2 ⁺	0.28 ps +13-8	A C G MNO	T _{1/2} : DSA in (n,n'γ). J ^π : log ft=5.91 from 1 ⁺ ; J=0 from γγ(θ) in (n,γ).
2335.24 5	0 ⁺		A C G M	J ^π : L(t,p)=0. But L(d,p)=1 for E=2360. It is possible that the (t,p) and (d,p) reactions correspond to the 2335 level.
2361.85 14	(0 ⁺)		FG L	
2507.32& 5	3 ⁻	6.2 ps 14	A DEFG MNOP R	B(E3)↑=0.027 3 (2002Ki06,1974Ba80) B(E3)↑: from Coul. ex. J ^π : L(p,p') and vector analyzing power in (p,p'). T _{1/2} : recoil-distance method in (α,2nγ) (1987Sc07). J ^π : L(t,p)=2. T _{1/2} : DSA in (n,n'γ). J ^π : γ to 4 ⁺ suggests 2 ⁺ to 6 ⁺ .
2536.94 4	2 ⁺	0.055 ps 7	A C FG MNO	J ^π : γ(θ) and polarization in (α,2nγ). T _{1/2} : DSA in (α,2nγ).
2546.3 3			G	
2546.51# 15	6 ⁺	0.49 ps 14	D M	J ^π : γ(θ) and polarization in (α,2nγ). T _{1/2} : DSA in (α,2nγ).
2560?	(1 ⁻ ,2 ⁻ ,3 ⁻)		L	E(level): no uncertainty available. May correspond to adjacent level. J ^π : L(d,p)=(2). J ^π : γ to 4 ⁺ .
2629.6 5			D	
2647.472 13	(1,2) ⁺		A C G MNO	J ^π : log ft=6.24 from 1 ⁺ ; γ's to 2 ⁺ and 3 ⁺ .
2682.110 16	4 ⁺		A FG MNO	J ^π : L(t,p)=4, L(p,p')=4. J ^π inconsistent with possible primary transition in (n,γ) and log f ^l t from 2 ⁻ small, but decay mode of 2682 level is consistent in (n,γ), β ⁻ , and (p,p'γ); so only one level appears to be involved.

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued)

⁷⁸Se Levels (continued)

E(level) [†]	J ^{π‡}	T _{1/2}	XREF			Comments
2719.3 5					M	
2735.0 [@] 6	(5 ⁺)	0.62 ps 21	D		M	J ^π : γ(θ) and band assignment in (α,2nγ). T _{1/2} : DSA in (α,2nγ) (1987Sc07).
2742.52 ^{&} 14	4 ⁻	0.42 ns 14	D		N	J ^π : γ(θ) and polarization in (α,2nγ) (1987Sc07). E2 γ from 6 ⁻ and E1 γ to 4 ⁺ . T _{1/2} : recoil-distance method in (α,2nγ) (1987Sc07).
2753.03 18	0 ⁺			F		J ^π : L(t,p)=0.
2754.46 17	2 ⁺			G	M O	J ^π : γ(circ pol) in (n,γ); γ to 0 ⁺ . E(level): from primary transition in (n,γ). The 757 and 2140 γ's are not seen in (n,γ), and the 2156γ is not seen in (n,n'γ) or (p,p'γ). It is possible that the γ transitions define more than one level, in particular, the 2753 10+ level reported in (t,p) is perhaps being excited. Transitions from the 2754.46 level are both included in the least-squares fit for determining the energies of other levels.
2838.49 7	(2 ⁺)		A	G	MN	J ^π : γ's to 0 ⁺ and 4 ⁺ .
2864.12 7				G	N	J ^π : γ to 3 ⁺ suggests 1 ⁺ :5 ⁺ .
2889.90 ^{&} 11	5 ⁻	18 ps 5	D F		M O	XREF: F(2893). T _{1/2} : recoil-distance method in (α,2nγ) (1987Sc07). J ^π : L(t,p)=5; γ(θ) and polarization measurements in (α,2nγ).
2898.13 6	2		C	G	MN	J ^π : γγ(θ) in (n,γ).
2914.7 5	4 ⁺	0.24 ns +15-8		F	MNO	T _{1/2} : DSA in (n,n'γ) (1989Do14). J ^π : L(t,p)=4.
2949.19 16	4 ⁻	>1.4 ps	D		LMNO	J ^π : γ(θ) in (α,2nγ); L(d,p)=4. T _{1/2} : DSA in (α,2nγ).
3003 9	3 ⁻			F		J ^π : L(t,p)=3.
3005.70 17	1,2 ⁺		C	G	J MNO	J ^π : log ft=6.28 from 1 ⁺ ; γ to 0 ⁺ .
3013.96 ^a 13	6 ⁻	3.0 ns 5	D F			J ^π : γ(θ) and polarization data in (α,2nγ). T _{1/2} : γγ(t) in (α,2nγ) (1987Sc07).
3039.81 6	(1 ⁺ to 4 ⁺)			G		J ^π : γ's to 2 ⁺ and 3 ⁺ .
3048.6 10	(3 ⁻)				NO	J ^π : L(p,p')=(3); γ to 4 ⁺ .
3061 12	0 ⁺ &5 ⁻			F		J ^π : L(t,p)=0+5.
3088.7 21	(5 ⁻)			f	N	J ^π : L(p,p')=5. L(t,p)=0+4 for a doublet.
3089.73 15	(0 ⁺)		C	fG	M	J ^π : L(t,p)=0+4 for a doublet; γ to 2 ⁺ .
3130?	0 ⁺ ,1 ⁺ ,2 ⁺				L	E(level): may Be same as 3090 level. J ^π : L(d,p)=1.
3133.3 5	3 ⁻			F	M	J ^π : L(t,p)=3.
3139.7 15	4 ⁺				NO	J ^π : L(p,p')=4.
3140.2 [@] 4	(6 ⁺)	0.28 ps +14-7	D			J ^π : γ(θ) and band assignment in (α,2nγ). T _{1/2} : DSA in (α,2nγ) (1987Sc07).
3144.46 11	3 ⁻		A	FG	M	J ^π : L(t,p)=3; γ's to 2 ⁺ and 4 ⁺ .
3181.9 5	(2 ⁺)			f	MN	J ^π : L(d,p)=1; γ to 0 ⁺ ; L(t,p)=2.
3186.37 14	2 ⁺			fG		J ^π : L(t,p)=2; γ to 2 ⁺ .
3229.71 13	(1 ⁻ ,2,3)		A		M	J ^π : γ's to 3 ⁻ and 2 ⁺ ; log ft=6.5 from 2 ⁻ .
3242.68 7	2 ⁺			G	MN	J ^π : L(p,p')=2. E(level): from primary transition in (n,γ). Deexciting transitions 3241.8 and 2627.87 (doubly placed) are placed by 1979BrZE, with additional transitions reported and placed by 1987Su05 (all from (n,γ)), and give excitation energies of 3242.8 3, 3242.8 2, 3241.5 2, 3243.3 3 and 3243.4 1. The spread in

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued)

⁷⁸Se Levels (continued)

E(level) [†]	J ^π [‡]	T _{1/2}	XREF	Comments
3254.83 20	(0,1,2) ⁺		C G M	excitation energies suggests that either one or more transitions are misplaced, or that there is more than one level at this energy. Transitions of energy 2629 and 3242 are reported also in (n,n'γ) and placed from a 3242 level. The 1484γ is not reported in (n,n'γ). Transitions from this level are not used in the least-squares fit for determining the energies of the other levels.
3288.27 6	1 ⁻		FG M	J ^π : γ to 2 ⁺ ; log ft=5.93 from 1 ⁺ .
3294.35 23	4 ⁺		A F N	J ^π : L(t,p)=1. XREF: N(3288).
3306.79& 16	6 ⁻	11 ps 4	D	J ^π : L(t,p)=4; L(p,p')=4. J ^π : γ(θ) and polarization data in (α,2nγ). T _{1/2} : recoil-distance method in (α,2nγ) (1987Sc07).
3309.9 20				E(level): multiplet.
3329 10			F L N	J ^π : L(d,p)=1+4 suggests a doublet, with opposite parities.
3372.6 3	3 ⁻		A F N	J ^π : L(p,p')=3.
3383.69 13	0 ⁺ to 4 ⁺		C G	J ^π : γ to 2 ⁺ .
3386.0 5	(2 ⁺)		f M	J ^π : γ's to 2 ⁺ and 0 ⁺ ; L(t,p)=2+5 for doublet.
3391? 8	(5 ⁻)		f	J ^π : L(t,p)=2+5 for a doublet.
3411.29 18	3 ⁻		A F N	J ^π : L(p,p')=3. L(t,p)=(4) is inconsistent.
3439.6 4	(1)		F I M	J ^π : γ to 0 ⁺ ; γ from 0 ⁻ resonance.
3450.94 14	0 ⁺		FG	J ^π : L(t,p)=0; γ to 2 ⁺ .
3453 4	3 ⁻		L N	J ^π : L(p,p')=3.
3488.2? 6		0.12 ps 4	D	J ^π : γ to 6 ⁺ and population in (α,2nγ) suggests 6,7,8 ⁺ . T _{1/2} : DSA in (α,2nγ) (1987Sc07).
3494.40 8	1,2 ⁽⁺⁾		G	J ^π : γ to 0 ⁺ .
3496.26 11			A	J ^π : γ's to 2 ⁺ and 3 ⁻ .
3522.91& 22	7 ⁻	1.4 ps +7 -4	D	J ^π : γ(θ) in (α,2nγ); M1 γ to 6 ⁻ . T _{1/2} : from DSA in (α,2nγ).
3523.5 5	1,2 ⁽⁺⁾		G	J ^π : γ to 0 ⁺ .
3527 14	1 ⁻		F	J ^π : L(t,p)=1.
3546 4	(2 ⁻ ,3 ⁻ ,4 ⁻)		F L N	J ^π : L(d,p)=(3).
3550.15 ^a 24	(7 ⁻)	3.5 ps 21	D	J ^π : band assignment in (α,2nγ). T _{1/2} : DSA and recoil-distance methods in (α,2nγ).
3585.0 [#] 3	8 ⁺	0.42 ps 14	D	J ^π : γ(θ) and polarization data in (α,2nγ). T _{1/2} : DSA in (α,2nγ).
3591.64 15	(1 ⁻)		FG	J ^π : L(t,p)=1, assuming 3598 9 corresponds to 3591.6 level and not 3603.8; γ to 2 ⁺ .
3603.8 10	2 ⁺			J ^π : L(p,p')=2; γ to 2 ⁺ .
3624.2 4	1,2 ⁽⁺⁾		fG	J ^π : L(t,p)=2 for a possible doublet; γ to 0 ⁺ .
3628.1 5			fG	J ^π : γ to 2 ⁺ .
3632.2 4	(1 ⁺ ,2 ⁺)			J ^π : γ's to 0 ⁺ and 3 ⁺ .
3686.50 16	3 ⁻		FG LMN	J ^π : L(t,p)=3; L(d,p)=2.
3704.0 [@] 8	(7 ⁺)	0.83 ps 21	D	J ^π : γ(θ) and band assignment in (α,2nγ). T _{1/2} : DSA in (α,2nγ).
3711.3 5	(1,2,3)		A	J ^π : log ft=7.0 from 2 ⁻ ; γ to 2 ⁺ .
3735.03 17	0 ⁺ to 4 ⁺		G	J ^π : γ to 2 ⁺ .
3754 15			F	
3774 4	3 ⁻		F N	E(level): from (p,p'). J ^π : L(t,p)=3; L(p,p')=3.
3830	1 ⁻ ,2 ⁻ ,3 ⁻		L	J ^π : L(d,p)=2.
3830.7 [@] 3	8 ⁺	0.55 ps 14	D	J ^π : γ(θ) and polarization measurements in (α,2nγ). E(level): the 8 ⁺ member of β band is either 3831 or 4121 level. T _{1/2} : DSA in (α,2nγ).

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued)

⁷⁸Se Levels (continued)

E(level) [†]	J ^π [‡]	T _{1/2}	XREF		Comments
3881 4	3 ⁻			N	J ^π : L(p,p')=3.
3894.55 15	2 ⁺		FG		J ^π : L(t,p)=2.
3933 9	2 ⁺		F		J ^π : L(t,p)=2.
3959.93 24	1,2 ⁽⁺⁾		G		J ^π : γ to 0 ⁺ .
3995 4	5 ⁻			N	J ^π : L(p,p')=5.
3999.33 15	1 ⁻		FG		J ^π : L(t,p)=1.
4037.01 21	(1 ⁻ ,3 ⁻)		fG		J ^π : L(t,p)=1+3 for a doublet; γ to 2 ⁺ .
4038 10	(1 ⁻ ,3 ⁻)		f		J ^π : L(t,p)=1+3 for a doublet.
4048.0& 6	8 ⁻	0.9 ps 3	D		J ^π : γ(θ) and polarization data in (α,2nγ). T _{1/2} : DSA in (α,2nγ).
4050 4	(5 ⁻)			N	J ^π : L(p,p')=(5).
4079.7 3	1,2 ⁽⁺⁾		G		J ^π : γ to 0 ⁺ .
4106 12	1 ⁻		F		J ^π : L(t,p)=1.
4120?	0 ⁻ ,1 ⁻			L	J ^π : L(d,p)=0.
4121.2 3	8 ⁺	>0.7 ps	D		J ^π : γ(θ) and polarization data in (α,2nγ). E(level): this level may be the 8 ⁺ member of β band, although, 3831 level is presently assigned as the 8 ⁺ member. T _{1/2} : DSA in (α,2nγ). Upper limit is <0.35 ns from pulsed-beam γ-timing in (α,2nγ).
4122 4	4 ⁺		F	N	E(level): weighted average from (p,p') and (t,p). J ^π : L(t,p)=4; L(p,p')=4.
4153.10 16	(1)		G I		J ^π : γ from 0 ⁻ resonance.
4155 4	3 ⁻		F	N	J ^π : L(p,p')=3. E(level): weighted average from (p,p') and (t,p).
4181.85 14	0 ⁺		FG		J ^π : L(t,p)=0.
4190?	0 ⁻ ,1 ⁻			L	J ^π : L(d,p)=0.
4214.1 ^a 4	(8 ⁻)	>1.4 ps	D		J ^π : γ(θ) and band assignment in (α,2nγ). T _{1/2} : DSA in (α,2nγ).
4224 10	3 ⁻		F		E(level): an unplaced 6274.40 16 transition in (n,γ), if a primary, would define a level at 4222.75 17, but the transition would be 1 ⁻ to 3 ⁻ . J ^π : L(t,p)=3.
4245.4 5	(1)			I	J ^π : γ from 0 ⁻ resonance.
4253.11 12	(2 ⁺)		fG		J ^π : L(t,p)=5+2 for a doublet; γ's to 2 ⁺ .
4253.64 17	(5 ⁻)		f	N	E(level): from (p,p'). J ^π : L(t,p)=5+2 for a doublet; L(p,p')=(4) seems inconsistent unless S=1 is involved.
4265 10	0 ⁺		F		J ^π : L(t,p)=0.
4297.38 15	2 ⁺		FG		J ^π : L(t,p)=2.
4341.61 13	1,2 ⁽⁺⁾		G		J ^π : γ to 0 ⁺ .
4345 11	3 ⁻		F		J ^π : L(t,p)=3.
4366.61 15	(1 ⁻)		fG I	L	J ^π : L(t,p)=3+1 for a doublet; L(d,p)=2; γ's to 0 ⁺ and 2 ⁺ ; γ from 0 ⁻ resonance.
4369 11	(3 ⁻)		f		J ^π : L(t,p)=3+1 for a doublet.
4386.68 13	(1,2 ⁺)		G		J ^π : γ to 0 ⁺ . Doubly-placed γ to 0 ⁺ .
4409 11	2 ⁺		F		E(level): an unplaced 6091.81 18 transition in (n,γ), if a primary, would define a level at 4405.65 19. J ^π : L(t,p)=2.
4412.02& 24	(9 ⁻)		D		J ^π : band assignment in (α,2nγ).
4424 4	(2 ⁺)			N	E(level): an unplaced 6077.24 18 transition in (n,γ), if a primary, would define a level at 4420.22 19. J ^π : L(p,p')=(2).
4448.24 15	1,2 ⁽⁺⁾		G		J ^π : γ's to 0 ⁺ and 2 ⁺ .
4451 11	(0 ⁺ &3 ⁻)		F		J ^π : L(t,p)=0+3.
4468.6 4	1,2 ⁽⁺⁾		G		J ^π : γ to 0 ⁺ .

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued)

⁷⁸Se Levels (continued)

E(level) [†]	J ^π [‡]	T _{1/2}	XREF		Comments
4483 11	4 ⁺		F		J ^π : L(t,p)=4.
4493 4	(3) ⁻			L N	J ^π : L(d,p)=2. L(p,p')=(3).
4509 11	2 ⁺		F		J ^π : L(t,p)=2.
4528.8 4			G		J ^π : 0 ⁺ to 4 ⁺ from possible γ to 2 ⁺ .
4557 4				N	
4569 11	(0 ⁺ &4 ⁺)		F		E(level): an unplaced 5932.03 21 transition in (n,γ), if a primary, would define a level at 4565.45 22.
					J ^π : L(t,p)=0+4.
4591 11	(3) ⁻		F	L	E(level): from (t,p).
					J ^π : L(t,p)=(3); L(d,p)=2.
4616 11	4 ⁺		F		J ^π : L(t,p)=4.
4622 4	5 ⁻			N	J ^π : L(p,p')=5.
4625.1 [#] 5	(10 ⁺)		D		J ^π : band assignment in (α,2nγ).
4639 11	3 ⁻		F		J ^π : L(t,p)=3.
4672.8 3			G		
4684.30 17			G		
4689.8 3	(2 ⁺)		fG		J ^π : γ to 0 ⁺ ; L(t,p)=2.
4697.07 13	(2 ⁺)		fG		J ^π : γ to 0 ⁺ ; L(t,p)=2.
4723.21 18	2 ⁺		FG		J ^π : L(t,p)=2.
4758 11	4 ⁺ &1 ⁻		F	N	XREF: N(4741).
					E(level): doublet from mixed L-transfer.
					J ^π : L(p,p')=4; L(t,p)=4+1.
4786.9 [@] 5	(10 ⁺)	>1.4 ps	D		J ^π : γ(θ), pol in (α,2nγ).
					T _{1/2} : DSA in (α,2nγ).
4787.93 21	(1) ⁻		G	L	J ^π : L(d,p)=0; γ to 2 ⁺ .
4791.5 5	0 ⁺		FG		J ^π : L(t,p)=0.
4811.5 3	2 ⁺		FG		J ^π : L(t,p)=2.
4819.2 ^a 6	(9 ⁻)	0.9 ps 3	D		J ^π : band assignment in (α,2nγ).
					T _{1/2} : DSA in (α,2nγ).
4857.0 [@] 9	(9 ⁺)	1.1 ps 4	D		J ^π : γ(θ) and band assignment in (α,2nγ).
					T _{1/2} : DSA in (α,2nγ).
4857 11	1 ⁻		F		J ^π : L(t,p)=1.
4879 11	3 ⁻		F		J ^π : L(t,p)=3.
4902 4	3 ⁻			L N	J ^π : L(p,p')=3; L(d,p)=2.
4904 10	2 ⁺		F		J ^π : L(t,p)=2.
4944 11	2 ⁺		F		J ^π : L(t,p)=2.
4957.3 3	1,2 ⁽⁺⁾		G		J ^π : γ to 0 ⁺ .
4972.3 3	1 ⁻		FG	L	XREF: F(4980)L(4970).
					J ^π : L(t,p)=1; L(d,p)=2.
4998.3 5			G		
5004.65 23	1,2 ⁽⁺⁾		G		J ^π : γ's to 0 ⁺ and 2 ⁺ .
5022.14 17			G		
5029.63 24	2 ⁺		FG		J ^π : L(t,p)=2.
5055 12			F		
5090.8 3			FG		XREF: F(5081).
5094.8 8			D		
5101.9 5			FG		
5120?	0 ⁻ ,1 ⁻			L	J ^π : L(d,p)=0.
5126.52 16	(2,3,4)		FG		J ^π : γ's to 2 ⁺ and 3 ⁺ ; multiply-placed γ to 4 ⁺ .
5136? 15			F		E(level): may Be same as 5126 level.
5164.05 16			FG		XREF: F(5169).
					J ^π : doubly-placed γ's to 2 ⁺ .
5180.75 22	1 ⁽⁺⁾ ,2 ⁽⁺⁾		FG		J ^π : γ's to 0 ⁺ and 3 ⁺ .
5205 15	1 ⁻ ,2 ⁻ ,3 ⁻		F	L	XREF: L(5210).
					J ^π : L(d,p)=2.

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{78}Se Levels (continued)

E(level) [†]	J^π [‡]	$T_{1/2}$	XREF		Comments
5235 15				F	
5247 15				F	
5290.22 18	1,2(+)			G	J^π : γ 's to 0 ⁺ and 2 ⁺ .
5295.2 3	3 ⁻		FG	N	J^π : L(p,p')=3.
5339.7 3	1,2(+)			G	J^π : γ 's to 0 ⁺ and 2 ⁺ .
5356.51 17	(2 ⁺)		G	L	J^π : L(d,p)=(2); γ to 2 ⁺ .
5391.0 3				FG	
5422 15				F	
5440.3 3				G	
5451.2 4	1,2(+)			G	J^π : γ to 0 ⁺ .
5480?	(1 ⁺ ,2 ⁺ ,3 ⁺)			L	J^π : L(d,p)=(2).
5513.26 19	1,2(+)			G	J^π : γ to 0 ⁺ ; multiply-placed γ to (4 ⁺).
5580 15				F	
5610?	2 ⁺			L	J^π : L(d,p)=2.
5689.1 8				D	
5709 15				F	
5783.8 [#] 7	(12 ⁺)	>0.6 ps		D	J^π : band assignment. $T_{1/2}$: DSA in (α ,2n γ).
5837 15				F	
6161 15				F	

[†] From (n, γ), (α ,2n γ) or other γ -ray studies if populated in these sets. In addition to the states shown, broad peaks are reported at 1450, 1790, and 3560 in (^{16}O , ^{14}C), and at 2360, 2550, 2730, 2830, 2990, 3170, 3270, 3370, 3500, and 3560 in (d,d').

[‡] Target $J^\pi=1/2^-$ for L(d,p) and 0⁺ for L(t,p).

[#] Band(A): g.s. band.

@ Band(B): Probable β band.

& Band(C): Probable octupole band.

^a Band(D): $\Delta J=1$ band based on 6⁻.

Adopted Levels, Gammas (continued)

$E_i(\text{level})$	J_i^π	$\gamma(^{78}\text{Se})$		E_f	J_f^π	Mult. [‡]	δ^{\ddagger}	Comments
		E_γ [†]	I_γ [†]					
613.727	2 ⁺	613.725 3	100	0.0	0 ⁺	E2		B(E2)(W.u.)=33.5 8 Mult.: from $\gamma(\theta)$ and $\gamma(\text{pol})$ in ($\alpha, 2n\gamma$).
1308.644	2 ⁺	694.916 4	100.0 20	613.727	2 ⁺	E0+M1+E2	+3.5 5	B(M1)(W.u.)=0.00067 19; B(E2)(W.u.)=22.2 18 Mult., δ : mult from $\gamma(\theta)$ in Coul. ex., δ from (n, γ). Others: +4.0 7 in ($\alpha, 2n\gamma$), +2.7 +9-6 in Coulomb excitation. X(E0/E2)=0.10 1 in (n, γ). B(E2)(W.u.)=0.76 6 B(E2)(W.u.)=1.17 21
1498.599	0 ⁺	1308.59 4 884.861 15	75.0 7 100	0.0 613.727	0 ⁺ 2 ⁺	E2 E2		B(E2)(W.u.)=0.76 6 B(E2)(W.u.)=1.17 21
1502.825	4 ⁺	1498 ^b 889.099 12	100	0.0 613.727	0 ⁺ 2 ⁺	[E0] E2		X(E0/E2) \leq 0.07 in (n, γ). B(E2)(W.u.)=49.5 24 Mult.: from $\gamma(\theta)$ in ($\alpha, 2n\gamma$) and Coul. ex.
1758.689	0 ⁺	260.1 ^b 449.94 6 1144.959 17	3.7 4 100 4	1498.599 1308.644 613.727	0 ⁺ 2 ⁺ 2 ⁺	[E0] (E2) [E0]		X(E0/E2) \leq 1.36 in (n, γ). Mult.: Q from $\gamma\gamma(\theta)$. $\Delta\pi$ =no from level scheme. X(E0/E2) \leq 0.27 in (n, γ).
1853.927	3 ⁺	1758 ^b 351.49 17 545.300 13	2.7 4 51 7	1502.825 1308.644	4 ⁺ 2 ⁺	M1+E2	+0.42 4	B(M1)(W.u.)=0.032 12; B(E2)(W.u.)=25 10 δ : from $\gamma(\theta)$ in (n,n' γ). Others: +0.45 10 in ($\alpha, 2n\gamma$). Mult.: from angular distribution and polarization measurements in 1987Sc07 and 1982Ma45.
		1240.13 3	100 10	613.727	2 ⁺	M1+E2	-0.41 +13-31	B(M1)(W.u.)=(0.0054 20); B(E2)(W.u.)=(0.8 5) Mult., δ : M1+E2 from $\gamma(\theta, \text{pol})$ in ($\alpha, 2n\gamma$); δ from $\gamma\gamma(\theta)$ in (n, γ).
1995.897	2 ⁺	497.294 7 687.254 7	11 2 57 5	1498.599 1308.644	0 ⁺ 2 ⁺	[E2] M1+E2(+E0)	-0.30 19	B(E2)(W.u.)=10 +4-8 B(M1)(W.u.)=0.0034 +12-25; B(E2)(W.u.)=0.8 +10-8 Mult., δ : from $\alpha(\text{K})\text{exp}$ and $\gamma\gamma(\theta)$ (1987Su05) in (n, γ); δ =0.12 to 0.49; sign is negative. X(E0/E2)=0.26 to 9.5 in (n, γ). B(M1)(W.u.)=0.00039 +13-28; B(E2)(W.u.)=0.05 +3-4 X(E0/E2)=11 4 in (n, γ). Mult., δ : from $\alpha(\text{K})\text{exp}$ and $\gamma\gamma(\theta)$ (1987Su05) in (n, γ). B(E2)(W.u.)=0.09 +3-6 B(M1)(W.u.)=0.04 3 B(E2)(W.u.)=40 +50-40 E γ : from (n,n' γ).
		1382.16 3	58 5	613.727	2 ⁺	E0+M1+E2	+0.44 10	
2190.65	4 ⁺	1995.87 8 688.0 3 881.7	100 4 100 7 <276	0.0 1502.825 1308.644	0 ⁺ 4 ⁺ 2 ⁺	[E2] (M1) [E2]		
2267.07		1576 1 271.1 8 958.37 19 1653.28 15	24 7 24 8 40 6 100 9	613.727 1995.897 1308.644 613.727	2 ⁺ 2 ⁺ 2 ⁺ 2 ⁺			
2299.8	1,2 ⁽⁺⁾	2299.8 5	100	0.0	0 ⁺			
2327.329	2 ⁺	331.2 3	1.6 3	1995.897	2 ⁺			

Adopted Levels, Gammas (continued)

$\gamma(^{78}\text{Se})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	δ^\ddagger	Comments
2327.329	2 ⁺	568.7 4	2.2 3	1758.689	0 ⁺	[E2]		B(E2)(W.u.)=32 +11-16
		824.8# 4	2.0 5	1502.825	4 ⁺			
		1018.65 5	6.1 3	1308.644	2 ⁺			
		1713.55 3	100 6	613.727	2 ⁺	E0+M1+E2	-1.8 5	B(M1)(W.u.)=0.0031 +17-20; B(E2)(W.u.)=4.5 +15-22 Mult.: from $\alpha(\text{K})\text{exp}$ in (n, γ) (1987Su05). X(E0/E2)=1.21 23 in (n, γ). B(E2)(W.u.)=0.10 +6-7
		2327.26 6	8 4	0.0	0 ⁺	[E2]		
2335.24	0 ⁺	575.0#b 10	<41	1758.689	0 ⁺			
		1026.59 20	10.8 8	1308.644	2 ⁺			
		1721.50 5	100 6	613.727	2 ⁺	E2		Mult.: from $\alpha(\text{K})\text{exp}$ =0.00015 5 in (n, γ) (1987Su05).
2361.85	(0 ⁺)	1748.21 15	100	613.727	2 ⁺			
2507.32	3 ⁻	1004.73 20	20 4	1502.825	4 ⁺	[E1]		B(E1)(W.u.)=9.E-6 3
		1198.6 3	100 4	1308.644	2 ⁺	(E1(+M2))	+0.09 5	B(E1)(W.u.)=2.5 \times 10 ⁻⁵ 6; B(M2)(W.u.)=(0.6 + δ -6) Mult.: from $\gamma(\theta)$ in (α ,2n γ) (1987Sc07) and γ from 3 ⁻ to 2 ⁺ . δ : from $\gamma(\theta)$ in (n,n' γ). B(E1)(W.u.)=1.1 \times 10 ⁻⁶ 5 Mult.: D+Q, -0.05< δ <-3.0 from $\gamma\gamma(\theta)$ in (n, γ). $\Delta\pi$ =yes from level scheme.
		1893.46 6	18 6	613.727	2 ⁺	(E1)		
2536.94	2 ⁺	203.3# 5	4.1 10	2335.24	0 ⁺			
		1039.3 3	3 1	1498.599	0 ⁺	[E2]		B(E2)(W.u.)=10 4
		1228.25 17	28 2	1308.644	2 ⁺			
		1923.15 4	100 6	613.727	2 ⁺	(M1+E2)	-1.1 11	Mult.: D+Q, δ <2.2, sign=- from $\gamma\gamma(\theta)$ in (n, γ). $\Delta\pi$ =no from level scheme.
2546.3		279.0 8	100 17	2267.07				
		1043.6& 4	10& 4	1502.825	4 ⁺			
2546.51	6 ⁺	1043.9 3	100	1502.825	4 ⁺	E2		B(E2)(W.u.)=47 14 Mult.: from ce measurements in (α ,2n γ).
2629.6		1126.8 5	100	1502.825	4 ⁺			
2647.472	(1,2) ⁺	286.4 4	15 5	2361.85	(0 ⁺)			
		320.3 3	11 4	2327.329	2 ⁺			
		651.573 11	43 3	1995.897	2 ⁺			
		793.5 3	14.2 20	1853.927	3 ⁺			
		1338.78 5	100 7	1308.644	2 ⁺			
2682.110	4 ⁺	174.2 3	2.2 5	2507.32	3 ⁻			E_γ : from β^- decay.
		354.735 25	21 4	2327.329	2 ⁺			
		686.3 2	12 2	1995.897	2 ⁺			E_γ : from β^- decay.
		828.189 13	100 8	1853.927	3 ⁺	(M1+E2)	+1.0 7	Mult.: D+Q, δ =+0.32 to +1.63 from $\gamma\gamma(\theta)$ in (n, γ). $\Delta\pi$ =no from level scheme.
		1373.48 6	54 4	1308.644	2 ⁺			
		2068.4 4	6.5 14	613.727	2 ⁺			
2719.3		1410.6 5	100	1308.644	2 ⁺			

Adopted Levels, Gammas (continued)

$\gamma(^{78}\text{Se})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	δ^\ddagger	$\alpha^@$	Comments
2735.0	(5 ⁺)	1232.2 6	100 14	1502.825	4 ⁺				
2742.52	4 ⁻	551.9 2	100 6	2190.65	4 ⁺	E1			B(E1)(W.u.)=3.1×10 ⁻⁶ 11
		889 ^b 1	10	1853.927	3 ⁺	[E1]			B(E1)(W.u.)=7.4×10 ⁻⁸ 25 I _γ : from coin. No uncertainty given.
		1239.4 3	59	1502.825	4 ⁺	[E1]			B(E1)(W.u.)=1.6×10 ⁻⁷ 6 I _γ : from coin. No uncertainty given.
2754.46	2 ⁺	757.2 5	35 8	1995.897	2 ⁺				E _γ : from (n,n'γ). Observed only in (n,n'γ) and (p,p'γ).
		1256.7 4	38 8	1498.599	0 ⁺				E _γ : reported only in (n,γ).
		1445.8 2	100 15	1308.644	2 ⁺				
		2140.8 9	35 11	613.727	2 ⁺				E _γ : from (n,n'γ). Observed only in (n,n'γ) and (p,p'γ).
2838.49	(2 ⁺)	156.6 3	3.7 9	2682.110	4 ⁺				E _γ : from ⁷⁸ As β ⁻ decay only.
		503.7 2	16.7 16	2335.24	0 ⁺				E _γ : from ⁷⁸ As β ⁻ decay only.
		842.36 19	32 4	1995.897	2 ⁺				
		1079.67 22	46 4	1758.689	0 ⁺				I _γ : I _γ (842γ):I _γ (1080γ):I _γ (1530γ) from (n,γ). Values from (n,n'γ) are 233 67:100 33:100 33 and from β ⁻ decay are 43 5:65 5: 100 7.
		1529.60 17	100 6	1308.644	2 ⁺				
		2224.7 3	37 5	613.727	2 ⁺				E _γ : from ⁷⁸ As β ⁻ decay only.
		2839.0 3	2.2 11	0.0	0 ⁺				E _γ : from ⁷⁸ As β ⁻ decay only.
2864.12		504.4 ^b 2	43 10	2361.85	(0 ⁺)				E _γ : very poor fit in level scheme. Level-energy difference=502.3. Placement is suspect.
		1010.19 6	100 10	1853.927	3 ⁺				
2889.90	5 ⁻	343.5 2	15.9 8	2546.51	6 ⁺	E1			B(E1)(W.u.)=5.4×10 ⁻⁵ 16
		382.42 17	33.3 15	2507.32	3 ⁻	E2		0.00650	Mult.: from γ(θ) and polarization data in (α,2nγ). B(E2)(W.u.)=43 13
		1387.4 2	100 5	1502.825	4 ⁺	E1			B(E1)(W.u.)=5.2×10 ⁻⁶ 15 Mult.: from γ(θ) and polarization data in (α,2nγ).
2898.13	2	391.3 [#] 5	5 2	2507.32	3 ⁻				
		902.3 [#] 3	11 3	1995.897	2 ⁺				
		2284.37 6	100 12	613.727	2 ⁺	D+Q	-0.9 8		Mult.: from γγ(θ) in (n,γ), δ=0.11 to 1.69; sign=negative.
2914.7	4 ⁺	1411.9 5	100	1502.825	4 ⁺				
2949.19	4 ⁻	441.7 2	100 11	2507.32	3 ⁻	M1+E2	-0.6 3		B(M1)(W.u.)<0.076; B(E2)(W.u.)<250
		1095.2 5	56	1853.927	3 ⁺	[E1]			Mult.,δ: from (α,2nγ).
		1446.7 5	67	1502.825	4 ⁺	[E1]			B(E1)(W.u.)<5.1×10 ⁻⁵ B(E1)(W.u.)<2.6×10 ⁻⁵
3005.70	1,2 ⁺	2391.93 ^{&} 17	100 ^{&} 11	613.727	2 ⁺				
		3005.9 10	13 2	0.0	0 ⁺				E _γ : observed only in ⁷⁸ Br ε decay.
3013.96	6 ⁻	124.1 1	32.3 16	2889.90	5 ⁻	M1		0.0566	B(M1)(W.u.)=0.00077 14
		271.4 1	100 3	2742.52	4 ⁻	(E2)		0.0211	Mult.: from (α,2nγ). B(E2)(W.u.)=4.0 7 Mult.: from (α,2nγ).

Adopted Levels, Gammas (continued) $\gamma(^{78}\text{Se})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	δ^\ddagger	Comments
3013.96	6 ⁻	467.4 2	24.2 16	2546.51	6 ⁺	E1		B(E1)(W.u.)= 1.8×10^{-7} 4 Mult.: from ($\alpha, 2n\gamma$).
3039.81	(1 ⁺ to 4 ⁺)	1043.6 & 4 1186.02 12 1731.11 7	14 & 5 52 7 100 7	1995.897 2 ⁺ 1853.927 3 ⁺ 1308.644 2 ⁺				
3048.6	(3 ⁻)	1545.8 10	100	1502.825 4 ⁺				
3089.73	(0 ⁺)	2475.96 15	100	613.727 2 ⁺				
3133.3	3 ⁻	2519.5 5	100	613.727 2 ⁺				
3139.7	4 ⁺	1831.0 15	100	1308.644 2 ⁺				
3140.2	(6 ⁺)	593.7 5 949.6 4	61 6 100 12	2546.51 6 ⁺ 2190.65 4 ⁺		M1(+E2) [E2]	-0.2 2	B(M1)(W.u.)=0.14 +4-8; B(E2)(W.u.)=(20 +40-20) B(E2)(W.u.)=82 +24-43
3144.46	3 ⁻	462.2 2 637.1 2 1290.6 6 1642.0 3 1835.8 2	41 4 14 2 7 2 11 3 100 7	2682.110 4 ⁺ 2507.32 3 ⁻ 1853.927 3 ⁺ 1502.825 4 ⁺ 1308.644 2 ⁺				E_γ : weighted average from β^- decay and ($n, n'\gamma$). E=1834.58 23 is reported in (n, γ) but is probably not the same transition.
3181.9	(2 ⁺)	3181.8 5	100 17	0.0 0 ⁺				
3186.37	2 ⁺	2572.60 14	100	613.727 2 ⁺				
3229.71	(1 ⁻ , 2, 3)	722.4 2 1732 ^b 1 1921.3 3 2615.8 2	11 1 100 24 52 8	2507.32 3 ⁻ 1498.599 0 ⁺ 1308.644 2 ⁺ 613.727 2 ⁺				E_γ : from ($n, n'\gamma$).
3242.68	2 ⁺	595.89 10 976.31 23 1387.56 20 1484.12 17 1744.24 23 2627.87 & 14 3241.8 4	28 3 15 3 36 4 94 6 28 4 82 & 10 100 14	2647.472 (1,2) ⁺ 2267.07 1853.927 3 ⁺ 1758.689 0 ⁺ 1498.599 0 ⁺ 613.727 2 ⁺ 0.0 0 ⁺				
3254.83	(0, 1, 2) ⁺	2641.05 20	100	613.727 2 ⁺				
3288.27	1 ⁻	1292.49 10 1979.57 8 2674.36 13	22 3 6.9 23 100 15	1995.897 2 ⁺ 1308.644 2 ⁺ 613.727 2 ⁺				
3294.35	4 ⁺	756.9 3 968.2 7 1440.9 7 1791.9 7 2681.3 7	5 1 9 3 19 6 56 6 100 6	2536.94 2 ⁺ 2327.329 2 ⁺ 1853.927 3 ⁺ 1502.825 4 ⁺ 613.727 2 ⁺				

Adopted Levels, Gammas (continued)

$\gamma(^{78}\text{Se})$ (continued)									
$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	δ^\ddagger	$\alpha^@$	Comments
3306.79	6 ⁻	357.3 3	21.4 18	2949.19	4 ⁻	E2		0.00816	B(E2)(W.u.)=50 19 Mult.: from ($\alpha, 2n\gamma$).
		416.9 2	100 6	2889.90	5 ⁻	M1+E2	-0.4 1		B(M1)(W.u.)=0.012 5; B(E2)(W.u.)=15 9
		564.4 4	27 4	2742.52	4 ⁻	E2			B(E2)(W.u.)=6 3 Mult.: Q from $\gamma(\theta)$ in (n, γ) and RUL.
		760.4 3	42.9 18	2546.51	6 ⁺	(E1)			B(E1)(W.u.)=1.7 $\times 10^{-5}$ 7 Mult.: from ($\alpha, 2n\gamma$).
3372.6	3 ⁻	2064.1 5	100 33	1308.644	2 ⁺				
		2758.8 3	100 19	613.727	2 ⁺				
3383.69	0 ⁺ to 4 ⁺	2769.91 13	100	613.727	2 ⁺				
3386.0	(2 ⁺)	2772.0 5	100 25	613.727	2 ⁺				
		3387 1	50 13	0.0	0 ⁺				
3411.29	3 ⁻	903.6 4	39 13	2507.32	3 ⁻				
		2797.6 2	100 13	613.727	2 ⁺				
3439.6	(1)	3439.5 4	100	0.0	0 ⁺				
3450.94	0 ⁺	2837.16 14	100	613.727	2 ⁺				
3488.2?		941.7 5	100	2546.51	6 ⁺				
3494.40	1,2 ⁽⁺⁾	655.90 7	100 8	2838.49	(2 ⁺)				
		1159.09 10	82 22	2335.24	0 ⁺				
		1499.1 3	65 16	1995.897	2 ⁺				
3496.26		657.9 2	58 6	2838.49	(2 ⁺)				
		959.0 2	100 10	2536.94	2 ⁺				
		988.2 4	20 5	2507.32	3 ⁻				
		1169.5 4	26 7	2327.329	2 ⁺				
		2187.8 2	78 8	1308.644	2 ⁺				
3522.91	7 ⁻	216.1 2	12.9 16	3306.79	6 ⁻	M1		0.01327	
		509 1	64	3013.96	6 ⁻				
		633.0 5	100	2889.90	5 ⁻	E2			
		976.7 4	53 5	2546.51	6 ⁺	(E1)			
3523.5	1,2 ⁽⁺⁾	3523.4 5	100	0.0	0 ⁺				
3550.15	(7 ⁻)	536.2 2	100	3013.96	6 ⁻				
3585.0	8 ⁺	1038.6 3	100	2546.51	6 ⁺	E2			B(E2)(W.u.)=56 19 Mult.: from ce data in ($\alpha, 2n\gamma$).
3591.64	(1 ⁻)	2977.85 15	100	613.727	2 ⁺				
3603.8	2 ⁺	2990 1	100	613.727	2 ⁺				
3624.2	1,2 ⁽⁺⁾	3624.1 & 4	100 &	0.0	0 ⁺				
3628.1		2319.4 5	100	1308.644	2 ⁺				
3632.2	(1 ⁺ , 2 ⁺)	1778.3 5		1853.927	3 ⁺				
		1873.5 5		1758.689	0 ⁺				
		3632 1		0.0	0 ⁺				
3686.50	3 ⁻	3072.71 16	100	613.727	2 ⁺				

Adopted Levels, Gammas (continued)

$\gamma(^{78}\text{Se})$ (continued)									
$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	δ^\ddagger	α^\oplus	Comments
3704.0	(7 ⁺)	969.0 5	100 8	2735.0	(5 ⁺)	E2			B(E2)(W.u.)=36 10 Mult.: from ($\alpha, 2n\gamma$).
3711.3	(1,2,3)	1158.7 ^{ab} 5	12 ^a	2546.51	6 ⁺				
3735.03	0 ⁺ to 4 ⁺	3097.5 5	100	613.727	2 ⁺				
3830.7	8 ⁺	3121.24 17	100	613.727	2 ⁺				
		245.6 2	33 2	3585.0	8 ⁺	M1		0.00960	B(M1)(W.u.)=0.67 18 Mult.: from ($\alpha, 2n\gamma$).
		1284.1 3	100 6	2546.51	6 ⁺	E2			B(E2)(W.u.)=11 3 Mult.: from ($\alpha, 2n\gamma$).
3894.55	2 ⁺	2391.93 ^{&} 17	100 ^{&} 17	1502.825	4 ⁺				
		3893.7 3	5.8 17	0.0	0 ⁺				
3959.93	1,2 ⁽⁺⁾	3345.8 4	86 15	613.727	2 ⁺				
		3960.0 3	100 15	0.0	0 ⁺				
3999.33	1 ⁻	1672.8 4	74 29	2327.329	2 ⁺				
		2003.1 6	74 29	1995.897	2 ⁺				
		2240.1 8	58 29	1758.689	0 ⁺				
		3385.88 21	100 6	613.727	2 ⁺				
		3998.2 3	19 3	0.0	0 ⁺				
4037.01	(1 ⁻ , 3 ⁻)	3423.20 21	100	613.727	2 ⁺				
4048.0	8 ⁻	741.2 5	100	3306.79	6 ⁻	E2			B(E2)(W.u.)=140 50 Mult.: from ($\alpha, 2n\gamma$).
4079.7	1,2 ⁽⁺⁾	4079.6 3	100	0.0	0 ⁺				
4121.2	8 ⁺	290.5 2	100 11	3830.7	8 ⁺	M1		0.00633	B(M1)(W.u.)<0.55 Mult.: from ($\alpha, 2n\gamma$).
		536.2 2	56	3585.0	8 ⁺	M1+E2	-0.4 3		B(M1)(W.u.)<0.051; B(E2)(W.u.)<70 Mult., δ : from ($\alpha, 2n\gamma$).
		1574 1	78 22	2546.51	6 ⁺	(E2)			B(E2)(W.u.)<1.4 Mult.: $\Delta J=2$, (Q) from ($\alpha, 2n\gamma$). RUL and $\Delta\pi$ =no from level scheme.
4181.85	0 ⁺	2186.0 10	59 24	1995.897	2 ⁺				
		2873.15 14	100 11	1308.644	2 ⁺				
4214.1	(8 ⁻)	664.0 3	80 10	3550.15	(7 ⁻)				
		1200 1	\approx 100	3013.96	6 ⁻	[E2]			B(E2)(W.u.)<5
4253.11	(2 ⁺)	2257.53 20	100 20	1995.897	2 ⁺				
		2944.20 14	54 6	1308.644	2 ⁺				
		3639.7 5	22 4	613.727	2 ⁺				
4297.38	2 ⁺	2988.67 15	100	1308.644	2 ⁺				
4341.61	1,2 ⁽⁺⁾	2843.02 ^{&} 14	114 ^{&} 15	1498.599	0 ⁺				
		4341.2 3	100 8	0.0	0 ⁺				
4366.61	(1 ⁻)	3057.90 16	100 17	1308.644	2 ⁺				
		4366.5 3	33 11	0.0	0 ⁺				

Adopted Levels, Gammas (continued)

γ(⁷⁸Se) (continued)

<u>E_i(level)</u>	<u>J_i^π</u>	<u>E_γ[†]</u>	<u>I_γ[†]</u>	<u>E_f</u>	<u>J_f^π</u>	<u>Mult.[‡]</u>	<u>Comments</u>
4386.68	(1,2 ⁺)	2627.87 & 14 3773.2 3	222 & 29 100 11	1758.689 613.727	0 ⁺ 2 ⁺		
4412.02	(9 ⁻)	363.1 4 862.0 5	26 3 77 9	4048.0 3550.15	8 ⁻ (7 ⁻)		
4448.24	1,2 ⁽⁺⁾	889.1 ^b 1 2452.27 16 4448.2 3	100 67 11 100 21	3522.91 1995.897 0.0	7 ⁻ 2 ⁺ 0 ⁺		
4468.6	1,2 ⁽⁺⁾	3855.0 & 4 4468.0 5	500 & 50 100 25	613.727 0.0	2 ⁺ 0 ⁺		
4528.8		3220.1 & 4	100 &	1308.644	2 ⁺		
4625.1	(10 ⁺)	794.6 ^b 4 1040.3 6	<21 100 24	3830.7 3585.0	8 ⁺ 8 ⁺		
4672.8		4059.0 3	100	613.727	2 ⁺		
4684.30		3375.73 20 4070.1 3	48 5 100 7	1308.644 613.727	2 ⁺ 2 ⁺		
4689.8	(2 ⁺)	4689.6 3	100	0.0	0 ⁺		
4697.07	(2 ⁺)	2843.02 & 14 4697.2 3	526 & 68 100 37	1853.927 0.0	3 ⁺ 0 ⁺		
4723.21	2 ⁺	3220.1 & 4 3224.4 5 3414.57 21	112 & 29 60 30 100 12	1502.825 1498.599 1308.644	4 ⁺ 0 ⁺ 2 ⁺		
4786.9	(10 ⁺)	161.9 2 955.9 5 1202.2 6	≈87 100 9 <13	4625.1 3830.7 3585.0	(10 ⁺) 8 ⁺ 8 ⁺	(E2) [E2]	B(E2)(W.u.)<13 B(E2)(W.u.)<0.3
4787.93	(1) ⁻	3479.36 22 4173.3 5	72 11 100 17	1308.644 613.727	2 ⁺ 2 ⁺		
4791.5	0 ⁺	4177.7 5	100	613.727	2 ⁺		
4811.5	2 ⁺	3503.6 5 4811.1 3	52 18 100 13	1308.644 0.0	2 ⁺ 0 ⁺		
4819.2	(9 ⁻)	1269.0 5	100	3550.15	(7 ⁻)	[E2]	B(E2)(W.u.)=10 4
4857.0	(9 ⁺)	1152.9 4 1273.2 ^b 5	100 6 50 13	3704.0 3585.0	(7 ⁺) 8 ⁺	[E2]	B(E2)(W.u.)=9 4
4957.3	1,2 ⁽⁺⁾	4957.1 3	100	0.0	0 ⁺		
4972.3	1 ⁻	4972.1 3	100	0.0	0 ⁺		
4998.3		3499.6 5	100	1498.599	0 ⁺		
5004.65	1,2 ⁽⁺⁾	3245.6 & 4 4391.2 3 5003.5 6	81 & 24 100 10 19 5	1758.689 613.727 0.0	0 ⁺ 2 ⁺ 0 ⁺		
5022.14		3168.14 & 17	100 &	1853.927	3 ⁺		

Adopted Levels, Gammas (continued) $\gamma(^{78}\text{Se})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. [‡]	Comments
5029.63	2 ⁺	3720.8 4	100 27	1308.644	2 ⁺		
		5029.5 3	100 18	0.0	0 ⁺		
5090.8		4476.9 3	100	613.727	2 ⁺		
5094.8		1046.8 6	100 17	4048.0	8 ⁻		
5101.9		4488.0 5	100	613.727	2 ⁺		
5126.52	(2,3,4)	3131.8 4	50 9	1995.897	2 ⁺		
		3272.13 19	100 14	1853.927	3 ⁺		
		3624.1 & 4	91 & 14	1502.825	4 ⁺		
5164.05		3168.14 & 17	46 & 8	1995.897	2 ⁺		
		3855.0 & 4	100 & 10	1308.644	2 ⁺		
5180.75	1 ⁽⁺⁾ ,2 ⁽⁺⁾	3326.4 3	100 10	1853.927	3 ⁺		
		3682.4 3	76 10	1498.599	0 ⁺		
5290.22	1,2 ⁽⁺⁾	3791.7 3	79 14	1498.599	0 ⁺		
		4676.2 3	100 14	613.727	2 ⁺		
		5290.0 3	86 14	0.0	0 ⁺		
5295.2	3 ⁻	4681.3 3	100	613.727	2 ⁺		
5339.7	1,2 ⁽⁺⁾	3840.9 3	100 16	1498.599	0 ⁺		
		4031.3 6	47 6	1308.644	2 ⁺		
5356.51	(2 ⁺)	3360.50 20	100 14	1995.897	2 ⁺		
		4742.7 3	67 14	613.727	2 ⁺		
5391.0		4777.1 3	100	613.727	2 ⁺		
5440.3		4826.4 3	100	613.727	2 ⁺		
5451.2	1,2 ⁽⁺⁾	3952.5 4	100	1498.599	0 ⁺		
5513.26	1,2 ⁽⁺⁾	3245.6 & 4	122 & 37	2267.07			
		4015.0 3	100 15	1498.599	0 ⁺		
		5512.9 3	35 7	0.0	0 ⁺		
5689.1		902.2 6	100	4786.9	(10 ⁺)		
5783.8	(12 ⁺)	1158.7 ^a 5	100 ^a	4625.1	(10 ⁺)	[E2] B(E2)(W.u.)<23	

[†] Weighted averages of all available data. For low-spin (up to about spin 4), the values are available from ⁷⁸As β^- decay; ⁷⁸Br ϵ decay; ($\alpha,2n\gamma$); (n,γ) E=thermal and ($n,n'\gamma$).

[‡] From $\gamma(\theta)$, $\gamma(\text{lin pol})$ and ce data (for a few transitions only) in ($\alpha,2n\gamma$) for transitions from high-spin ($J>4$) states. The multipolarity and mixing ratios for transitions from low-spin states (J up to about 4) are from $\gamma(\theta)$, $\gamma(\text{circ pol})$ and ce measurements in (n,γ) E=thermal; and some from $\gamma(\theta)$ in ($n,n'\gamma$).

γ only from (n,γ) E=thermal.

@ Total theoretical internal conversion coefficients, calculated using the BrIcc code (2008Ki07) with Frozen orbital approximation based on γ -ray energies, assigned multiplicities, and mixing ratios, unless otherwise specified.

& Multiply placed with undivided intensity.


^a Multiply placed with intensity suitably divided.

^b Placement of transition in the level scheme is uncertain.

Adopted Levels, Gammas

Level Scheme

Intensities: Relative photon branching from each level
 & Multiply placed: undivided intensity given
 @ Multiply placed: intensity suitably divided




Adopted Levels, Gammas

Level Scheme (continued)

Legend

Intensities: Relative photon branching from each level
& Multiply placed: undivided intensity given
@ Multiply placed: intensity suitably divided


-----> γ Decay (Uncertain)



Adopted Levels, Gammas

Level Scheme (continued)

Intensities: Relative photon branching from each level
 & Multiply placed: undivided intensity given
 @ Multiply placed: intensity suitably divided




Adopted Levels, Gammas

Level Scheme (continued)

Legend

Intensities: Relative photon branching from each level
 & Multiply placed: undivided intensity given
 @ Multiply placed: intensity suitably divided

-----▶ γ Decay (Uncertain)




⁷⁸Se₄₄

Adopted Levels, Gammas

Level Scheme (continued)

Intensities: Relative photon branching from each level
& Multiply placed: undivided intensity given
@ Multiply placed: intensity suitably divided




Adopted Levels, Gammas

Level Scheme (continued)

Legend

Intensities: Relative photon branching from each level
& Multiply placed: undivided intensity given
@ Multiply placed: intensity suitably divided

-----> γ Decay (Uncertain)



⁷⁸Se₄₄


Adopted Levels, Gammas

Level Scheme (continued)

Legend

Intensities: Relative photon branching from each level
& Multiply placed: undivided intensity given
@ Multiply placed: intensity suitably divided

-----▶ γ Decay (Uncertain)



Adopted Levels, Gammas

Level Scheme (continued)


Intensities: Relative photon branching from each level

& Multiply placed: undivided intensity given

@ Multiply placed: intensity suitably divided

Legend

-----▶ γ Decay (Uncertain)



⁷⁸Se₄₄

Adopted Levels, Gammas

