

Adopted Levels, Gammas

Type	Author	History Citation	Literature Cutoff Date
Full Evaluation	Balraj Singh	NDS 105,223 (2005)	22-Jun-2005

Q(β^-)=-1870.5 4; S(n)=9913.4 13; S(p)=11412 6; Q(α)=-6971.5 13 [2012Wa38](#)Note: Current evaluation has used the following Q record \$ -1870.5 3 9913.7 16 11412 5 -6971.8 16 [2003Au03](#).

Other reactions:

 $^{80}\text{Se}(e,e)$: [1988Kh02](#). $^{82}\text{Se}(\gamma,2n)$ GDR: [1976Ca06](#). $^{82}\text{Se}(n,3n)$: [1975FrZW](#).**Additional information 1.** $^{80}\text{Se}(d,^3\text{He})$: [1983Ro08](#) (g.s. proton occupation number for ^{80}Se). $^{79}\text{Se}(n,\gamma)$ resonances: [1979EnZZ](#), [1976Ca06](#), [1969Ma15](#), [1964Co31](#), [1962Ju01](#).Mass measurements: [1985El01](#) (also [1984EIZY](#)), [1977De20](#), [1964Ba03](#), [1963Ri07](#).IBM description of even-even Se isotopes: [1996Ra44](#).Nuclear structure theory (levels in ^{80}Se): [2004Da36](#). **^{80}Se Levels**Deformation parameters are available from (p,p'), (n,n'), (α,α') and Coul. ex. datasets. Only selected values are given here. See (p,p') for such data on many levels.**Cross Reference (XREF) Flags**

A	^{80}As β^- decay (15.2 s)	F	$^{80}\text{Se}(p,p')$,(pol p,p')	K	$^{80}\text{Se}(\alpha,\alpha')$
B	Muonic atom	G	$^{80}\text{Se}(p,p'\gamma),(\alpha,\alpha'\gamma)$	L	Coulomb excitation
C	^{80}Br ε decay (17.68 min)	H	$^{80}\text{Se}(n,n')$	M	$^{82}\text{Se}(p,t)$
D	$^{78}\text{Se}(t,p)$	I	$^{80}\text{Se}(n,n'\gamma)$	N	$^{176}\text{Yb}(^{28}\text{Si},X\gamma),(^{30}\text{Si},X\gamma)$
E	$^{80}\text{Se}(\gamma,\gamma')$	J	$^{80}\text{Se}(d,d')$,(pol d,d)		

E(level) [†]	J [‡]	T _{1/2} ^{&}	XREF	Comments
0.0 ^b	0 ⁺	stable	ABCDEFGHIJKLMN	%2 β^- =? $\langle r^2 \rangle^{1/2}=4.1399$ fm 19 (2004An14). 2 β decay: theoretical calculations: 2005Do07 , 2001Ka15 , 2000Bo05 . No experimental information is available. Additional information 2. B(E2) $\uparrow=0.253$ 6 (2001Ra27); $\beta_2=0.2318$ 27 (2001Ra27) $\mu=0.87$ 5 (1998Sp03) $Q=-0.31$ 7 (1977Le11 , 1989Ra17) J^π : L(t,p)=L(p,p')=2. T _{1/2} : from B(E2) taken from evaluation of 2001Ra27 . Other: 8.3 ps 8 (from (γ,γ'), 1976KaYY). μ : transient-field technique in Coul. Ex. (1998Sp03). Other: 0.84 24 (IMPAC in Coul. ex., 1969He11 , 1989Ra17). Q: reorientation effect in Coul. ex. (1977Le11). Other: -0.35 12 (1976VoZY). $\beta_2(p,p')=0.21$ (1993Mo05), 0.193 (1988Ba35 , 1986Og01), 0.22 1 (1986MoZR), 0.229 15 (1984De01), 0.195 30 (1983Ma59), 0.210 15 (1979Ma28), 0.234 (1970He10). $\beta_2(n,n')=0.225$ (1990Go13), 0.244 10 (1988Ba35 , 1984Ku09), 0.265 20 or 0.293 25 (1984De01), 0.25 (1979Ef01 , 1976La12). $\beta_2(\alpha,\alpha')=0.255$ or 0.190 (1988Ba35). $\beta_2(\text{Coul. ex.})=0.232$ 2 (1977Le11), 0.224 2 (1974Ba80), 0.245 (1962St02).
666.27 ^b 7	2 ⁺	8.52 ps 21	ABCDEFGHIJKLMN	

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) **^{80}Se Levels (continued)**

E(level) [†]	J ^π [‡]	T _{1/2} ^{&}	XREF	Comments
1449.35 7	2 ⁺	1.95 ps 7	A DEF GHIJKL	$\mu=0.70$ 20 (1998Sp03) μ : transient-field technique in Coul. Ex. (1998Sp03). 1449 and 1479 are unresolved in (α, α') . J^π : $L(p, p')=2$ and $\gamma\gamma(\theta)$ in (γ, γ') . $\beta_2(p, p')=0.047$ (from $\beta_2 R=0.25$ (1986Og01)), 0.082 20 or 0.065 5 (1986MoZR). $\beta_2(\alpha, \alpha')=0.05$ (1988Ba35). β_2 (Coul. ex.)=0.054 (1974Ba80). $T_{1/2}$: other: 0.2 ps +24–3 (DSAM in $(n, n'\gamma)$). XREF: F(?). J^π : $(812\gamma)(666\gamma)(\theta)$ in ^{80}Br decay. Parity from $\log ft=5.3$ 5 from 1 ⁺ .
1478.82 9	0 ⁺	11.4 ps 17	A C EFG HIJKL	
1701.45 ^b 11	4 ⁺	0.66 ps 2	FGHI JKLM N	$\mu=2.7$ 10 (1998Sp03) μ : transient-field technique in Coul. Ex. (1998Sp03). J^π : $L(p, p')=4$. $\beta_4(p, p')=-0.033$ (from $\beta_4 R=-0.18$ (1986Og01)), -0.026 8 or -0.034 10 (1983Ma59). Others: 1984De01 , 1986MoZR . $\beta_4(\alpha, \alpha')=0.07$ or -0.02 (1988Ba35). $T_{1/2}$: other: 0.7 ps +10–4 (DSAM in $(n, n'\gamma)$). J^π : $L(t, p)=0$ but $L(p, p')=2$. $\gamma\gamma(\theta)$ in (γ, γ') gives $J=0$ or 2; 0 ⁺ supported by comparison of experimental and theoretical yields in $(n, n'\gamma)$.
1873.40 12	(0) ⁺		A DEFG IJ	J^π : $\gamma\gamma(\theta)$ in (γ, γ') and $L(p, p')=2$. $T_{1/2}$: from DSAM in $(n, n'\gamma)$. Other: 2.8 ps +14–7 or 7 ps +9–3 (from B(E2) in Coul. ex.). XREF: D(2150?)J(2150).
1959.82 9	2 ⁺	0.38 ps +22–12	A D FG IJ L	J^π : from comparison of experimental and theoretical yields in $(n, n'\gamma)$.
2121.12 14	(3 ⁺)		D FG IJ	J^π : from comparison of experimental and theoretical yields in $(n, n'\gamma)$.
2311.29 9	(2 ⁺)	0.152 ps +28–14	A EFG Ij	J^π : from comparison of experimental and theoretical yields in $(n, n'\gamma)$.
2344.17 9	(1 ^{+,2+})	0.35 ps +17–10	D FG Ij	J^π : $L(t, p)=(2)$; 1 ⁺ from comparison of experimental and theoretical yields in $(n, n'\gamma)$.
2494.77 23	(4 ⁺)	1.1 ps 7	FG Ij	J^π : $L(p, p')=4$.
2513.57 10	(2 ⁺)	0.048 ps 7	A DEFG Ij	XREF: F(?). J^π : $L(t, p)=1$, but $\gamma\gamma(\theta)$ in (γ, γ') suggests $J=2$; 2 ⁺ also supported from comparison of experimental and theoretical yields in $(n, n'\gamma)$.
2627.40 19	(0 ⁺)		E I	J^π : primary transition in (γ, γ') from 1 ^(−) ; 0 ⁺ from comparison of experimental and theoretical yields in $(n, n'\gamma)$. $B(E3)\hat{\uparrow}=0.030$ 10 (2002Ki06) J^π : $L(p, p')=L(t, p)=3$.
2716.65 11	3 ⁻	0.38 ps 14	D FG HIJ L	$B(E3)$ adopted in evaluation by 2002Ki06 from (p, p') (1993Mo05 , 1986Og01 , 1979Ma28). Other: $B(E3)=0.0084$ 14 from Coul. ex. (1974Ba80). Average β_3 (from inelastic scattering)=0.154 from $\beta_3(\alpha, \alpha')=0.161$ (1988Ba35); $\beta_3(n, n')=0.151$ 10 (from $b_3 r=0.78$ 5, 1984Ku09); $\beta_3(p, p')=0.163$ (1993Mo05), 0.124 (deduced by 1988Ba35 from 1986Og01), 0.144 (deduced by 1988Ba35 from 1984De01), 0.17 1 (1986MoZR), 0.167 (1979Ma28). β_3 (from B(E3) in Coul. ex.)=0.083.
2774.3 10	(1,2 ⁺) [@]		A	
2787? 5			F	
2814.50 16	(2 ^{+,1+})		EF Ij	XREF: F(2819).
2825.55 23	(6 ⁺)		Ij	J^π : 2 ⁺ from $\gamma\gamma(\theta)$ in (γ, γ') and $L(p, p')=(2)$; 1 ⁺ from comparison of experimental and theoretical yields in $(n, n'\gamma)$. J^π : γ to 4 ⁺ . 6 ⁺ from comparison of experimental and

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) **^{80}Se Levels (continued)**

E(level) [†]	J [‡]	T _{1/2} ^{&}	XREF	Comments
2826.99 11	(2 ⁺)	0.18 ps 4	E G Ij	theoretical yields in (n,n'γ). J ^π : γγ(θ) in (γ,γ') and γ to 0 ⁺ . Parity from reduced strength for E1 transition in (γ,γ').
2836.3 10	(1,2 ⁺) [@]		A j	
2895.5 ^b 10	(6 ⁺) ^a		F I N	
2947.54 15	(2 ^{+,4⁺})	0.18 ps +11-6	F I	J ^π : L(p,p')=(2); 4 ⁺ from comparison of experimental and theoretical yields in (n,n'γ).
2998? 5			F	
3025.17 16	(1 ^{+,2⁺)[@]}	0.049 ps 14	A G I	J ^π : 1 ⁺ from comparison of experimental and theoretical yields in (n,n'γ).
3033 4	(4 ⁺)		F J	J ^π : L(p,p')=4.
3036 10	(6 ⁺)		d	J ^π : L(t,p)=(2+6).
3037.74 13	(1 ^{+,2⁺)}	0.13 ps +9-5	d I	E(level): doublet in (t,p). J ^π : L(t,p)=(2+6) and γ to 0 ⁺ ; 1 ⁺ from comparison of experimental and theoretical yields in (n,n'γ).
3125.79 16	(2 ⁺) [#]	0.028 ps 14	EF I	T _{1/2} : from DSAM in (n,n'γ) (1989Do14); not given by 1999Ko46.
3160 9	0 ⁺		D	J ^π : L(t,p)=0.
3176.92 19	(1,2 ^{+)@}		F I	
3199.4 3	(2) [#]		EF I	XREF: F(?).
3224.28 19	(1,2)	0.070 ps 28	I	J ^π : γ to 0 ⁽⁺⁾ .
3226 4	(4 ⁺)		F	J ^π : L(p,p')=4.
3248.3 5	(2 ^{+)#}		E	
3280.4 4	(1,2 ^{+)@}		d I	
3284 4	(3 ⁻)		d F	J ^π : L(p,p')=3.
3314? 5			F j	
3316.4 10	(0) [#]		EF j	XREF: F(?).
3349.95 20	(1 ⁺)		E I	J ^π : from γγ(θ) in (γ,γ').
3354 4	(3 ⁻)		D F J	XREF: J(3370).
3390.75 24	(2 ⁺)		DEF j	J ^π : L(p,p')=3 and L(t,p)=(3). XREF: j(3370). J ^π : L(t,p)=(2).
3441.88 22	(0 ^{+)#}		EF I	J ^π : L(p,p')=2 but γγ(θ) in (γ,γ') suggests 0 ⁺ . XREF: D(3484).
3491 5			D F	
3567 5			F	
3606.4 3	(2) [#]		A E	
3619.7 4	(0 ^{+,2^{+)#}}		dEF	XREF: d(3635). J ^π : L(t,p)=0 for a 3635 group suggests J ^π =0 ⁺ for 3620 or 3640 level, but L(p,p')=(2) suggests 2 ⁺ .
3635.5 ^b 15	(8 ^{+)a}		N	
3640 5			d F	XREF: d(3635).
3655.4 10	(0,1,2)		E	J ^π : primary transition from 1 ⁽⁻⁾ .
3675 5			F	
3727.2 5	(0,1,2)		A	J ^π : log ft=6.1 from 1 ⁽⁺⁾ . XREF: d(3760).
3753 4	(3 ⁻)		d F j	J ^π : L(p,p')=3. Also L(t,p)=(3) for a 3760 10 group. XREF: d(3760).
3774? 5			d F j	
3813.7 4	(6 ⁺)		I	J ^π : γ to 4 ⁺ ; comparison of experimental and theoretical yields in (n,n'γ).
3814.9 5	(8 ⁺)		I	
3826 5			F	
3845? 10			F	

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) **^{80}Se Levels (continued)**

E(level) [†]	J π [‡]	XREF	Comments		
3870.0 4	(1 $^-$)	DEF	J π : L(t,p)=(1).		
3931 4	(2 $^+$)	F	J π : L(p,p')=(2).		
3951.9 4	(2 $^+$)	EF	XREF: F(3960).		
			J π : L(p,p')=(2) for a 3960 4 group.		
3976 8	(1 $^-$)	D	J π : L(t,p)=(1).		
3996 4	(5 $^-$)	F	J π : L(p,p')=5.		
4039 4		F			
4047.1 5	(2 $^+$)	D I	XREF: D(4063).		
			J π : L(t,p)=(2).		
4062.2 4	(0 $^+)^{\#}$	EF	XREF: F(?).		
4129 8	0 $^+$	D	J π : L(t,p)=0.		
4130 4	(3 $^-$)	F	J π : L(p,p')=3.		
4173 4	2 $^+$	D F J	XREF: J(4180).		
			J π : L(t,p)=2.		
4225 4		F			
4247 7	2 $^+$	D	J π : L(t,p)=2.		
4295 4		F			
4322 4	(2 $^+$)	D F	J π : L(t,p)=(2).		
4352 4	2 $^+$	D F	J π : L(t,p)=2.		
4420 4	(2 $^+$)	F	J π : L(p,p')=(2).		
4436.6 4	(5 $^-$)	F I	J π : L(p,p')=5.		
4464 5	(1 $^-$)	D	J π : L(t,p)=1.		
4511 4	(4 $^+$)	F	J π : L(p,p')=4.		
4570 4		F			
4673.5 ^b 18	(10 $^+)^{\#}$	N			
4682 4	(4 $^+$)	D F	XREF: D(4712).		
			J π : L(p,p')=4.		
4950 4		F			
4993 4		F			
5180 30		D			
5325 4	(3 $^-$)	F	J π : L(p,p')=3.		
7818.52 9	1 $^{(-)}$	E	J π : γ to 0 $^+$. Parity from reduced strength for E1 transition in (γ, γ').		

[†] From least-squares fit to E γ 's for levels populated in γ -ray studies. For others weighted averages of values available from different reactions have been taken.

[‡] Above 2 MeV excitation energy, J π 's deduced from L(p,p') are given in parentheses due to high level density, ambiguity in level correspondence between different reactions, and tentative nature of L value.

[#] From $\gamma\gamma(\theta)$ in (γ, γ'). Parity is from a comparison of reduced strengths for E1 and M1 transitions with systematics of known E1 and M1 transitions in this mass region. The reduced strengths have been calculated by [1973Sz04](#) from relative intensities corrected for energy dependence, average level spacing and partial widths for the g.s. and the excited levels J π assignments based on (γ, γ') study are considered tentative; first, because $\gamma(\theta)$ data are reported at only two angles and, second because transitions are assumed pure dipole with no quadrupole admixture.

[@] γ to 0 $^+$.

[&] From B(E2) values in Coul. ex. for levels below 1900 keV. Above this, values are from DSA method in (n,n' γ) ([1999Ko46](#)).

^a Systematics of yrast sequences in even-even nuclides populated in heavy-ion reactions.

^b Band(A): Yrast sequence.

Adopted Levels, Gammas (continued)

$\gamma(^{80}\text{Se})$								
$E_i(\text{level})$	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. $\&$	$\delta\&$	Comments
666.27	2 ⁺	666.15 10	100	0.0	0 ⁺	E2 ^a		B(E2)(W.u.)=24.7 6
1449.35	2 ⁺	783.1 1	66.6 10	666.27	2 ⁺	E2+M1	-5 +2-6	B(M1)(W.u.)=0.0004 3; B(E2)(W.u.)=18.5 10 Mult., δ , from $\gamma(\theta)$ in Coul. ex.
								$\delta=-0.71 +12-17$ is also possible but less likely from systematics of second 2 ⁺ states in even-even nuclei.
1478.82	0 ⁺	1449.4 1	100 3	0.0	0 ⁺	[E2]		B(E2)(W.u.)=1.33 7
1701.45	4 ⁺	812.4 1	100	666.27	2 ⁺	E2 ^a		B(E2)(W.u.)=6.9 11
1873.40	(0) ⁺	1035.1 1	100	666.27	2 ⁺	E2		B(E2)(W.u.)=35.2 11
1959.82	2 ⁺	1207.1 1	100	666.27	2 ⁺			
		1293.7 2	100 5	666.27	2 ⁺	M1+E2	-1.1 +6-11	δ : from M1 and E2 matrix elements in Coul. Ex. (1995Ka29). Other: -0.31 5 or +10 +10-2 from $\gamma(\theta)$ in ($n,n'\gamma$). B(E2)(W.u.)=0.9 +3-6
2121.12	(3 ⁺)	1959.9 1	55 5	0.0	0 ⁺	[E2]		
		671.7 2	15 3	1449.35	2 ⁺			
		1454.9 2	100 8	666.27	2 ⁺			
2311.29	(2 ⁺)	861.9 1	15 5	1449.35	2 ⁺			
		1645.0 1	100 12	666.27	2 ⁺	D+Q		δ : +1.95 7 or -0.09 +2-6 from $\gamma(\theta)$ in ($n,n'\gamma$).
2344.17	(1 ^{+,2⁺)}	470.5 4	55 9	1873.40	(0) ⁺			
		894.8 ^b 1	100 9	1449.35	2 ⁺			
		1677.9 ^b 1	55 9	666.27	2 ⁺			
2494.77	(4 ⁺)	2344 ^b 1	9.1 18	0.0	0 ⁺			
		793.0 3	100 30	1701.45	4 ⁺	M1+E2	+1.1 1	B(M1)(W.u.)=0.012 9; B(E2)(W.u.)=28 21 δ : from $\gamma(\theta)$ in ($n,n'\gamma$) and T _{1/2} (2495 level).
2513.57	(2 ⁺)	1046 ^b	≈3	1449.35	2 ⁺	[E2]		
		1828.8 3	53 5	666.27	2 ⁺	[E2]		B(E2)(W.u.)=0.4 3
		813.3 ^b 2		1701.45	4 ⁺			
		1035.7 ^b 4	≈40	1478.82	0 ⁺			
								Reported in (γ,γ') only. The placement is considered suspect since with the quoted intensity in (γ,γ'), it would have been seen in ⁸⁰ As β^- decay and in ($n,n'\gamma$).
2627.40	(0 ⁺)	1063.8 4	4.3 14	1449.35	2 ⁺			
2716.65	3 ⁻	1847.3 1	100 9	666.27	2 ⁺			B(E2)(W.u.)=0.17 7
		2513.4 2	4.3 14	0.0	0 ⁺	[E2]		
		1178.2 ^b 2	100	1449.35	2 ⁺			
		405.1 3	7.7 23	2311.29	(2 ⁺)	[E1]		B(E1)(W.u.)=0.0010 5
		1015.1 2	7.7 15	1701.45	4 ⁺	[E1]		B(E1)(W.u.)=6.E-5 3
		2050.4 1	100 8	666.27	2 ⁺	[E1]		B(E1)(W.u.)=0.00010 4
		(2716.6)	0.15 7	0.0	0 ⁺	[E3]		B(E3)(W.u.)=10 6
								I γ : deduced (evaluator) from T _{1/2} and B(E3) for 2717 level.
2774.3	(1,2 ⁺)	2774.2 10	100	0.0	0 ⁺			
2814.50	(2 ^{+,1⁺)}	2148.0 ^b 3	29 14	666.27	2 ⁺			
		2814.6 2	100 14	0.0	0 ⁺			
2825.55	(6 ⁺)	1124.1 2	100	1701.45	4 ⁺			E γ : from (γ,γ'). E γ =2817.7 in ($n,n'\gamma$).
2826.99	(2 ⁺)	2160.7 1	100 15	666.27	2 ⁺			
		2826.9 3	7.7 24	0.0	0 ⁺	[E2]		
2836.3	(1,2 ⁺)	2836.2 10	100	0.0	0 ⁺			B(E2)(W.u.)=0.061 25

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) **$\gamma(^{80}\text{Se})$ (continued)**

E_i (level)	J^π_i	E_γ^\dagger	L_γ^\dagger	E_f	J^π_f	Mult. ^{&}	Comments
2895.5	(6 ⁺)	1194		1701.45	4 ⁺		
2947.54	(2 ^{+,4⁺)}	826.4 2 1498.1 2 2281.4 3	50 17 100 33 67 33	2121.12 1449.35 666.27	(3 ⁺) 2 ⁺ 2 ⁺		
3025.17	(1 ^{+,2⁺)}	1577.6 [‡] 3 2358.2 2 3024.8 3	50 17 100 25 30 20	1449.35 666.27 0.0	2 ⁺ 2 ⁺ 0 ⁺		E_γ : poor fit. Level-energy difference=1575.8. E_γ : level-energy difference=2358.86.
3037.74	(1 ^{+,2⁺)}	1078.6 2 1558.7 2 1587.9 2	100 20 80 20 56 12	1959.82 1478.82 1449.35	2 ⁺ 0 ⁺ 2 ⁺		E_γ : level-energy difference=1077.9.
3125.79	(2 ⁺)	1677.0 ^{‡b} 5 2459.3 2	≈1 100	1449.35 666.27	2 ⁺ 2 ⁺		
3176.92	(1,2 ⁺)	1697.8 5 3176.9 2	70 20 100 20	1478.82 0.0	0 ⁺ 0 ⁺		
3199.4	(2)	3199.5 [‡] 5	100	0.0	0 ⁺		
3224.28	(1,2)	1522.8 2 1745.5 3	100 13 43 22	1701.45 1478.82	4 ⁺ 0 ⁺		
3280.4	(1,2 ⁺)	2614.5 5 3280.0 5	73 21 100 27	666.27 0.0	2 ⁺ 0 ⁺		
3349.95	(1 ⁺)	3348.4 5	100	0.0	0 ⁺		
3390.75	(2 ⁺)	1909.9 5 1941.9 5	100 20 100 20	1478.82 1449.35	0 ⁺ 2 ⁺		E_γ : poor fit. Level-energy difference=1911.9.
3441.88	(0 ⁺)	1097 [‡] 1 2775.9 3	80 20 100 30	2344.17 666.27	(1 ^{+,2⁺) 2⁺}		
3606.4	(2)	2156.9 [#] 5 2940.3 [#] 10	100 50	1449.35	2 ⁺		
3619.7	(0 ^{+,2⁺)}	2953.7 5	100	666.27	2 ⁺		
3635.5	(8 ⁺)	740		2895.5	(6 ⁺)		
3727.2	(0,1,2)	1415.9 5 3060.8 ^b 20	100 50 50 50	2311.29 666.27	(2 ^{+) 2⁺)}		
3813.7	(6 ⁺)	2112.2 3	100	1701.45	4 ⁺		
3814.9	(8 ⁺)	989.3 4	100	2825.55	(6 ^{+) 0⁺)}		
3870.0	(1 ⁻)	2391.9 5	100	1478.82	0 ⁺		
3951.9	(2 ⁺)	3286.1 5	100	666.27	2 ⁺		
4047.1	(2 ⁺)	2597.7 5	100	1449.35	2 ⁺		
4062.2	(0 ⁺)	2612.7 5	100	1449.35	2 ⁺		
4436.6	(5 ⁻)	1941.8 3	100	2494.77	(4 ^{+) 0⁺)}		
4673.5	(10 ⁺)	1038		3635.5	(8 ^{+) 0⁺)}		
7818.52	1 ⁽⁻⁾	3756.1 4 3866.9 4 3949.1 5 4163 1 4199.1 5 4212.0 4 4376.8 3 4427.1 3 4468.2 2 4502 1 4570.1 5 4619.1 3 4692.4 2 4991.4 2 5004.3 5 5191.6 4	4.3 4 3.0 5 3.0 4 1.3 3 2.8 3 3.7 3 5.2 4 8.5 3 9.2 4 2.2 4 7.3 3 5.5 3 12.5 3 12.4 4 3.5 3 1.0 3	4062.2 3951.9 3870.0 3655.4 3619.7 3606.4 3441.88 3390.75 3349.95 3316.4 3248.3 3199.4 3125.79 2826.99 2814.50 2627.40	(0 ^{+) 0⁺) (2^{+) 0⁺) (1⁻) (0,1,2) (0^{+,2⁺) (2) (0⁺) (2^{+) 0⁺) (1^{+) 0⁺) (0) (2^{+) 0⁺) (2) (2^{+) 0⁺) (2^{+) 0⁺) (2^{+,1⁺) (0⁺)}}}}}}}}}	(E1) (E1) (E1) (E1) (E1) (E1) (E1) (E1) (E1) (E1) (E1) (E1) (E1) (E1) (E1) (E1) (E1)	

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) $\gamma(^{80}\text{Se})$ (continued)

E_i (level)	J_i^π	E_γ^\dagger	I_γ^\dagger	E_f	J_f^π	Mult. ^{&}
7818.52	$1^{(-)}$	5304.4 3	6.4 3	2513.57	(2^+)	
		5507.2 7	4.2 5	2311.29	(2^+)	
		5858.4 2	27.8 3	1959.82	2^+	(E1)
		5944.7 8	1.1 2	1873.40	$(0)^+$	
		6339.4 1	9.4 2	1478.82	0^+	
		6369.4 3	8.4 2	1449.35	2^+	
		7818.9 5	100.0 5	0.0	0^+	(E1)

[†] Weighted averages taken when data of comparable precision are available from more than one dataset.

[‡] Reported in $(n,n'\gamma)$ only.


[#] Reported in ^{80}As β^- only.

[@] Reported in $(p,p'\gamma)$ only.

[&] From $\gamma(\theta)$ in $(n,n'\gamma)$ and RUL deduced from $T_{1/2}$. Mult=E1 for transitions from 7819 level is from $\gamma(\theta)$ in (γ,γ') and transition strengths.

^a From $(813\gamma)(666\gamma)(\theta)$ in ^{80}Br ε decay and $T_{1/2}$ (levels).

^b Placement of transition in the level scheme is uncertain.




Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level


- - - - - γ Decay (Uncertain)

Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level

- - - - - γ Decay (Uncertain)

Adopted Levels, Gammas

Band(A): Yrast sequence

