

Adopted Levels, Gammas

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	B. Singh, A. Negret, and K. Zuber		NDS 110,2815 (2009)	30-Sep-2009

Q(β^-)=-6756 5; S(n)=11923 7; S(p)=8868 3; Q(α)=-5181.2 15 [2012Wa38](#)

Note: Current evaluation has used the following Q record -6757 5 11923 7 8867.5 27-5181.6 16 [2009AuZZ](#).

[Additional information 1](#).

Values in [2003Au03](#) are: Q=-6490 90, S(n)=11920 11, S(p)=8858 7, Q(α)=-5176 4.

⁸⁴Sr evaluated by B. Singh, A. Negret, and K. Zuber.

Theory/calculations:

[Additional information 2](#).

[1997Su08](#): energies of ground state and γ band members, IBA.

[1989Sa38](#): collective bands.

[1989Co02](#): octupole excitation.

[1986Ga04](#), [1979Bu20](#): nuclear deformation and potential energy surfaces.

[1985Na02](#): microscopic study of high-spin states.

[1982De05](#), [1983Bu09](#), [1984He07](#): interacting-boson model.

[1971Ki16](#), [1973Og01](#): shell-model calculations.

Other experiments:

Atomic mass measurements using Penning-trap system: [2007Ke09](#).

Measurements of isotope shift and mean square charge radius: [1992Ba55](#), [1990Bu12](#), [1988Si06](#), [1987An02](#), [1986An39](#), [1986Ea01](#), [1986Ma43](#), [1985Bu20](#), [1983El04](#), [1983Bo35](#), [1983Lo13](#).

⁸⁴Sr Levels

Cross Reference (XREF) Flags

A	⁸⁴ Rb β^- decay (32.82 d)	F	⁵⁹ Co(²⁸ Si,3p γ)	K	⁸⁴ Sr(d,d')
B	⁸⁴ Y ε decay (39.5 min)	G	⁷⁶ Ge(¹² C,4n γ), ⁸¹ Br(⁶ Li,3n γ)	L	⁸⁴ Sr(α , α'), (α , α' γ)
C	⁸⁴ Y ε decay (4.6 s)	H	⁸² Kr(³ He,n)	M	Coulomb excitation
D	⁵¹ V(³⁶ S,p2n γ)	I	⁸² Kr(α ,2n γ)	N	⁸⁵ Rb(p,2n γ)
E	⁵² Cr(³⁶ S,2p2n γ)	J	⁸⁴ Sr(p,p'), (p,p' γ)	O	⁸⁶ Sr(p,t)

E(level) [†]	J ^π	T _{1/2} [‡]	XREF	Comments
0.0 ^{&}	0 ⁺	stable	ABCDEFGHIJKLMNO	$\langle r^2 \rangle^{1/2}$ =4.2364 fm 17 (2004An14 evaluation). T _{1/2} : >7.3×10 ¹³ y (1952Fr23 , double β decay). Other:>10 ¹⁷ y probably for neutrino-less double β/ε decay, preliminary result from H.J. Kim, presented at 16th Int. Conf. on Supersymmetry and the Unification of Fundamental Interactions, Seoul, June 2008 Communication with the author on April 16, 2009 revealed that the analysis of this experiment is still in progress. J ^π : L(p,t)=0.
793.22 ^{& 6}	2 ⁺	3.23 ps 35	BCDEFG IJKLMNO	μ =+0.84 9 (1988Ku01 , 1989Ra17) μ : from g-factor=+0.419 47 measured in Coulomb ex. (1988Ku01). See also 2005St24 compilation. T _{1/2} : weighted average of 3.19 ps 35 (1982De05) and 4.2 ps +28-14 (1980Ek03). Other: 6.2 ps 21 (1982De05 value reanalyzed by 1994Ch28) 2001Ra27 evaluation gives adopted T _{1/2} =3.2 ps 5 and B(E2)($\uparrow$)=0.289 44. J ^π : E2 γ to 0 ⁺ ; L(p,t)=2.
1453.93 ^{d 10}	2 ⁺		BC G IJ L NO	J ^π : M1+E2 γ to 2 ⁺ ; $\gamma(\theta)$ not consistent with $\Delta J=1$ and δ . Also L(p,t)=(2).

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{84}Sr Levels (continued)

E(level) [†]	J ^π	T _{1/2} [‡]	XREF	Comments
1504.2 10	0 ⁺		BC J L O	J ^π : L(p,t)=0.
1767.69 & 9	4 ⁺	1.4 ps 4	B DEFG IJ L NO	J ^π : ΔJ=2, E2 to 2 ⁺ . L(p,t)=(4) is consistent. T _{1/2} : unweighted average of 1.73 ps 21 (1982De05) and 0.97 ps 28 (1980Ek03). Other: 4.16 ps 14 (1982De05) value reanalyzed by 1994Ch28).
2056.07 ^d 11	(3) ⁺		B G I N	J ^π : ΔJ=1, M1+E2 γ to 2 ⁺ ; γ to 4 ⁺ ; band member.
2071.6 8	0 ⁺		C J L O	J ^π : L(p,t)=0.
2297.93 14			G	
2390 5	2		O	J ^π : L(p,t)=2.
2448.11 ^c 11	3 ⁻		B G IJKL O	J ^π : L(p,t)=L(d,d')=3. Configuration=(g _{9/2} ,f _{5/2} ⁻¹) or (g _{9/2} ,p _{3/2} ⁻¹) (1982De05). B(E3)(↑)=0.043 18 (2002Ki06) evaluation, data from 1973Re01). Deduced B(E3)(W.u.)=15 6.
2525 5	(0 ⁺)		O	J ^π : L(p,t)=(0).
2598.23 ^d 22	(4 ⁺)		B G IJ L O	XREF: B(?). J ^π : 2 ⁺ or 4 ⁺ from 1145γ(θ) indicating ΔJ=0 or 2; J=4 favored by excitation function and band assignment.
2735.25 ^d 20	(5 ⁺)		B G J O	J ^π : ΔJ=1 γ to 4 ⁺ ; excitation function; band member.
2769.03 10	(5 ⁻)	9.5 @ ps 6	B DEFG IJ L NO	μ=+8.0 10 (1989Ku11,2005St24) μ: transient-field integral perturbed-angular correlation in ⁷⁴ Ge(¹² C,2nγ) (1989Ku11). J ^π : ΔJ=1, DIPOLE G TO 4 ⁺ ; L(p,t)=(5).
2807.87 & 11	6 ⁺	1.01 ps 21	B DEFG IJ L N	J ^π : ΔJ=2, E2 γ to 4 ⁺ . T _{1/2} : weighted average of 1.04 ps 21 (1982De05), 0.97 ps 28 (1994Ch28). Other: 2.6 ps 4 from (1982BrZO).
2886.99 14	2 ⁺		B J L O	J ^π : L(p,t)=2.
3041.25 ^c 13	(5 ⁻)		FG J L O	XREF: J(?)L(?). J ^π : ΔJ=(0), dipole γ to (5 ⁻); γ to 3 ⁻ ; L(p,t)=(4,5).
3098.67 13	6 ⁽⁺⁾		B G	J ^π : ΔJ=2 γ to 4 ⁺ ; γ to 6 ⁺ .
3157.05 ^d 22	(7 ⁺)		G	J ^π : ΔJ=2 γ to (5 ⁺); excitation function.
3175 5	(2 ⁺)		J L O	J ^π : L(p,t)=(2).
3255 30	3 ⁻		J L O	J ^π : L(p,t)=3.
3270.58 17	(4,5,6) ⁺		B G	J ^π : γ to 4 ⁺ ; M1,E2 γ to 6 ⁺ . The β feeding from (6 ⁺) disfavors 4.
3279.15 ^c 14	(6 ⁻)		FG I	J ^π : ΔJ=1 γ to (5 ⁻); band member.
3330 30	0 ⁺		H J L	J ^π : L(³ He,n)=0.
3331.91 ^b 13	8 ⁺	157 ps 5	DEFG I	μ=-1.2 6 (1981Br20,1989Ra17) J ^π : ΔJ=2, E2 γ to 6 ⁺ . μ: from g factor=-0.15 7 from spin precession in polarized hyperfine fields of a tilted multi-foil target (1981Br20). Other: -0.8 16 from g=-0.1 2 (1989Ku11, transient-field integral perturbed-angular correlation in ⁷⁴ Ge(¹² C,2nγ)) See also 2005St24 compilation. J ^π : Configuration=(vg _{9/2}) ⁻² ₈₊ ⊗(g.s. of ⁸⁶ Sr core) (1982De05). T _{1/2} : from 1982De05. Others: 163 ps 3 (1982De05) value reanalyzed by 1994Ch28), 170 ps 7 (1982BrZO).
3455 30			J L O	
3487.92 ^c 12	(7 ⁻)	4.4 @ ps 5	DEFG IJ	μ=+4.2 14 (1989Ku11,2005St24) μ: transient-field integral perturbed-angular correlation in ⁷⁴ Ge(¹² C,2nγ) (1989Ku11). J ^π : ΔJ=2, E2 γ to (5 ⁻); ΔJ=1 γ to 6 ⁺ . XREF: L(3520).
3511.77 16	(4 ⁺ ,5 ⁻)		B J L	J ^π : γ's to 3 ⁻ and 6 ⁺ ; β feeding from (6 ⁺) favors 5 ⁻ .

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{84}Sr Levels (continued)

E(level) [†]	J ^π	T _{1/2} [‡]	XREF	Comments
3578.23 [?] 25			B	
3650.15 ^c 13	(7 ⁻)		G L	J ^π : ΔJ(2) γ to (5 ⁻); ΔJ=1 γ to 6 ⁽⁺⁾ ; band member.
3679.94 ^a 13	8 ⁺	3.33 [@] ps 14	DEFG I	μ=+7.2 8 (1989Ku11,2005St24) J ^π : configuration=πg _{9/2} =2 g ₊ ⊗ ⁸² Kr core (1982De05). μ: transient-field integral perturbed-angular correlation in ⁷⁴ Ge(¹² C,2nγ) (1989Ku11).
3749.07 24	(7)		G	J ^π : ΔJ=1 γ to 6 ⁺ .
3750 30	(3 ⁻ ,4 ⁺)		O	J ^π : L(p,t)=(3,4).
3819.58 [?] 15			B	
3918.08 [?] 16			B	
3960 30			L	
4028.78 ^{&} 23	(8 ⁺)		G	J ^π : ΔJ=2 γ to 6 ⁺ .
4062.78 17	4 ⁺		B L O	XREF: L(?)O(4080). J ^π : L(p,t)=4. Note that 4 ⁺ is inconsistent with β feeding from (6 ⁺).
4260 30			L O	
4268.05 ^c 16	(8 ⁻)		FG	XREF: F(?).
4365.95 18	(4 ⁺)		B L	J ^π : ΔJ=2 γ to (6 ⁻); γ to (7 ⁻); band member. XREF: L(4360). J ^π : γ's to 2 ⁺ and 6 ⁺ . Note that (4 ⁺) is inconsistent with β feeding from (6 ⁺).
4370.4 ^d 3	(9 ⁺)		G	J ^π : ΔJ=(2) γ to (7 ⁺); excitation function; band member.
4447.61 ^b 14	10 ⁺	2.22 [@] ps 35	DEFG I	μ=+2.0 10 (1989Ku11,2005St24) J ^π : ΔJ=2, E2 γ to 8 ⁺ . configuration=(vg _{9/2}) ₈₊ ⁻² ⊗ (2 ⁺ of ⁸⁶ Sr core) (1982De05). μ: transient-field integral perturbed-angular correlation in ⁷⁴ Ge(¹² C,2nγ) (1989Ku11).
4534.06 ^a 15	10 ⁺	1.66 [@] ps 14	DEFG I	μ=+8.0 20 (1989Ku11,2005St24) μ: transient-field in ⁷⁴ Ge(¹² C,2nγ) (1989Ku11). J ^π : ΔJ=2, E2 γ to 8 ⁺ .
4540 30			L O	
4636.13 ^c 14	(9 ⁻)	2.5 [@] ps 4	DEFG I	μ=0.00 36 (1989Ku11,2005St24) J ^π : ΔJ=2, E2 γ to (7 ⁻). Configuration=vg _{9/2} ⁻² ⊗ (3 ⁻) (1982De05). μ: transient-field integral perturbed-angular correlation in ⁷⁴ Ge(¹² C,2nγ) (1989Ku11).
4660 30			L	
4740 30			L	
4745.72 24	(8,9,10 ⁺)		G	E(level): γ to 8 ⁺ . It is unlikely that this level is same as 4740 in (α,α').
5150.7 [?] 3			B	
5444.48 ^c 15	(11 ⁻)	7.5 [@] ps 10	DEFG	J ^π : ΔJ=2, E2 γ to (9 ⁻); γ to 10 ⁺ .
5653.25 ^a 16	12 ⁺	0.61 ps 21	DEFG	J ^π : ΔJ=2, E2 γ to 10 ⁺ . T _{1/2} : weighted average of 0.83 ps 28 (1982De05), 0.49 ps 21 (1994Ch28).
5891.6 ^b 10	(12 ⁺)	0.24 [#] ps 10	D F	J ^π : γ to 10 ⁺ ; band member.
6069.43 ^c 17	(12 ⁻)	0.42 [#] ps 14	EFG	J ^π : ΔJ=1, dipole γ to (11 ⁻); band member.
6484.34 ^c 21	(13 ⁻)	0.62 [#] ps 28	F	J ^π : γ to (12 ⁻); possible γ's to 12 ⁺ and (11 ⁻); band member.
6739.65 ^a 19	14 ⁺	0.42 [#] ps 14	DEFG	J ^π : ΔJ=2, E2 γ to 12 ⁺ ; band member.
6916.8 ^c 4	(14 ⁻)		F	J ^π : γ to (13 ⁻); band member.
7822.8 7	(15 ⁺)		D	J ^π : γ to 14 ⁺ and a low-energy γ from 16 ⁺ .
8006.4 ^a 5	16 ⁺	0.21 [#] ps 7	DEF	J ^π : γ to 14 ⁺ ; band member.

Continued on next page (footnotes at end of table)

Adopted Levels, Gammas (continued) ^{84}Sr Levels (continued)

E(level) [†]	J ^π	T _{1/2} [‡]	XREF	Comments
9098.4 8	(17 ⁺)		D	J ^π : γ's to 16 ⁺ and (15 ⁺).
9424.9 ^a 10	18 ⁺	0.14 [#] ps 6	DEF	J ^π : γ to 16 ⁺ ; band member.
11059.9 ^a 22	20 ⁺	<0.18 ps	DEF	J ^π : γ to 18 ⁺ ; band member. T _{1/2} : 0.14 ps 4, effective half-life from 1994Ch28, not corrected for side feeding.
12920 ^a 3	22 ⁺		DE	J ^π : γ to 20 ⁺ ; band member.
15080? ^a 4	(24 ⁺)		E	J ^π : possible γ to 22 ⁺ ; band member.

[†] Level energies with ΔE<5 keV are deduced from least-square fit to the adopted gammas. The others are from (p,t), (α,α'), or weighted averages from (p,t), (α,α'), and (p,p').

[‡] From Doppler-shift attenuation method (DSAM) and/or recoil-distance Doppler shift (RDDS) methods. Measurements are from 1994Ch28 using line-shape analysis in DSA in $^{59}\text{Co}(^{28}\text{Si},3p\gamma)$ reaction for levels above 5600 keV. For levels up to 5700 keV, measurements are from 1982De05 using recoil-distance Doppler-shift method in $^{76}\text{Ge}(^{12}\text{C},4n\gamma)$ reaction. For the 5653.5 level, values are measured in both studies. Values from recoil-distance method are also available from 1980Ek03 for 793 and 1768 levels using (α,2nγ) reaction and from 1982BrZO for 2808 and 3331 levels using $^{76}\text{Ge}(^{12}\text{C},4n\gamma)$ reaction.

[#] From 1994Ch28.

[@] From 1982De05.

[&] Band(A): g.s. band.

^a Band(B): $\pi(g_{9/2}^{-2})_{8+} \otimes (^{82}\text{Kr} \text{ core})$.

^b Band(C): $\nu(g_{9/2}^{-2})_{8+} \otimes (^{86}\text{Sr} \text{ core})$.

^c Band(D): Octupole band.

^d Band(E): quasi γ band.

Adopted Levels, Gammas (continued)

E _i (level)	J _i ^π	γ(⁸⁴ Sr)							Comments
		E _γ [†]	I _γ [†]	E _f	J _f ^π	Mult. ^a	δ ^b	α ^d	
793.22	2 ⁺	793.22 6	100	0.0	0 ⁺	E2		0.00106	B(E2)(W.u.)=26 3
1453.93	2 ⁺	660.85 9	100 3	793.22	2 ⁺	M1+E2	+0.59 5	0.00145	
		1453.9 3	13.3 9	0.0	0 ⁺				
1504.2	0 ⁺	711 [#]		793.22	2 ⁺				B(E2)(W.u.)=21 6
1767.69	4 ⁺	974.48 7	100	793.22	2 ⁺	E2 ^c			
2056.07	(3) ⁺	288.3 [#] 5	8.3 [#] 21	1767.69	4 ⁺				I _γ : others: 11.3 14 in (⁶ Li,3nγ); 57 in (p,2nγ).
		602.3 1	100 [#] 5	1453.93	2 ⁺	M1+E2 ^c	+0.24 8		
		1262.6 2	28 [#] 3	793.22	2 ⁺	D+Q ^c			
2071.6	0 ⁺	617 [@]		1453.93	2 ⁺				
		1279 [@]		793.22	2 ⁺				
2297.93		844.0 1	100	1453.93	2 ⁺				
2448.11	3 ⁻	994.4 4	100 [#] 10	1453.93	2 ⁺	D ^c			A 679γ with an intensity 3 times that of 994γ is reported only in (α,2nγ).
		1654.6 [#] 2	63 [#] 5	793.22	2 ⁺				
2598.23	(4 ⁺)	1144.3 2	100 9	1453.93	2 ⁺				
		1805.0 ^{#e} 10	5 5	793.22	2 ⁺				
2735.25	(5 ⁺)	680.6 [#] 4	100 [#] 8	2056.07	(3) ⁺				E _γ : poor fit, level-energy difference=679.2. Additional information 3. Additional information 4.
		967.2 [#] 2	31 [#] 3	1767.69	4 ⁺	D+Q ^c			
2769.03	(5 ⁻)	321.0 1	2.8 ^{&} 5	2448.11	3 ⁻	[E2]		0.0153	B(E2)(W.u.)=22 5
		1001.28 7	100 ^{&} 9	1767.69	4 ⁺	(E1)			B(E1)(W.u.)=3.6×10 ⁻⁵ 5 Mult.: ΔJ=1, dipole from γ(θ); ΔJ ^π requires E1.
2807.87	6 ⁺	1040.11 9	100	1767.69	4 ⁺	E2			B(E2)(W.u.)=21 5
2886.99	2 ⁺	1119.6 [#] 2	100 [#] 10	1767.69	4 ⁺				
		2093.3 [#] 2	45 [#] 15	793.22	2 ⁺				
3041.25	(5 ⁻)	272.2 1	100 ^{&} 3	2769.03	(5 ⁻)	(D) ^c			
		593.3 2	27 ^{&} 3	2448.11	3 ⁻				
3098.67	6 ⁽⁺⁾	290.8 1	37 ^{&} 3	2807.87	6 ⁺				
		1331.0 2	100 ^{&} 6	1767.69	4 ⁺	Q			
3157.05	(7 ⁺)	421.8 1	100	2735.25	(5 ⁺)	Q			
3270.58	(4,5,6) ⁺	462.8 [#] 2	100 [#] 5	2807.87	6 ⁺	M1,E2			
		1502.8 [#] 2	62 [#] 6	1767.69	4 ⁺				I _γ : other: 30 10 in ⁷⁶ Ge(¹² C,4nγ), ⁸¹ Br(⁶ Li,3nγ).
3279.15	(6 ⁻)	237.9 1	17 ^{&} 2	3041.25	(5 ⁻)	D			
		510.1 [‡] 5	≈100 ^{‡&}	2769.03	(5 ⁻)				
3331.91	8 ⁺	524.0 1	100	2807.87	6 ⁺	E2 ^c			B(E2)(W.u.)=4.18 14

Adopted Levels, Gammas (continued)

$\gamma(^{84}\text{Si})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ [†]	I_γ [†]	E_f	J_f^π	Mult. ^a	Comments
3487.92	(7 ⁻)	680.0 2	21.1 21	2807.87	6 ⁺	(E1) ^c	B(E1)(W.u.)=4.4×10 ⁻⁵ 7
		718.9 1	100 5	2769.03	(5 ⁻)	E2 ^c	Mult.: ΔJ=1, dipole for a doublet (680.0+679.1); E1 from ΔJ ^π .
3511.77	(4 ⁺ ,5 ⁻)	241.2 [#] 5	5.1 [#] 34	3270.58	(4,5,6) ⁺		B(E2)(W.u.)=25 4
		703.6 [#] 2	100 [#] 10	2807.87	6 ⁺		
		1063.5 [#] 3	13 [#] 4	2448.11	3 ⁻		
		1744.4 [#] 2	38 [#] 4	1767.69	4 ⁺		
3578.23?		980.2 ^{#e} 10	82 [#] 45	2598.23	(4 ⁺)		
		1129.6 ^{#e} 4	36 [#] 18	2448.11	3 ⁻		
		1810.8 ^{#e} 3	100 [#] 45	1767.69	4 ⁺		
3650.15	(7 ⁻)	162.2 ^{&} 2	91 ^{&} 4	3487.92	(7 ⁻)		
		371.0 ^{&} 1	22 ^{&} 4	3279.15	(6 ⁻)	D ^c	
		551.5 ^{&} 2	39 ^{&} 4	3098.67	6 ⁽⁺⁾	D ^c	
		608.9 ^{&} 1	100 ^{&} 4	3041.25	(5 ⁻)		
		881.1 ^{&} 2	52 ^{&} 4	2769.03	(5 ⁻)	(Q) ^c	
3679.94	8 ⁺	348.0 1	29.2 9	3331.91	8 ⁺	(M1+E2) ^c	
		581.3 2	4.4 9	3098.67	6 ⁽⁺⁾		
		872.1 1	100 3	2807.87	6 ⁺	E2 ^c	B(E2)(W.u.)=11.5 7
3749.07	(7)	650.4 2	100	3098.67	6 ⁽⁺⁾	D+Q ^c	
3819.58?		932.2 ^{#e} 2	60 [#] 5	2886.99	2 ⁺		
		1370.8 ^{#e} 3	21 [#] 11	2448.11	3 ⁻		
		1763.6 ^{#e} 2	100 [#] 11	2056.07	(3) ⁺		
		2052.9 ^{#e} 3	26 [#] 13	1767.69	4 ⁺		E _γ : poor fit, level-energy difference=2051.9.
3918.08?		1110.3 ^{#e} 2	100 [#] 10	2807.87	6 ⁺		
		1469.9 ^{#e} 2	29 [#] 10	2448.11	3 ⁻		
		2150.9 ^{#e} 5	17 [#] 8	1767.69	4 ⁺		
4028.78	(8 ⁺)	1220.9 2	100	2807.87	6 ⁺	Q	
4062.78	4 ⁺	1255.0 [#] 2	100 [#] 10	2807.87	6 ⁺		
		1463.3 ^{#e} 2	6 [#] 3	2598.23	(4 ⁺)		
		1614.5 [#] 2	27 [#] 3	2448.11	3 ⁻		
		2006.7 ^{#e} 5	4.5 [#] 30	2056.07	(3) ⁺		
		2295.3 [#] 4	33 [#] 5	1767.69	4 ⁺		
4268.05	(8 ⁻)	780.1 ^{&} 2	48 ^{&} 4	3487.92	(7 ⁻)		
		988.9 ^{&} 1	100 ^{&} 4	3279.15	(6 ⁻)	Q ^c	

Adopted Levels, Gammas (continued)

$\gamma(^{84}\text{Sr})$ (continued)

$E_i(\text{level})$	J_i^π	E_γ [†]	I_γ [†]	E_f	J_f^π	Mult. ^a	α^d	Comments
4365.95	(4 ⁺)	1479.2 [#] 2	39 [#] 13	2886.99	2 ⁺			
		1557.6 [#] 3	26 [#] 13	2807.87	6 ⁺			
		1918.0 [#] 4	100 [#] 13	2448.11	3 ⁻			
		2309.5 [#] 4	52 [#] 9	2056.07	(3) ⁺			
4370.4	(9 ⁺)	1213.3 2	100	3157.05	(7 ⁺)	(Q) ^c		
4447.61	10 ⁺	1115.7 1	100	3331.91	8 ⁺	E2 ^c		B(E2)(W.u.)=6.7 11
4534.06	10 ⁺	86.3 2	14 5	4447.61	10 ⁺	[M1+E2]	1.0 8	E _γ , I _γ : from (²⁸ Si,3pγ).
		854.1 1	100 5	3679.94	8 ⁺	E2 ^c		B(E2)(W.u.)=27 5
								I _γ : from (²⁸ Si,3pγ).
								B(E2)(W.u.)=5.2 9
4636.13	(9 ⁻)	1148.2 1	100	3487.92	(7 ⁻)	E2 ^c		
4745.72	(8,9,10 ⁺)	1413.8 2	100	3331.91	8 ⁺			
5150.7?		1232.9 ^{#e} 3	38 [#] 3	3918.08?				
		1330.7 ^{#e} 4	100 [#] 10	3819.58?				
		1638.6 ^{#e} 7	12 [#] 9	3511.77	(4 ⁺ ,5 ⁻)			
5444.48	(11 ⁻)	808.35 10	100 4	4636.13	(9 ⁻)	E2 ^c		B(E2)(W.u.)=7.7 12
		996.9 1	30 4	4447.61	10 ⁺			
5653.25	12 ⁺	1119.2 1	100 4	4534.06	10 ⁺	E2 ^c		B(E2)(W.u.)=18 7
		1205.6 2	35 4	4447.61	10 ⁺	E2 ^c		B(E2)(W.u.)=4.3 16
5891.6	(12 ⁺)	1444 1	100	4447.61	10 ⁺	[E2]		B(E2)(W.u.)=17 8
6069.43	(12 ⁻)	625.0 1	100	5444.48	(11 ⁻)	(M1) ^c	0.00157	B(M1)(W.u.)=0.21 8
								Mult.: ΔJ=1, dipole from γ(θ), ΔJ ^π requires M1.
6484.34	(13 ⁻)	415.1 2	100 13	6069.43	(12 ⁻)	[M1]	0.00407	B(M1)(W.u.)=0.44 22
		830.9 2	<12	5653.25	12 ⁺	[E1]		B(E1)(W.u.)=5.E-5 +6-5
		1040 1	<12	5444.48	(11 ⁻)	[E2]		B(E2)(W.u.)=1.8 +21-18
6739.65	14 ⁺	1086.4 1	100	5653.25	12 ⁺	E2 ^c		B(E2)(W.u.)=41 14
6916.8	(14 ⁻)	432.5 3	100	6484.34	(13 ⁻)			
7822.8	(15 ⁺)	1084		6739.65	14 ⁺			
8006.4	16 ⁺	184		7822.8	(15 ⁺)	[M1]	0.0314	E _γ : from (³⁶ S,p2nγ) only.
		1266.5 5	100 17	6739.65	14 ⁺	[E2]		B(E2)(W.u.)<54
9098.4	(17 ⁺)	1092		8006.4	16 ⁺			
		1276		7822.8	(15 ⁺)			
9424.9	18 ⁺	327		9098.4	(17 ⁺)			E _γ : from (³⁶ S,p2nγ) only.
		1418 1	100 25	8006.4	16 ⁺	[E2]		B(E2)(W.u.)<50
11059.9	20 ⁺	1635 2	100	9424.9	18 ⁺	[E2]		B(E2)(W.u.)>12
12920	22 ⁺	1860 2	100	11059.9	20 ⁺			
15080?	(24 ⁺)	2160 ^e	100	12920	22 ⁺			E _γ : A 2125γ is tentatively assigned in (³⁶ S,p2nγ) from a 24 ⁺ to 22 ⁺ .

Adopted Levels, Gammas (continued) $\gamma(^{84}\text{Sr})$ (continued)


- [†] From weighted averages of all available data. Energies from ($\alpha, 2n\gamma$) have not been used in the averaging procedure due to consistently low values.
- [‡] Doublet. Approximate intensity given.
- [#] From ^{84}Y ε decay (39.5 min).
- [@] From ^{84}Y ε decay (4.6 s).
- [&] From [1982De05](#) in $^{81}\text{Br}(^6\text{Li}, 3n\gamma)$ reaction.
- ^a From ce data in ^{84}Y ε decay (39.5 min) unless otherwise stated.
- ^b From $\gamma\gamma(\theta)$ in ^{84}Y ε decay (39.5 min), unless otherwise stated.
- ^c From $\gamma(\theta)$ data in in-beam γ -ray studies. From RUL, $\Delta J=2$, quadrupole transitions are assigned as E2.
- ^d Total theoretical internal conversion coefficients, calculated using the BrIcc code ([2008Ki07](#)) with Frozen orbital approximation based on γ -ray energies, assigned multipolarities, and mixing ratios, unless otherwise specified.
- ^e Placement of transition in the level scheme is uncertain.

Adopted Levels, Gammas

Legend

Level Scheme

Intensities: Relative photon branching from each level


-----► γ Decay (Uncertain)

Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level


-----► γ Decay (Uncertain)


Adopted Levels, Gammas

Legend

Level Scheme (continued)

Intensities: Relative photon branching from each level

-----► γ Decay (Uncertain) $^{84}_{38}\text{Sr}_{46}$

Adopted Levels, Gammas