

${}^9\text{Be}({}^{13}\text{O},\text{X}): {}^{11}\text{O}$ 2p decay [2019We03](#)

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	J. H. Kelley, B. Grees		ENSDF	31-July-2020

[2019We03](#): ${}^9\text{C}$ is populated in the 2p-decay of ${}^{11}\text{O}$. A beam of 69.5 MeV/nucleon ${}^{13}\text{O}$ ions, from the NSCL/A1900 fragment separator, was purified in the Radio Frequency Fragment Separator before impinging on a 1-mm thick ${}^9\text{Be}$ target. The reaction products were detected using the HiRA High-Resolution position sensitive $\Delta\text{E-E}$ telescope array, which covered the polar angles $\theta_{\text{lab}}=2.1^\circ$ to 12.1° . A broad peak near $E_{\text{res}}(2\text{p}+{}^9\text{C})\approx 4.5$ MeV was observed in the total energy spectrum and attributed to a collection of four 2p-unbound ${}^{11}\text{O}$ states.

See additional discussion and theoretical analysis in ([2019Fo10](#), [2019Ka50](#), [2019Wa16](#)).

 ${}^9\text{C}$ LevelsE(level)

0