

$^{176}\text{Yb}(^{28}\text{Si},\text{F}\gamma)$ 2005Pa48

Type	Author	History	Citation	Literature Cutoff Date
Full Evaluation	D. Abriola(a), A. A. Sonzogni		NDS 107, 2423 (2006)	1-Jan-2006

Includes $^{176}\text{Yb}(^{31}\text{P},\text{X}\gamma)$ also from 2005Pa48.

$^{176}\text{Yb}(^{28}\text{Si},\text{x}\gamma)$: E=145 MeV. Measured $E\gamma$, $I\gamma$, $\gamma\gamma$ with the EUROGAM II array, consisting of 54 escape-suppressed Ge detectors, 30 of which were of large-volume coaxial design with the remaining 24 being of the four-element clover type.

$^{176}\text{Yb}(^{31}\text{P},\text{X}\gamma)$: E=152 MeV. Measured $E\gamma$, $I\gamma$, $\gamma\gamma$ with the EUROBALL IV array, comprising 15 cluster, 26 clover and 30 tapered single-crystal Ge detectors. Each cluster detector consisted of seven closely-packed, large-volume encapsulated Ge crystals.

 ^{94}Zr Levels

$E(\text{level})^\dagger$	$J^\pi^\ddagger$	$E(\text{level})^\dagger$	$J^\pi^\ddagger$	$E(\text{level})^\dagger$	$J^\pi^\ddagger$	$E(\text{level})^\dagger$	$J^\pi^\ddagger$
0.0	0^+	3143.0 5	(6^+)	4371.4 6		6009.5 7	(13^+)
919.0 3	2^+	3444.1 6	(7^-)	4480.5 6	(10^+)	6374.8 8	(14)
1470.0 4	4^+	3596.4 6		4813.8 7	(12^+)	7058.2 8	(15)
2330.2 4	4^+	3632.7 6	(8^+)	5492.9 7	(11^+)	7795.0 9	(16)
2606.2 5	5^-	4225.7 6		5806.9 7	(12^+)		

† From least-squares fit to $E\gamma$'s; $\Delta E\gamma=0.3$ keV assumed for each transition.

‡ Tentative assignments based on well known low-spin states from literature, comparison with theoretical calculations and by assuming spin values increase with excitation energy of levels. Tentative assignments are supported by comparison of corresponding states in neighboring Zr isotopes using a weak-coupling scheme.

 $\gamma(^{94}\text{Zr})$

$E_\gamma^\dagger$	I_γ	$E_i(\text{level})$	J_i^π	E_f	J_f^π	$E_\gamma^\dagger$	I_γ	$E_i(\text{level})$	J_i^π	E_f	J_f^π
145.7	1.9 2	4371.4		4225.7		781.6	1.6 3	4225.7		3444.1	(7^-)
152.3	4.7 3	3596.4		3444.1	(7^-)	812.8	46.4 16	3143.0	(6^+)	2330.2	4^+
202.6	5.9 3	6009.5	(13^+)	5806.9	(12^+)	837.9	24.9 12	3444.1	(7^-)	2606.2	5^-
314.0	9.0 4	5806.9	(12^+)	5492.9	(11^+)	847.8	37.5 15	4480.5	(10^+)	3632.7	(8^+)
333.3	10.5 5	4813.8	(12^+)	4480.5	(10^+)	919.0	100	919.0	2^+	0.0	0^+
365.3	15.3 6	6374.8	(14)	6009.5	(13^+)	927.2	5.9 5	4371.4		3444.1	(7^-)
489.7	49.6 14	3632.7	(8^+)	3143.0	(6^+)	1012.4	12.7 8	5492.9	(11^+)	4480.5	(10^+)
536.8	7.5 5	3143.0	(6^+)	2606.2	5^-	1136.2	28.6 13	2606.2	5^-	1470.0	4^+
551.0	46.8 18	1470.0	4^+	919.0	2^+	1195.7	8.4 7	6009.5	(13^+)	4813.8	(12^+)
629.3	2.3 3	4225.7		3596.4		1411.2	48.9 23	2330.2	4^+	919.0	2^+
683.4	5.1 5	7058.2	(15)	6374.8	(14)	1672.9	1.9 3	3143.0	(6^+)	1470.0	4^+
736.8	2.6 4	7795.0	(16)	7058.2	(15)						

† Gamma energies from this dataset are consistently higher than those from other datasets.


$^{176}\text{Yb}(^{28}\text{Si},\text{F}\gamma)$ 2005Pa48

Level Scheme

Intensities: Relative I_γ

Legend

- $\longrightarrow$ $I_\gamma < 2\% \times I_\gamma^{\text{max}}$
- $\longrightarrow$ $I_\gamma < 10\% \times I_\gamma^{\text{max}}$
- $\longrightarrow$ $I_\gamma > 10\% \times I_\gamma^{\text{max}}$

 $^{94}_{40}\text{Zr}_{54}$