

New nuclear data libraries for Pb and Bi

Arjan Koning

Marieke Duijvestijn

Robin Klein Meulekamp

Steven van der Marck

Alfred Hogenbirk

NRG Petten

CSEWG Meeting, Brookhaven National Laboratory, November 4-6, 2003

Contents

- Nuclear data needs for Pb and Bi
- TALYS code
- TALYS vs. experiments
- New ENDF-6 formatted data libraries
- Integral benchmarks
- Conclusions

Nuclear data problems in lead cooled fast systems

Various independent checks of cross section sensitivity for lead/bismuth cooled ADS:

- IAEA ADS benchmark: reactivity burnup swing
- PDS-XADS
- MUSE
- Other ADS designs (CIEMAT, ANL (Palmiotti, Salvatores))

Result: (n,n'), (n,2n) and feeding from > 20 MeV yield considerable differences, among world data libraries, in source importance ϕ^* and up to 2000 pcm in k_{eff} , *on the basis of Pb cross sections only*.

Uncertainties in lead cross sections

Is this uncertainty really necessary?

Much experimental data from past 10 years not taken into account in data libraries.

Attacking the lead problem:

- Collect all relevant experimental data
- Compare nuclear model code with experiments and adjust nuclear model parameters, if needed.
- Produce all reaction channels with nuclear model code → complete ENDF-6 datafile
- Do not stop at the usual 20 MeV but at 200 MeV. (HINDAS-project)

Why do we need a model code?

Low energies ($E < 1$ MeV):

- Only a few open reaction channels: Elastic scattering, capture and fission.
- For important nuclei: (almost) Complete measurements can be made.

Higher energies ($E > 1$ MeV):

- Many reaction channels open up and the existing experimental database can never cover what happens in reality: *Predictive nuclear models are essential*.
- A model code can reveal inconsistencies between uncorrelated experiments.

Important for nuclear applications:


- Number of (expensive) experiments is very finite. A model code can produce nuclear data for *all* nuclides, energies and channels → Required accuracy from applications is crucial information to decide on use of experiment or model code or both!

TALYS

TALYS: nuclear reaction software by NRG Petten (Koning, Duijvestijn) and CEA Bruyères-le-Châtel (Hilaire)

- Energy range 1 keV - 200 MeV.
- Neutrons, protons, deuterons, tritons, helions, alphas and photons.
- Many nuclear reaction models implemented.
- Continuous, smooth description over a wide energy and mass range.
- Automatic generation of data in ENDF-6 format.
- TALYS is not yet generally available.

TALYS: CALCULATIONAL SCHEME







Data produced by TALYS

- Total, elastic and reaction cross sections.
- Inelastic cross sections.
- Elastic and inelastic angular distributions.
- All exclusive reaction channels (e.g. (n,p), (n,2n), (n,nd2a)).
- All exclusive double-differential spectra.
- All exclusive isomeric production cross sections.
- All exclusive discrete and continuum gamma ray production cross sections.
- Photonuclear reactions.
- Reactions on isomeric targets.

Data produced by TALYS


- Total particle production cross sections (n,xn).
- Total particle production double-differential spectra.
- Residual production cross sections (including isomers).
- Activation libraries in EAF and ENDF-6 format.
- Transport libraries in ENDF-6 format.






$^{208}\text{Pb}(n,n_1')$

$J^\pi = 3^-, E_x = 2.61 \text{ MeV}$




$^{208}\text{Pb}(n,n'\gamma)$

Level 1 $\rightarrow$ Level 0: $E_\gamma = 2.61$ MeV




$^{208}\text{Pb}(n,n_2')$

$J^\pi = 5^-, E_x = 3.20 \text{ MeV}$




$^{208}\text{Pb}(n,n')$

Total




$^{208}\text{Pb}(n, n)$




Pb(n,xn) at 14.23 MeV

Angle-integrated spectrum




$^{208}\text{Pb}(n,2n)^{207}\text{Pb}$




$^{208}\text{Pb}(n,2n\gamma)$

Level 1 $\rightarrow$ Level 0: $E_\gamma=0.57$ MeV




$^{208}\text{Pb}(n,2n\gamma)$

Level 2 $\rightarrow$ Level 0: $E_\gamma = 0.89 \text{ MeV}$




$^{208}\text{Pb}(n,p)^{208}\text{Tl}$




$^{207}\text{Pb}(n,n_1')$

$J^\pi = 2.5^-, E_x = 0.57 \text{ MeV}$




$^{207}\text{Pb}(n,n')$

$J^\pi = 1.5^-, E_x = 0.90 \text{ MeV}$




$^{206}\text{Pb}(n,\alpha)$




$^{209}\text{Bi}(n,n')$


$J^\Pi = 3.5^-, E_x = 0.90$




$^{209}\text{Bi}(n,2n)^{208}\text{Bi}$




$^{209}\text{Bi}(n,3n)^{207}\text{Bi}$




$^{209}\text{Bi}(n,4n)^{206}\text{Bi}$




$^{209}\text{Bi}(n,5n)^{205}\text{Bi}$




$^{209}\text{Bi}(n,6n)^{204}\text{Bi}$




$^{209}\text{Bi}(n,7n)^{203}\text{Bi}$




$^{209}\text{Bi}(n,8n)^{202}\text{Bi}$




$^{209}\text{Bi}(n,9n)^{201}\text{Bi}$




$^{209}\text{Bi}(n,10n)^{200}\text{Bi}$



$^{209}\text{Bi}(n,11n)^{199}\text{Bi}$



$^{209}\text{Bi}(n,12n)^{198}\text{Bi}$



	Library	Resonance range	URR - 20 MeV	> 20 MeV
Pb-204	ENDF/B-VI.8	—	—	—
	JENDL-3.3	Mizumoto (1987)	Mizumoto(1987), Shibata(2001)	—
	JEFF-3.0	Mizumoto (1987)	Mizumoto(1987)	—
	BROND-2.2	Blokhin (1984)	Blokhin(1990)	—
	NRG	Larson (1989)	Koning(2003)	Koning(2003)

	Library	Resonance range	URR - 20 MeV	> 20 MeV
Pb-206	ENDF/B-VI.8	Larson (1989)	Fu (1989)	Chadwick (1996)
	JENDL-3.3	Mizumoto (1987)	Mizumoto(1987), Shibata(2001)	—
	JEFF-3.0	Mizumoto (1987)	Mizumoto(1987)	—
	BROND-2.2	Blokhin (1984)	Blokhin(1990)	—
	NRG	Larson (1989)	Koning(2003)	Koning(2003)


	Library	Resonance range	URR - 20 MeV	> 20 MeV
Pb-207	ENDF/B-VI.8	Larson (1989)	Fu (1989)	Chadwick (1996)
	JENDL-3.3	Mizumoto (1987)	Mizumoto(1987), Shibata(2001)	—
	JEFF-3.0	Mizumoto (1987)	Mizumoto(1987)	—
	BROND-2.2	Blokhin (1984)	Blokhin(1990)	—
	NRG	Larson (1989)	Koning(2003)	Koning(2003)

	Library	Resonance range	URR - 20 MeV	> 20 MeV
Pb-208	ENDF/B-VI.8	Larson (1989)	Young (1996)	Chadwick (1996)
	JENDL-3.3	Larson (1989)	Mizumoto(1987), Shibata(2001)	—
	JEFF-3.0	Larson (1989)	Mizumoto(1987)	—
	BROND-2.2	Blokhin (1984)	Blokhin(1990)	—
	NRG	Larson (1989)	Koning(2003)	Koning(2003)

	Library	Resonance range	URR - 20 MeV	> 20 MeV
Bi-209	ENDF/B-VI.8	Smith (1989)	Smith (1989)	Chadwick (1998)
	JENDL-3.3	Yamamuro (1989)	Yamamuro(1989), Shibata(2001)	—
	JEFF-3.0	Smith (1989)	Smith (1989)	—
	BROND-2.2	Blokhin (1984)	Blokhin(1990)	—
	NRG	Yamamuro (1989)	Koning(2003)	Koning(2003)

ENDF-6 philosophy used by TALYS


- Data file should be complete, i.e. should include all required transport and activation data.
- Use one and the same file structure for every isotope:
 - MF1
 - MF2: use best possible resonance data from other library.
 - MF3: include all open reaction channels. Use MT5 for $(n, \gamma n)$ and reaction cross section above 20 MeV.
 - MF4: Use only for elastic scattering angular distribution (MT2).




ENDF-6 philosophy used by TALYS

- MF6: Use for all discrete angular distributions (apart from elastic), all double-differential exclusive spectra, all discrete and continuum photon production and recoils. Above 20 MeV, use MF6/MT5 for particle production cross sections and spectra and residual production cross sections.
- MF8: In case of isomers: point to MF (6, 9 or 10) where info exists.
- MF9: Isomeric cross sections for non-threshold reactions.
- MF10: Isomeric cross sections for threshold reactions.

Benchmark leu-comp-therm-010




Impact of new Pb data in fast spectrum





Pb(n,xn) at 14.23 MeV


Angle-integrated spectrum




TU-Dresden (Seidel) benchmark: leakage through Pb-sphere (2.5 - 25cm)







Neutron spectrum outside lead buffer for incident protons of 150 MeV



Conclusions

New ENDF-6 formatted data libraries for $^{204,206,207,208}\text{Pb}$ and ^{209}Bi

- Older Pb and Bi evaluations from the major world libraries (ENDF/B-VI, JENDL and JEFF) all suffer from problems below 20 MeV, at the microscopic and benchmark level.
- Existing experimental data < 20 MeV consistently analyzed with TALYS code.
- New experiments/model developments used for $20 < E < 200$ MeV.
- Evaluated datafiles improved from unresolved resonance range up to 200 MeV with TALYS:
 - Consistent good description of many reaction channels.
 - Complete ENDF-6 datafile including all secondary distributions, photons, etc.
 - Successful NJOY processing.

Conclusions

- Data files will be proposed for the next version of JEFF.
- First integral benchmarks with MCNP(X) encouraging.
- Sensitivity studies and further integral benchmarks should reveal whether good is good enough.
- TALYS is ready to give other nuclides the same treatment (All Ca, Sc, Fe, and Ge isotopes under construction).