NEW APPROACH FOR THE DETERMINATION OF THE NUCLEAR LEVEL DENSITY PARAMETERS

Said F. Mughabghab
Brookhaven National Laboratory
Building 197D
Upton, NY 11973
(516) 344-5085

Charles L. Dunford
Brookhaven National Laboratory
Building 197D
Upton, NY 11973
(516) 344-2804

ABSTRACT

The nuclear level density parameters of the stable nuclides have been determined from the neutron resonance data on the basis of new values for the spin dispersion parameter. The latter parameter is deduced from the spin dependent average level spacings of s-wave neutron resonances of odd target nuclides. In addition, the temperature dependence of the level density parameter is studied and an analytical relation for it is presented.

I. INTRODUCTION

The nuclear level density plays a major role in statistical calculations in reactor physics, nuclear model calculations, astrophysics, intermediate-energy heavy-ion collision investigations, and spallation neutron calculations. However, up until very recently[1], little attention was paid into a critical examination of the derivation of the level density parameter. This is largely due to a lack of a detailed knowledge of the spin dispersion parameter. Previous analysis [2,3,4] relied on two theoretical expressions for this parameter, one due to Gilbert and Cameron, [5] the other to Bohr and Mottelson [6] . The validity of the theoretical values of the spin dispersion parameter were not verified previously, particularly over an extended atomic mass range.

II. ANALYSIS AND RESULTS

The spin dispersion parameter was determined from the ratio of the average level spacings of s-wave neutron resonances[7, 8] for the two spin states I + 0.5, and I - 0.5 , where I is the spin of t

$$\rho(U,J) = \frac{(2J+1)}{24U^{5/4}2^{1/2}\sigma^3} e^{2(aU)^{1/2}} \, e^{\frac{-(J+1/2)^2}{2\sigma^2}} \qquad \textbf{(1)}$$

$$\sigma^2 = 0.0494 aTA^{2/3} \qquad \textbf{(2)}$$